

Evidence Synthesis

Number 205

Aspirin Use to Prevent Preeclampsia and Related Morbidity and Mortality: An Evidence Update for the U.S. Preventive Services Task Force

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
5600 Fishers Lane
Rockville, MD 20857
www.ahrq.gov

Contract No. HHS-290-2015-00007-I-EPC5, Task Order No. 6

Prepared by:

Kaiser Permanente Research Affiliates Evidence-based Practice Center
Kaiser Permanente Center for Health Research
Portland, OR

Investigators:

Jillian T. Henderson, PhD, MPH
Kimberly K. Vesco, MD, MPH
Caitlyn A. Senger, MPH
Rachel G. Thomas, MPH
Nadia Redmond, MSPH

AHRQ Publication No. 21-05274-EF-1
February 2021

This report is based on research conducted by the Kaiser Permanente Research Affiliates Evidence-based Practice Center (EPC) under contract to the Agency for Healthcare Research and Quality (AHRQ), Rockville, MD (Contract No. HHS-290-2015-00007-I-EPC5, Task Order No. 6). The findings and conclusions in this document are those of the authors, who are responsible for its contents; the findings and conclusions do not necessarily represent the views of AHRQ. Therefore, no statement in this report should be construed as an official position of AHRQ or of the U.S. Department of Health and Human Services.

The information in this report is intended to help health care decisionmakers—patients and clinicians, health system leaders, and policymakers, among others—make well-informed decisions and thereby improve the quality of healthcare services. This report is not intended to be a substitute for the application of clinical judgment. Anyone who makes decisions concerning the provision of clinical care should consider this report in the same way as any medical reference and in conjunction with all other pertinent information (i.e., in the context of available resources and circumstances presented by individual patients).

The final report may be used, in whole or in part, as the basis for development of clinical practice guidelines and other quality enhancement tools, or as a basis for reimbursement and coverage policies. AHRQ or U.S. Department of Health and Human Services endorsement of such derivative products may not be stated or implied.

Acknowledgments

The authors gratefully acknowledge the following individuals for their contributions to this project: Iris Mabry-Hernandez, MD, at AHRQ; current and former members of the U.S. Preventive Services Task Force who contributed to topic deliberations; the National Heart, Lung, and Blood Institute, the Office of Research on Women's Health, and the Centers for Disease Control and Prevention, for providing federal partner review of the draft report; Lisa Askie, PhD, MPH, BN, CM, NICC, Robert Silver, MD, Shireen Meher, MD, and James Roberts, MD, who provided expert review of the draft report; Jennifer Lin, MD, for mentoring and project oversight; Melinda Davies, MA, and Neon Brooks, PhD, for technical and editorial assistance at the Center for Health Research.

Structured Abstract

Objective: We conducted this review to support the United States Preventive Services Task Force (USPSTF) in updating its 2014 recommendation on daily low dose aspirin use during pregnancy for individuals at increased risk for preeclampsia. We systematically reviewed updated evidence on the effectiveness and potential harms of daily aspirin use in pregnancy to prevent morbidity and mortality associated with preeclampsia.

Data Sources: We performed comprehensive searches of MEDLINE, PubMed (publisher-supplied only), Embase, and the Cochrane Collaboration Registry of Controlled Trials for studies published between January 2013 and July 2019. A research librarian developed and executed the search strategy. Studies included in the prior review to support the 2014 recommendation and studies referenced in recently published reviews were also considered for inclusion.

Study Selection: We reviewed 3749 newly identified abstracts and 183 articles against prespecified inclusion criteria developed for the current review. Studies were included if they evaluated daily treatment during pregnancy with aspirin not combined with any other medication at a dosage of ≥ 50 mg daily compared with placebo or no treatment. Included studies reported incidence of preeclampsia or serious maternal, perinatal, and longer-term health outcomes. For studies evaluating aspirin effectiveness, eligible study designs included randomized controlled trials (RCTs) and individual participant data meta-analyses that enrolled pregnant persons who were at increased risk for preeclampsia based on personal sociodemographic characteristics, medical history, diagnostic measurements or assays, or risk prediction models. For studies evaluating harms of aspirin use, the study design criteria were expanded to include observational studies and trials conducted in average-risk populations.

Data Analysis: Eligible studies that were determined to have high risk of bias were excluded for poor quality according to standard USPSTF procedures. Descriptions of study populations, study design characteristics, and outcomes were developed for all included studies. The most consistently reported health outcomes were analyzed using random effects meta-analytic model to calculate pooled intervention effects and subgroup comparisons were tested with meta-regression tests for interaction. We examined whether the results were influenced by small-study effects using the Peters test and visual inspection of forest plots.

Results: Twenty-three studies (33 articles) met our inclusion criteria. Four studies included in the previous review (including two observational studies) were not included and four new trials were added. We included 18 trials evaluating aspirin effectiveness in individuals at increased risk for preeclampsia; all but one used a matching placebo comparator. Dosages of aspirin ranged from 50-150 mg per day, usually starting in the second or third trimester and continuing until delivery or near term. These interventions were associated with reduced risks of perinatal mortality (pooled RR 0.79 [95% CI 0.66, 0.96], I^2 0%), preterm birth (pooled RR 0.80 [95% CI 0.67, 0.95], I^2 49%), and small for gestational age/intrauterine growth restriction (SGA/IUGR) (pooled RR 0.82 [95% CI 0.68, 0.99], I^2 41%). There was also a statistically significant reduction in the risk of preeclampsia (pooled RR, 0.85 [95% CI, 0.75 to 0.95], I^2 0%). Rare maternal health outcomes, such as eclampsia and maternal mortality, occurred too infrequently to estimate preventive effectiveness. In meta-analytic subgroup comparisons, we found no consistent

evidence for effect differences related to intervention or population characteristics such as the timing of treatment initiation, the dosage of aspirin used, or participant characteristics. There was evidence of small-study effects for several of the pooled health outcomes.

Studies in average and increased-risk populations did not provide any clear evidence of harms associated with daily aspirin use (<150mg) taken during the second or third trimester of pregnancy, and there was no evidence of differences in harms by aspirin dosage or timing, or for specific populations identified in limited subgroup comparisons. Bleeding related harms were uncommon; our analyses showed null effects for differences in risk of postpartum hemorrhage (pooled RR 1.03, [95% CI, 0.94, 1.12], I^2 0%, k=9), or intracranial fetal bleeding (pooled RR, 0.90 [95% CI, 0.51, 1.57]; I^2 19%, k=6) between intervention and control groups. There was also no difference in rates of placental abruption (pooled RR, 1.15 [95% CI, 0.76, 1.72], I^2 25%, k=10). Longer-term followup from one large trial found no differences in child developmental outcomes between aspirin and placebo exposed groups. No differences were found in a limited set of studies reporting other rare perinatal harms.

Limitations: Our search was limited to English-language literature and trials conducted in settings other than very high Human Development Index settings were excluded. Large trials conducted in other settings could provide additional relevant information, but evidence from other reviews without this exclusion do not find substantively different results. The meta-analysis results for some outcomes were limited by low numbers of included studies and very few events. Conservative approaches to estimation of pooled effects and testing of subgroup comparisons were used but cannot fully address some limitations in the data. The two largest studies used the same low aspirin dosage (60mg) and enrolled a majority of participants after 16 weeks of gestation and small-study effects cannot be entirely disentangled from other design and population features of the included studies. Evidence was lacking on potential longer term health consequences of low dose aspirin exposure during pregnancy.

Conclusions: Daily aspirin use in pregnancy for individuals at increased risk for preeclampsia consistently led to beneficial effects on perinatal mortality, preterm birth, fetal growth restriction, and preeclampsia diagnosis across a clinically heterogeneous set of trials. A large body of trial evidence shows no clear signal of serious harms associated with daily low dose aspirin use in the second and third trimesters of pregnancy. Further research from pragmatic and comparative effectiveness trials is needed to identify the best way to identify patients at increased preeclampsia risk who are most likely to benefit from aspirin use, and to obtain greater clarity on the optimal regimen.

Table of Contents

Chapter 1. Introduction.....	1
Condition Background.....	1
Condition Definition.....	1
Prevalence and Burden.....	2
Etiology and Natural History.....	3
Risk Factors.....	4
Treatment Approaches.....	5
Preeclampsia Prevention.....	5
Previous USPSTF Recommendation and Recommendations of Others.....	6
Chapter 2. Methods.....	7
Scope and Purpose.....	7
Key Questions and Analytic Framework.....	7
Data Sources and Searches.....	7
Study Selection.....	8
Quality Assessment and Data Abstraction.....	8
Data Synthesis and Analysis.....	9
Grading the Strength of the Body of Evidence.....	10
Expert Review and Public Comment.....	11
USPSTF Involvement.....	11
Chapter 3. Results.....	12
Description of Included Studies.....	12
Population and Setting Details.....	12
Intervention Characteristics.....	14
KQ1. Does Aspirin Reduce Adverse Maternal, Perinatal, Child, or Combined Health Outcomes in Pregnant Persons at Increased Risk of Preeclampsia?.....	14
Summary of Results.....	14
Detailed Results.....	15
Small-Study Effects.....	16
KQ1a. Does the Effectiveness of Aspirin for Reducing Adverse Health Outcomes Vary by Subpopulations Defined by Personal Characteristics or Preeclampsia Risk Factors?.....	16
Aspirin Timing and Dosage.....	17
Personal Characteristics and Clinical History.....	17
KQ2. Does Aspirin Prevent Preeclampsia in Pregnant Persons at Increased Risk for Preeclampsia?.....	18
Summary of Results.....	18
Detailed Results.....	18
KQ2a. Does the Effectiveness of Aspirin for Reducing Preeclampsia Vary by Subpopulations Defined by Personal Characteristics or Preeclampsia Risk Factors?.....	19
Aspirin Timing and Dosage.....	19
Personal Characteristics and Clinical History.....	20
Preeclampsia Risk Assessment Approach.....	20
KQ3. What Are the Harms of Aspirin Use to Prevent Preeclampsia During Pregnancy?.....	21
Summary of Results.....	21
Detailed Results.....	21

KQ3a. Do the Harms of Aspirin Use to Prevent Preeclampsia Vary by Subpopulations Defined by Personal Characteristics or Preeclampsia Risk Factors?.....	23
Chapter 4. Discussion	25
Summary of Evidence.....	25
Comparison of Our Results With Findings From Other Systematic Reviews	27
Limitations	28
Exclusions for Publication Language and Study Setting.....	28
Small-Study Effects	29
Confounding of Aspirin Initiation Timing Effects	29
Observational Evidence Exclusions.....	30
Adherence	30
Identifying Patients at Increased Risk for Preeclampsia.....	31
Evidence Gaps and Future Research.....	34
Applicability of Evidence and Health Equity	35
Chronic and Gestational Hypertension	36
Obesity	38
Identifying Serious Harms	38
Longer-Term Health Outcomes	39
Conclusions.....	39
References.....	41

Figures

- Figure 1. Analytic Framework
- Figure 2. Perinatal Mortality, Sorted by Study Size
- Figure 3. Preterm Birth Before 37 Weeks of Gestation, Sorted by Study Size
- Figure 4. Small for Gestational Age or Intrauterine Growth Restriction, Sorted by Study Size
- Figure 5. Small for Gestational Age or Intrauterine Growth Restriction, Stratified by Timing of Aspirin Initiation
- Figure 6. Preterm Birth, Stratified by Timing of Aspirin Initiation
- Figure 7. Preeclampsia, Sorted by Study Size
- Figure 8. Preeclampsia, Stratified by Timing of Aspirin Initiation
- Figure 9. Placental Abruption, Stratified by Risk and Sorted by Study Size
- Figure 10. Postpartum Hemorrhage Stratified by Risk and Sorted by Study Size
- Figure 11. Fetal Intracranial Bleeding Stratified by Risk and Sorted by Study Size

Tables

- Table 1. Existing Clinical Recommendations
- Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population
- Table 3. Population, Study Design, and Intervention Characteristics, Average-Risk Population
- Table 4. Patient Risk Factors and Clinical Tests Used to Identify Study Populations at Increased Risk for Preeclampsia
- Table 5. Summary of Meta-Analysis Results
- Table 6. Summary of Evidence
- Table 7. Theoretical Number Needed to Treat (NNT) Across Different Levels of Outcome Incidence
- Table 8. Preeclampsia Risk Factors Based on Patient Medical History

Appendixes

Appendix A. Detailed Methods

Appendix B. Literature Flow Diagram

Appendix C. Included Studies

Appendix D. Excluded Studies

Appendix E. Detailed Tables

Appendix F. Evidence Tables

Appendix G. Ongoing Studies

Chapter 1. Introduction

Condition Background

Condition Definition

Preeclampsia is a pregnancy-specific disease that occurs after the 20th week of pregnancy and is defined as the new onset of hypertension (systolic blood pressure [SBP] \geq 140 mm Hg or diastolic blood pressure [DBP] \geq 90 mm Hg, on two occasions at least 4 hours apart) in a person with previously normal blood pressure, together with additional specified signs or symptoms.^{1,2} Although preeclampsia was historically diagnosed as hypertension accompanied by new onset proteinuria (300 mg/dL of protein in a 24-hour urine collection or a protein-to-creatinine ratio of 0.30 or more), in 2013 the American College of Obstetricians and Gynecologists (ACOG) revised the diagnostic criteria such that proteinuria is no longer requisite for diagnosis. In the absence of proteinuria, preeclampsia is diagnosed when hypertension is accompanied by any of the following signs or symptoms: thrombocytopenia; impaired liver function as indicated by elevated blood concentrations of liver enzymes and severe persistent right upper quadrant or epigastric pain unresponsive to medication; renal insufficiency; pulmonary edema; new-onset headache unresponsive to medication; or visual disturbances.¹ The presence of any of these systemic signs or symptoms, or of severe hypertension (systolic \geq 160mmHg or diastolic $>$ 110mmHg), also define the clinical designation of “preeclampsia with severe features.”¹

Gestational hypertension alone (i.e. new onset hypertension after 20 weeks of gestation that is not accompanied by the additional signs and symptoms that define preeclampsia), is also associated with adverse pregnancy outcomes and an increased risk for future chronic hypertension and cardiovascular disease.¹ Gestational hypertension is defined as new onset of elevated blood pressure (SBP of 140 mmHg or more or DBP of 90 mmHg or more, on two occasions at least 4 hours apart) after 20 weeks gestation in the absence of proteinuria or other signs or symptoms of preeclampsia.¹ Gestational hypertension requires enhanced surveillance as up to 50 percent of women develop proteinuria or other signs of preeclampsia, and progression is more likely if hypertension arises prior to 32 weeks of gestation.¹ Chronic hypertension with superimposed preeclampsia is diagnosed when there is new onset of proteinuria beyond 20 weeks in a woman with chronic hypertension but without prior proteinuria, or sudden onset of the following: substantial sustained increases in protein excretion, exacerbation of hypertension with need to escalate antihypertensive medication dose, or manifestation of systemic signs and symptoms associated with preeclampsia.

Preeclampsia can remain stable until delivery, but occasionally can quickly and unpredictably take a more serious turn. Severe hypertension; eclampsia; hemolysis, elevated liver enzymes, and low platelet counts (HELLP syndrome); and other organ or systemic complications can lead to maternal or fetal injury and death.^{2,3} There is some variation in the diagnostic criteria for HELLP syndrome, but the following are widely agreed upon: lactate dehydrogenase elevated to 600 IU/L or more, aspartate aminotransferase and alanine aminotransferase elevated more than twice the upper limit of normal, and platelet counts less than $100 \times 10^9/L$.^{1,4,9} The majority of HELLP

syndrome cases are seen in the third trimester, but about a third of cases present or progress postpartum. The primary symptoms of HELLP syndrome include right upper quadrant pain and generalized malaise, with nausea and vomiting seen in approximately half of cases.

Prevalence and Burden

Preeclampsia and eclampsia are the second and third most common causes of maternal morbidity and mortality worldwide, respectively.⁵⁻⁷ Access to standard medical care has been shown to greatly improve health outcomes, as demonstrated by higher proportions of maternal mortality attributed to hypertensive disorders of pregnancy in lower income countries.⁸

In the U.S., preeclampsia occurs in about 1 in 25 pregnancies that continue beyond 20 weeks of gestation. In a large retrospective cohort analysis of nearly 37 million deliveries (n=36,985,729) in the United States from 2006 to 2015, 3.8 percent received a diagnosis of preeclampsia. Severe preeclampsia increased as a proportion of preeclampsia cases over the study period, as did superimposed preeclampsia (preeclampsia arising in women with pre-existing chronic hypertension).⁹ A rising trend in the proportion of hospital deliveries in the US complicated by preeclampsia and/or preeclampsia is evident from Healthcare Cost and Utilization Project data on all hospital deliveries in the United States, a 21 percent increase from 38.4 per 1,000 deliveries in 2005 to 46.6 per 1,000 deliveries in 2014).¹⁰ Of over 175,000 deliveries with preeclampsia estimated in 2014 from these data, over one third (37%) were categorized as severe preeclampsia, 15 percent were cases among individuals with preexisting hypertension, and 1 percent resulted in eclampsia.¹⁰

In the United States, the risk of preeclampsia and of serious morbidity and mortality from the condition is greatest for non-Hispanic Black women.^{10, 11} Rates of preeclampsia are more than twice as high among Black women and more likely to manifest as severe disease. National estimates have shown that fewer than half of non-Hispanic White women had severe preeclampsia whereas nearly two-thirds of cases among Black women were severe.^{10, 12} Consequently, overall maternal mortality ratios and the contribution of preeclampsia/eclampsia to maternal mortality are higher for Black women.¹³ Disparities in overall health, chronic health conditions, health care access, psychosocial stress, and systemic racial biases in health care are thought to contribute to the greater risk and worse outcomes of preeclampsia for Black women.^{12, 14}

From 2011-2016, hypertensive disorders of pregnancy were responsible for 6.9 percent of pregnancy-related deaths in the United States.^{13, 15} Significant maternal morbidities associated with preeclampsia include cerebrovascular bleeding, retinal detachment, and organ damage and failure complications from HELLP syndrome.^{16, 17} Eclamptic seizures occur in approximately 1 to 2 percent of preeclampsia cases, and can lead to death or serious complications such as brain damage, aspiration pneumonia, pulmonary edema, placental abruption, disseminated coagulopathy, acute renal failure, cardiopulmonary arrest, and coma.⁵ Serious morbidity is more common than mortality; cohort data from obstetric patients attending 25 U.S.-based medical centers encompassing the Maternal-Fetal Medicine Units Network from 2008 to 2011 indicated that at least 21 percent of severe maternal morbidity (i.e., requiring surgical intervention, intubation, blood transfusion, or intensive care unit admission or diagnosed failure of at least one

organ system) was attributable to hypertensive disorders of pregnancy.¹⁸ The serious threat to maternal health associated with preeclampsia contributes to clinical decision-making related to the timing of delivery and other interventions to reduce maternal risk.

Preeclampsia increases the risk of adverse fetal, neonatal, and child health outcomes. These risks include intrauterine growth restriction, small for gestational age, low birth weight, preterm birth, placental abruption, stillbirth, and neonatal death.² The majority of preeclampsia cases occur after 34 weeks, but perinatal morbidity and mortality are greatest for early onset disease.¹⁹ Due to the fact that the only effective treatment for preeclampsia is delivery, the condition is a leading cause of medically induced preterm birth and low birth weight. It has been estimated that preeclampsia contributes to 6 percent of preterm births and 19 percent of medically indicated preterm births.²⁰ Infants born before term (<37 weeks of gestation) are at increased risk of morbidity and mortality, with risks rising dramatically with earlier delivery.

There is also growing evidence that having preeclampsia is associated with increased risk of maternal chronic hypertension and cardiovascular disease later in life, including congestive heart failure,²¹ myocardial infarction, and stroke.²² The risk appears to be greatest among women who have experienced preeclampsia in more than one pregnancy,²³ required preterm delivery, or had a pregnancy complicated by fetal growth restriction.²⁴ Other data suggest that having had any hypertensive disorder of pregnancy may also be associated with long-term cardiovascular health risk.^{25, 26} It is not yet clear whether this is due to common underlying risk factors for cardiovascular conditions across the life span, or whether the occurrence of a hypertensive disorder in pregnancy alters risk and predisposes one to future cardiovascular disease.

Etiology and Natural History

The underlying cause of preeclampsia is not fully understood, however, the placenta is thought to play a central role in its development. Preeclampsia can occur even in pregnancies without fetal development (e.g. hydatidiform mole) and it does not appear to regress until the placenta is removed or resorbed.²⁷ The development of preeclampsia is usually presented as involving at least two stages—the placental stage and the maternal stage. The placental stage involves abnormal placentation, which contributes to early-onset preeclampsia (delivery <34 weeks gestation), and/or placental microvillus overcrowding, which develops with advancing pregnancy and is thought to contribute to late-onset of preeclampsia.^{28, 29} The consequence of these changes is reduced placental perfusion, leading to hypoxia, placental ischemia, oxidative stress, and ultimately the release of damaging factors (i.e., cellular debris, oxidized lipids, antiangiogenic factors, soluble endoglin) into the maternal blood stream.^{27, 30-32} The second, “maternal” stage involves the development of systemic maternal sequelae resulting from placental dysfunction.^{27, 30} Placental damage leads to activation of platelets and the maternal clotting system,³³ and a systemic inflammatory response.²⁸ Changes in the renin-angiotensin-aldosterone system and increased sensitivity of blood vessels to contractile agents result in vasoconstriction.³⁰ Reduced perfusion, resulting from vasoconstriction, vascular occlusion by microthrombi, reduced vascular volume from leaking of fluid from the intravascular compartment, and vascular inflammation affect virtually every maternal organ in a women with preeclampsia.^{27, 29} Placental perfusion abnormalities may also affect the fetus, leading to increased-risk of fetal growth restriction among women with preeclampsia.

Risk Factors

It is not clear why abnormal placentation occurs and preeclampsia develops, but the process may be influenced by maternal genetic, environmental, and immunologic factors.^{27, 30, 34} In addition, the observation of heightened risk of preeclampsia during first pregnancies and in women who undergo in vitro fertilization with donor eggs has led to numerous investigations regarding the potential interaction of maternal physiology with fetal/paternal genes.^{2, 35}

Factors associated with risk of preeclampsia can be classified into those obtained by clinical history, clinical exam, laboratory tests, and imaging. Patient characteristics associated with increased risk for preeclampsia include high pre-pregnancy BMI; nulliparity; Black non-Hispanic race/ethnicity;^{10, 36} maternal age greater than 35 years; family history of preeclampsia, specifically a mother or sister with preeclampsia; and family history of early-onset cardiovascular disease. Current or past medical conditions that increase the risk of preeclampsia include preeclampsia, placental abruption, or stillbirth in a prior pregnancy; the presence of multifetal gestation; autoimmune disease, such as systemic lupus erythematosus or antiphospholipid antibody syndrome; pregestational diabetes mellitus; chronic hypertension; renal disease; and conception via assisted reproductive technology.³⁷ Individuals at highest risk for developing preeclampsia are those with autoimmune disease; preeclampsia or hypertensive disease in previous pregnancy; preexisting diabetes mellitus; and/or renal disease. Rates of preeclampsia associated with different risk factors vary. The medical history risk factors described above are more strongly associated with incidence of preeclampsia than patient characteristics such as nulliparity and maternal age, doubling or tripling the risk of developing preeclampsia. The presence of multiple risk factors, whether personal characteristics or medical history risk factors, further heightens risk for the condition. Individuals at lowest risk are those who have previously had an uncomplicated term pregnancy and do not have any conditions or circumstances known to increase risk.

Various maternal measures, including lab tests, blood pressure response to a stimulus, and ultrasound assessments of uterine artery blood flow, have been evaluated as means to identify women at increased risk of developing preeclampsia. In addition, several models have been developed that aim to identify women who are at risk of developing preeclampsia. Many of these models include variables for medical history, patient characteristics, blood serum biomarkers (e.g., serum placental growth factor), mean arterial pressure (MAP), and ultrasound readings (e.g., Doppler uterine artery pulsatility index). The most extensively researched of these are various iterations of the Fetal Medicine Foundation (FMF) model; a recent publication provides the most complete development and validation evidence to date.³⁸ As in earlier studies of the model, performance was best for prediction of the more rare cases of preeclampsia requiring delivery early in pregnancy (<34 weeks). Evidence to support clinical application of available risk prediction models is limited, as there are very few external validation and implementation studies, and it is unclear whether risk prediction models are superior to risk assessment based on clinical history taking.³⁹

Treatment Approaches

The only curative treatment for preeclampsia is delivery of the placenta. Current recommendations for delivery of patients with preeclampsia consider the risks and benefits both to the mother and the fetus. For women with gestational hypertension and preeclampsia without severe features, delivery is recommended when the patient reaches 37 0/7 weeks of gestation. For preeclampsia with severe features, delivery is recommended at 34 0/7 weeks of gestation, or at the time it occurs if after 34 0/7 weeks. Given the serious morbidity associated with HELLP syndrome, delivery is recommended regardless of gestational age at which it arises. Similarly, women with eclampsia should be delivered in a timely manner once the patient is stabilized. Close antenatal surveillance of women with gestational hypertension or preeclampsia without severe features arising prior to 37 weeks of gestation is required to assess for worsening disease or fetal compromise that might require more expedient delivery.¹

Secondary prevention measures to avoid serious complications from preeclampsia include administration of antihypertensive medications to prevent stroke and administration of magnesium sulfate to prevent eclamptic seizures.^{1, 40-43}

Preeclampsia Prevention

Interventions aimed at reducing the risk of preeclampsia have focused on several of the pathways potentially involved in its development, including vasoconstriction and platelet aggregation as well as metabolic abnormalities and nutritional deficiencies such as inadequate dietary calcium.⁴⁴ In a systematic review of 27 randomized trials of calcium supplementation for the prevention of preeclampsia, high-dose calcium supplementation (≥ 1 g/day) was shown to reduce the risk of preeclampsia and preterm birth, among women with a history of a low calcium diet (< 900 mg/day).⁴⁵ Currently, the World Health Organization⁴⁶ recommends daily calcium supplementation (1.5-2.0 g oral elemental calcium) for pregnant women with low dietary calcium intake to reduce the risk of preeclampsia, whereas ACOG does not.¹ Medications such as heparin, metformin, sildenafil, and statins, nutritional supplements, and dietary modifications have been or are being explored for the prevention of preeclampsia, but, aside from calcium, none have been identified as efficacious.⁴⁷⁻⁵²

Acetylsalicylic acid (aspirin) demonstrates inhibitory effects on cyclooxygenases (COX) leading to shifts in the synthesis of prostacyclin and thromboxane A₂.⁵³ This results in a two-fold action, reducing both platelet aggregation and vasoconstriction. Aspirin also leads to reduction in the synthesis of other prostaglandins, such as PGE₂, which are involved with inflammation, contributing to aspirin's effect as an anti-inflammatory agent.⁵⁴ The inhibition of platelet aggregation and reduced vasoconstriction, as well as aspirin's anti-inflammatory effects⁵⁵ are thought to provide the basis by which aspirin administration may reduce the risk of development of gestational hypertension and preeclampsia.^{53, 56, 57}

While no interventions eliminate preeclampsia risk, evidence from a moderate-sized body of trial evidence that includes several large trials has established that aspirin modestly reduces the risk of preeclampsia in increased-risk populations.⁵⁸⁻⁶⁰ Trials conducted among average-risk populations

where preeclampsia prevalence ranges from 3 to 5 percent have not consistently demonstrated preventive benefits, and concerns about potential bleeding from aspirin use as well as lack of data regarding long-term effects on the offspring⁵⁵ have supported guidelines focused on increased-risk populations.^{1, 61} There have, however, been modeling studies and clinical discussions in recent years arguing for broader use of aspirin for prevention of preeclampsia. Most recently, a large multisite trial conducted in low income and middle income countries reported a statistically significant benefit of low dose aspirin use for prevention of preterm birth and perinatal mortality among nulliparous individuals.⁶²

Previous USPSTF Recommendation and Recommendations of Others

In 2014, the USPSTF concluded with moderate certainty that there is a substantial net benefit of daily low dose aspirin use to reduce the risk for preeclampsia, preterm birth, and intrauterine growth restriction in women at high risk for preeclampsia, and recommended the prescription of low dose (81 mg/d) aspirin after 12 weeks of gestation to asymptomatic pregnant women who are at high risk for preeclampsia (B recommendation).

The USPSTF recommendation of low dose aspirin for women at high risk of preeclampsia was evaluated in a retrospective cohort study using a database of deliveries from two hospitals within a single academic institution.⁶³ Rates of recurrent preeclampsia for women with a history of preeclampsia in a prior pregnancy were compared for periods before and after the 2014 recommendation, spanning recommendation August 2011 to June 2016. Results showed that the risk of recurrent preeclampsia was 30% lower in the “after” group (adjusted relative risk, 0.70; 95% CI, 0.52, 0.95).⁶³

Aspirin for women at high risk of preeclampsia is currently the only recommended method of prevention.¹ Low dose aspirin is recommended by the World Health Organization,⁶⁴ the American College of Obstetricians and Gynecologists,⁶⁵ the National Institute for Health and Clinical Excellence,⁶⁶ and the American Heart Association/American Stroke Association (**Table 1**).⁶⁷

Chapter 2. Methods

Scope and Purpose

This systematic review will provide updated evidence regarding the effectiveness of aspirin in reducing adverse maternal, perinatal, and child outcomes, as well as the effectiveness of aspirin in preventing preeclampsia in those at risk. In addition, the harms of aspirin use to prevent preeclampsia will be evaluated. The USPSTF will use this review to update its 2014 recommendation for primary care practices.⁶⁸ This review includes all trials from the previous review⁶⁹ that met current inclusion/exclusion criteria as well as newly identified studies.

Key Questions and Analytic Framework

We followed USPSTF procedures and methods to define study inclusion and exclusion criteria (**Appendix A Table 1**) and developed an analytic framework (**Figure 1**) with three Key Questions (KQs).

The KQs are:

1. Does aspirin reduce adverse maternal, perinatal, child, or combined health outcomes in pregnant persons at increased risk of preeclampsia?
 - a. Does the effectiveness of aspirin for reducing adverse health outcomes vary by subpopulations defined by personal characteristics or preeclampsia risk factors?
2. Does aspirin prevent preeclampsia in pregnant persons at increased risk for preeclampsia?
 - a. Does the effectiveness of aspirin for reducing preeclampsia vary by subpopulations defined by personal characteristics or preeclampsia risk factors?
3. What are the harms of aspirin use to prevent preeclampsia during pregnancy?
 - a. Do the harms of aspirin use to prevent preeclampsia vary by subpopulations defined by personal characteristics or preeclampsia risk factors?

Data Sources and Searches

We considered all studies from the previous review⁷⁰ conducted to support the 2014 USPSTF recommendation⁷¹ on this topic for inclusion in the current review and performed a comprehensive search of MEDLINE, PubMed (publisher-supplied only), Embase, and the Cochrane Collaboration Registry of Controlled Trials for relevant studies published between January 2013 and May 15, 2020. This search was designed to partially overlap with the end search dates of the previous USPSTF-commissioned review of this topic, and to identify new studies. Studies included in the previous USPSTF review were evaluated for inclusion against the inclusion and exclusion criteria for the current review. A research librarian developed and executed the search, which was peer-reviewed by a second research librarian (**Appendix A**).

We also examined the reference lists of other previously published reviews, meta-analyses, and

primary studies to identify additional potential studies for inclusion. We supplemented our searches with suggestions from experts and articles identified through news and table-of-contents alerts. We also searched ClinicalTrials.gov (<https://ClinicalTrials.gov/>) for ongoing trials (**Appendix G**). We imported the literature from these sources directly into EndNote® X9 (Thomson Reuters, New York, NY).

Study Selection

Two reviewers independently reviewed the titles and abstracts of all identified articles to determine whether studies met inclusion and exclusion criteria for design, population, intervention, and outcomes (**Appendix A Table 1**). Two reviewers then independently evaluated the full-text article(s) of all potentially included studies against the complete inclusion and exclusion criteria. Disagreements regarding the abstract and/or full-text review were resolved by discussion and consultation with a third reviewer if necessary. Excluded studies and reasons for exclusion are listed in **Appendix D**.

We developed an *a priori* set of criteria for inclusion and exclusion of studies with input from USPSTF members and based on our understanding of the literature (**Appendix A Table 1**). For the KQ1 and KQ2 questions of effectiveness, RCTs and individual participant data meta-analyses (IPD-MAs) of pregnant persons at increased risk for preeclampsia were considered for inclusion. Risk of preeclampsia was determined based on personal sociodemographic characteristics, medical history, diagnostic measurements or assays, or risk prediction models. For the KQ3 question evaluating harms, these criteria were expanded to include (1) RCTs conducted in lower or average-risk populations; and (2) comparative observational studies of pregnant persons exposed to aspirin for preeclampsia prevention over the course of pregnancy, as well as their similarly exposed fetuses, infants, and children. **Appendix A Table 1** lists all outcomes that were considered for inclusion in the review. Studies limited to persons seeking fertility treatments were not included. Only studies of daily aspirin (≥ 50 mg) for the primary prevention of preeclampsia were considered for inclusion, but studies evaluating nonaspirin antiplatelet medications or aspirin combined with other potentially active interventions (e.g., dietary supplements, weight loss), or studies of aspirin aimed at preventing other complications of pregnancy such as miscarriage, were not. The review was limited to studies conducted in countries with “very high” Human Development Index (2016) scores as published by the United Nations Development Programme.⁷² We also limited our included studies to those published in English and deemed good- or fair-quality based on USPSTF quality rating standards.⁷³

Quality Assessment and Data Abstraction

Two reviewers applied USPSTF design-specific criteria to assess the methodological quality of all eligible studies (**Appendix A Table 2**). We assigned each study a quality rating of “good,” “fair,” or “poor.” Discordant quality ratings were resolved by discussion or adjudicated by a third reviewer as needed. Studies rated as poor quality were excluded from the review.

Good-quality RCTs were those that met all or nearly all of the specified quality criteria (e.g.,

comparable groups were assembled initially and maintained throughout the study, followup was 90% or higher, assessment procedures were described and blinded if they involved direct interviews, randomization methods were described, allocation was concealed). Fair-quality studies did not meet all criteria but did not have serious threats to their internal validity related to design, execution, or reporting. Intervention studies rated as poor quality generally had several important limitations, including at least one of the following risks of bias: very high attrition (generally > 40%); differential attrition between intervention arms (generally > 20%); lack of baseline comparability between groups without adjustment; or problematic issues in trial conduct, analysis, or reporting of results (e.g., possible selective reporting; inappropriate exclusion of participants from analyses; questionable validity of allocation or assessment procedures).

For all of the included studies, one reviewer extracted key elements into standardized abstraction forms in DistillerSR (Evidence Partners, Ottawa, Canada). Key elements included general characteristics of the study (e.g., author, year, study design), clinical and demographic characteristics of the sample and setting (e.g., age, race/ethnicity, baseline clinical characteristics, setting, country), analytic methods, and results. A second reviewer checked the data for accuracy.

When abstracting reported data on health outcomes, we included the number of participants experiencing an event, as well as incidence rates, as appropriate. For KQ1, we abstracted the following maternal health outcomes: eclampsia, puerperal cerebrovascular disorder, cerebrovascular hemorrhage, edema, or embolus; renal or hepatic injury/failure; pulmonary edema; mortality; mental health diagnoses or symptoms; and any reported measures of well-being. Fetal, neonatal, and child health outcomes included preterm delivery; mean gestational age; low birth weight; intrauterine growth restriction/small for gestational age; and stillbirth or neonatal mortality. The study defined definitions of outcomes were described and the most commonly reported were combined for meta-analysis. Where available, we aimed to abstract information on the number of early and very early preterm births, and separate cases of spontaneous from induced preterm birth. For KQ2 (preeclampsia prevention effectiveness), we abstracted the incidence of preeclampsia reported in each RCT. For KQ 3 (harms of aspirin use for preeclampsia prevention), we abstracted reports of abruptio placentae, intracranial fetal bleeding, postpartum hemorrhage or estimated blood loss, and any other major harm to the mother or fetus reported.

Data Synthesis and Analysis

This review was conducted in accordance with current USPSTF procedures and systematic review methods.⁷³ Tables describing key study design and participant characteristics were created. Pooled effect estimates of the relative risks of health outcomes associated with aspirin use were generated using random effects model with restricted maximum likelihood (REML) and Knapp-Hartung adjustment methods.^{74, 75} This analytic approach overcomes limitations of other estimators for tau-square that have been found to generate overly precise confidence limits, especially when pooling fewer than 20 studies.⁷⁶ For rare outcomes (i.e., perinatal mortality) we conducted sensitivity analyses using the Peto odds ratio method,⁷⁷ as well as pooled analyses

including and excluding studies that had no events in either study arm, to assess the robustness of the final RR pooled estimates.⁷⁸ We identified study design and participant characteristics that could possibly explain heterogeneity in aspirin effectiveness among the included studies. To examine subgroup differences, we generated stratified forest plots using the Cochran Q statistical test for interaction and conducted REML meta-regression analyses using the Knapp Hartung correction. Variables anticipated *a priori* to be potential sources of heterogeneity included aspirin dosage, timing, and duration; population risk characteristics (e.g., incidence of preeclampsia in the control condition, strategy for selecting participants); study size; and control condition (i.e., placebo versus no treatment). We defined comparison groups based on clinically meaningful distinctions. For comparisons of effects by population risk of preeclampsia, we compared studies where more than 12 percent of participants developed preeclampsia in the control group to studies with lower incidence rates. This threshold was selected to distinguish trials with approximately 3 times incidence seen in the general population. We also compared studies that used a Doppler ultrasound test as part of the risk assessment procedure with those that did not, and studies that used any lab or imaging tests compared to those relying only on clinical history and exams. For each KQ, the statistical heterogeneity of included studies was estimated with I^2 statistics and Tau^2 . The distribution of trial results was examined with funnel plots and Peters tests to assess whether there was evidence of small-study effects.⁷⁹⁻⁸¹

In accordance with published methodological guidance on evaluating subgroup effects within trials,⁸² we abstracted the results of within study differences to support inferences related to subgroup effects only if statistically valid tests for interaction were reported.

For each outcome measure, we calculated the number needed to treat (NNT) by first estimating the absolute risk reduction based on the pooled risk ratio and two to three estimated levels of baseline risk of the outcome of interest (i.e., absolute risk reduction=(risk ratio-1)*population risk). Because there was a wide range of risks for some of the outcomes (preeclampsia, SGA/IUGR, preterm birth), we selected population risk levels empirically using the included studies corresponding to the 20th, 50th, and 80th percentiles of outcome incidence in the control arm of the trials. For perinatal mortality we used the lowest and highest incidence levels reported in trials included in the pooled analysis for this outcome. From the absolute risk reduction for every level of baseline risk tested, we calculated the NNT as the inverse of the absolute risk ratio.⁸³

We used Stata version 16.1 (StataCorp LP, College Station, TX) for all analyses. All significance testing was two-sided, and results were considered statistically significant if the p-value was 0.05 or less.

Grading the Strength of the Body of Evidence

The strength of the overall body of evidence for each KQ was graded using an adaptation of the Evidence-based Practice Center approach,⁸⁴ which is based on a system developed by the Grading of Recommendations Assessment, Development and Evaluation (GRADE) Working Group.⁸⁵ This adaptation explicitly addresses four of the five Evidence-based Practice Center-required domains: consistency (similarity of effect direction and size), precision (degree of

certainty around an estimate), reporting bias (potential for bias related to publication, selective outcome reporting, or selective analysis reporting), and study quality (i.e., study limitations, risk of bias). We do not evaluate the fifth domain—directness—as it is implied in the structure of the KQs (i.e., pertains to whether the evidence links the interventions directly to a health outcome).

Consistency was rated as reasonably consistent, inconsistent, or not applicable (e.g., single study). Precision was rated as reasonably precise, imprecise, or not applicable (e.g., no evidence). Reporting bias was rated as suspected, undetected, or not applicable (e.g., when there was insufficient evidence for a particular outcome). Study quality summarizes the quality ratings of the individual trials included for an outcome and indicates the degree to which the results are likely to have adequately low risk of bias. The limitations domain highlighted important restrictions in answering the overall KQ (e.g., lack of replication of interventions, nonreporting of outcomes important to patients).

The overall strength of evidence was graded as “high,” “moderate,” “low,” or “insufficient.” “High” indicates high confidence that the evidence reflects the true effect and that further research is very unlikely to change our confidence in the estimate of effects. “Moderate” indicates moderate confidence that the evidence reflects the true effect and that further research may change our confidence in the estimate of effect and may change the estimate. “Low” indicates low confidence that the evidence reflects the true effect and that further research is likely to change our confidence in the estimate of effect and to change the estimate. A grade of “insufficient” indicates that evidence is either unavailable or does not permit estimate of an effect. At least two independent reviewers rated the overall strength of evidence for each intervention type. We resolved discrepancies through consensus discussion involving more reviewers.

Expert Review and Public Comment

A draft Research Plan for this review was posted for public comment on the USPSTF website from June 20 to July 23, 2019. Several minor additions and wording changes were made to improve the clarity and specificity of the inclusion criteria. Additional clinically important categories used for defining preeclampsia and preterm birth were included to more fully represent health outcomes that may be reported in the literature. A final research plan was posted on the USPSTF website on September 26, 2019.

USPSTF Involvement

We worked with four USPSTF members at key points throughout this review, particularly when determining the scope and methods for the review and when developing the Analytic Framework and Key Questions. After revisions reflecting the public comment period, the USPSTF members approved the final analytic framework, KQs, and inclusion and exclusion criteria. AHRQ funded this review under a contract to support the work of the USPSTF. An AHRQ Medical Officer provided project oversight, reviewed the draft report, and assisted in the external review of the report.

Chapter 3. Results

Description of Included Studies

Our literature search yielded 3749 unique citations. From these, we reviewed the full text of 183 articles (**Appendix B Figure 1**). Of these, 23 studies (33 articles) met our inclusion criteria. We excluded the remaining 150 full-text articles (**Appendix C**).

Of the 23 included studies (n=26,850) 19 are carried forward from the previous USPSTF review conducted in 2014.^{56, 86-103} These studies met the inclusion and exclusion criteria of the current review and were combined with four newly identified RCTs.^{58, 104-106} Four studies included in the previous review were excluded based on risk of bias and eligibility determinations according to the scope of the current review.¹⁰⁷⁻¹¹⁰ Among the included studies, 18 (10 good-quality) met the inclusion criteria for KQ 1 (maternal and perinatal health outcomes); 16 (10 good-quality) met the inclusion criteria for KQ 2 (preeclampsia); and 21 (14 good-quality) met the inclusion criteria for KQ 3 (harms). One IPD-MA was identified but was not included because our prespecified inclusion and exclusion criteria for intervention and setting were not met. The results of this study are discussed in the next chapter. For KQs 1 and 2, we included trials that enrolled pregnant individuals at increased risk of preeclampsia. For KQ 3 we included studies reporting harms from trials enrolling pregnant individuals at increased risk (16 included trials) and at average risk of preeclampsia (5 included trials) (**Appendix B Figure 1**).

Population and Setting Details

Tables 2 and 3 provide summary information on study population, setting, design, and intervention characteristics of the included studies. For KQ 1 and KQ 2, assessing the effectiveness of aspirin in preventing preeclampsia and related health outcomes for pregnant individuals at increased risk for preeclampsia, we included a large Maternal Fetal Medicine Unit Network (MFMU) trial (n=2,539) funded by National Institute of Child Health and Human Development that recruited participants from 13 clinical sites in the United States from May 1991 to June 1995.⁹⁸ Two large international trials coordinated from the United Kingdom were also included:^{58, 94} The Collaborative Low dose Aspirin Study in Pregnancy (CLASP) (n=9,364) trial enrolled participants from 213 centers across 16 countries from January 1988 to December 1992,⁹⁴ and the Combined Multimarker Screening and Randomized Treatment with Aspirin for Evidence-based Preeclampsia Prevention (ASPRE) trial (n=1,776) enrolled participants 13 maternity hospitals based in 6 countries (United Kingdom, Spain, Italy, Belgium, Greece, Israel) from April 2014 to April 2016.⁵⁸ There were 15 smaller trials included from various countries.^{56, 86-89, 92, 93, 95-97, 100, 102, 103, 105, 106} For KQ 3, assessing harms associated with aspirin use during pregnancy, we included five additional RCTs of women considered to be at average risk for preeclampsia.^{90, 91, 99, 101, 104} These five RCTs included a good-quality multisite study conducted in the United States MFMU sites that enrolled average- or low-risk participants (n=3,135),⁹¹ a smaller U.S.-based study (n=606),⁹⁰ a large good-quality multisite study in France and Belgium (n=3,294),¹⁰¹ a large good-quality hospital-based study in Barbados (n=3,647),⁹⁹ and a smaller fair-quality study from Ireland (n=362).¹⁰⁴

Among trials included to assess the effectiveness of aspirin, a variety of study procedures and criteria were used to identify populations of pregnant persons at increased risk for preeclampsia (**Table 4**). The most common method was examination of participant characteristics and personal or family medical history (e.g., age, BMI, parity, multifetal gestation, history of hypertensive disorders, history of pregnancy complications). Among these studies, history of hypertensive disorders (preeclampsia, gestational hypertension, or chronic hypertension), alone or in combination with other risk factors, was used as the primary method to identify trial participants in ten studies,^{56, 87, 88, 92, 94, 96-98, 100, 103} and multifetal gestation was used in six trials.^{56, 88, 93, 94, 97, 98} Other clinical risk factors used included metabolic disease,^{56, 97} diabetes (prepregnancy or gestational),^{98, 103} history of IUGR/SGA,^{87, 100, 103} spontaneous abortion,^{56, 97, 103} or stillbirth;^{87, 100} renal disease;^{94, 96} and maternal age.^{56, 94, 97}

Most risk factors and clinical tests used for identifying increased-risk populations, whether used alone or in combination with other factors, were associated with at least a two-fold higher risk for preeclampsia. An exception to this was nulliparity, which is understood to be a more modest independent risk factor for preeclampsia. Nulliparity was combined with other risk factors to identify an increased-risk population in three of the effectiveness trials.^{56, 88, 97, 106} Nulliparity alone was used to select participants in four of the five studies enrolling average-risk populations that were included for evaluating harms of aspirin use in pregnancy.^{90, 91, 101, 104} These studies, which reported preeclampsia incidence in the control condition ranging from 2 to 6 percent (similar to the population average), were not included for evaluating KQ1 and KQ2 as nulliparity alone was not considered sufficient evidence of increased risk in accordance with the scope defined for the previous and current review.

Risk assessment also frequently involved diagnostic measurements or assays, either as the primary means to identify individuals at risk, or in combination with medical history and personal characteristics. Abnormal Doppler readings were used in six studies.^{58, 89, 102, 103, 105, 106} One study relied on angiotensin-II sensitivity identified via blood pressure response to IV infusion of angiotensin II.⁸⁶ Another required a positive rollover test in which rolling the participant from her left side to her back resulted in rise of diastolic blood pressure >15 mg Hg above baseline.⁸⁸ One study used a second trimester hemoglobin concentration greater than 13.2 g/dL to determine the risk of an individual participant in the trial.⁹⁵ Finally, one trial utilized a prediction model that combined maternal demographic factors and medical history, mean arterial pressure, uterine-artery pulsatility index, and maternal serum pregnancy-associated plasma protein A and placental growth factor to identify persons at increased risk for preterm birth secondary to preeclampsia.⁵⁸

The inclusion and exclusion criteria in the trials were usually well described, but details on the baseline demographic characteristics and risk factors of enrolled participants were sparsely and inconsistently reported (**Appendix E Table 1**). Overall, the population of women included in the trials was young (mean age ranging from 20.4 to 33.5 years) and predominantly White, with only three trials reporting majority populations of Black women (ranging from approximately 50% to 72%).^{90, 91, 98} Parity was reported in most trials, with the proportion nulliparous ranging from approximately 15 to 100 percent of study participants. Similarly, the proportion of participants with a history of chronic hypertension at baseline ranged from no participants (in low or average-risk population studies included for KQ3) to nearly all having this history (91%). History of

smoking and current smoking were reported in 13 trials and ranged from approximately 10 to 40 percent of participants.^{58, 86, 87, 89, 91, 94, 95, 98, 101-105} Mean BMI ranged from 22.3 to 29.7 kg/m² in the studies reporting this baseline characteristic.

Intervention Characteristics

Included trials varied widely in timing of and dosages of aspirin treatment (**Table 2, Table 3**). The gestational age at which aspirin therapy was initiated across the trials varied substantially with nine trials starting therapy in participants as early 11 to 12 weeks of gestation and five allowing initiation to continue as late as 36 to 38 weeks of gestation. The most common date of aspirin discontinuation was delivery, but eight trials stopped aspirin prophylaxis before delivery,^{58, 88, 96, 100-104} as early as 34 weeks¹⁰¹ or at the point when preeclampsia developed.⁹⁸ Aspirin dosages ranged from 50 to 150 mg daily. The majority of trials used either dosages of 60 mg (6 trials)^{86, 87, 90, 91, 94, 98} or 100 mg (9 trials);^{56, 88, 93, 96, 97, 100, 101, 103, 106} two of the newly identified trials used a higher dose of 150 mg per day.^{58, 105} A matching placebo was the comparator in all trials except in one study included for harms, in which participants in the control arm received usual care with no placebo.¹⁰⁴

Meta-analysis was possible for seven outcomes that were reported with enough consistency across multiple studies to support valid estimation of effects. The relevant results related to each Key Question are presented in detail below (**Table 5**). The three largest included trials, MFMU, CLASP, and ASPRE,^{58, 94, 98, 111-116} conducted prespecified or exploratory comparisons and reported statistical tests for interaction suitable for evaluating subgroup differences (**Appendix F Table 1**).

KQ1. Does Aspirin Reduce Adverse Maternal, Perinatal, Child, or Combined Health Outcomes in Pregnant Persons at Increased Risk of Preeclampsia?

Summary of Results

Eighteen trials (n=15,908) reported maternal, perinatal, or child health outcomes for daily aspirin use (50mg to 150mg daily) compared with a matching placebo starting as early as 11 weeks or as late as 32 weeks of gestation and continuing until late pregnancy or delivery. In pooled analyses, aspirin was consistently associated with a reduced risk of perinatal mortality, preterm birth, and SGA/IUGR (**Table 5**). Preventive effects of aspirin on the three main outcomes ranged from 20 to 30 percent reduced risk. Evidence from one within-trial comparison indicated that aspirin may be more effective among nonsmokers than among smokers. Aggregate comparisons between different design and participant characteristics of the included studies suggested that the benefit of aspirin for preventing preterm birth may be larger when aspirin was started before 16 weeks of gestation than when started later. There also was evidence of a larger effect of aspirin for preventing preterm birth in studies enrolling a larger proportion of nulliparous participants (who also had other risk factors). We did not observe statistically significant differences between subgroups for the preeclampsia or SGA/IUGR outcomes, however, and differences for preterm

birth could be attributed to multiple comparisons. Small-study effects could not be ruled out and might lead to some overestimation of pooled effect sizes. Confounding of study size with other study and participant characteristics could have influenced subgroup comparisons. For example, the two largest studies reported the smallest risk reductions for all outcomes, but these studies also both started treatment later than 16 weeks of gestation and used very low aspirin dosages. Other rare health outcomes, such as eclampsia and maternal mortality, occurred too infrequently to estimate preventive effectiveness.

Detailed Results

Maternal Health

Direct maternal health consequences of preeclampsia are extremely rare and include eclampsia, HELLP syndrome, stroke, organ failure, and death. Few trials reported these events, and pooled estimates were not possible to compute. In the few large studies with these and other maternal health outcomes, risk estimates were too imprecise to assess differences by study group (**Appendix E Table 2**). Many of the perinatal outcomes described below also have implications for maternal health and well-being, as do outcomes described as potential harms for KQ3.

Perinatal Mortality

Fifteen trials reported on perinatal mortality (N = 15,527). Four of these reported no perinatal deaths in either study arm and were not included in the pooled analysis.^{88, 95, 97, 106} Seven reported two or fewer events in one or both study arms.^{56, 86, 87, 89, 92, 93, 96} No individual study reported a statistically significant difference in perinatal mortality and all were underpowered for this outcome, but the pooled effect estimate showed a statistically significant 21 percent reduction in the risk of perinatal mortality associated with aspirin use (pooled RR 0.79, 95% CI 0.66, 0.96; I^2 0%; k=11, n=13,860) (**Figure 2, Appendix E Table 3**). The pooled effect was consistent with the findings from the three largest trials, which reported more cases of perinatal mortality in the control group than in the aspirin group, with relative risks ranging from 0.59 to 0.80. Smaller trials reported effects that were less consistently in the direction of a benefit, but confidence intervals were overlapping.

Preterm Birth

Thirteen trials reported on preterm birth (<37 weeks of gestation) (N = 15,213). Results across these trials were consistently in the direction of a preventive benefit of daily aspirin use, with no trial finding more cases of preterm birth in the aspirin group than in the placebo group. The estimated pooled effect showed a 20 percent reduced risk of preterm birth associated with intervention (pooled RR 0.80, 95% CI 0.67, 0.95; I^2 49%; k=13, n=13,619) (**Figure 3, Appendix E Table 4**). The primary outcome of the ASPRE trial⁵⁸ was “preterm preeclampsia” a composite of preeclampsia and preterm birth. For pooled analyses, we used reported data from this trial to compute a preterm birth outcome consistent with what was reported in all other trials: preterm birth with or without a preeclampsia diagnosis. The combination of preeclampsia and preterm birth is a less common outcome, with ASPRE reporting 1.6 percent (13/798) in the aspirin group and 4.3 percent (35/822) in the placebo group, and the reported effectiveness of aspirin for the

outcome (OR 0.38, 95% CI, 0.20, 0.74) was greater than for the prevention of any preterm birth. Nevertheless, the computed relative risk of any preterm birth for the ASPRE trial also favored aspirin (RR 0.65, 95% CI, 0.47, 0.90). Trials rarely reported cases of spontaneous preterm birth separately from induced preterm birth, and few studies reported the number of cases of very early and early preterm birth included in the preterm birth outcome.

Small for Gestational Age or Intrauterine Growth Restriction

Sixteen trials reported on SGA/IUGR (N = 15,757) and all but two^{89, 98} reported fewer cases in the aspirin group than in the placebo group. The estimated pooled effect indicated a 18 percent reduced risk of having an SGA/IUGR infant for women who took aspirin (pooled RR 0.82, 95% CI 0.68, 0.99; I² 41%; k=16; n=14,385) (**Figure 4, Appendix E Table 5**).

Studies did not consistently report birthweight or gestational age at birth, but the results that were reported are consistent with the aspirin effects on SGA/IUGR and preterm birth (**Appendix E Tables 6 and 7, Appendix F Figure 1**). Combined mean birthweight (grams) was higher with aspirin use than with placebo across eleven studies (n=10,987) that reported the outcome (weighted mean difference 126.9; [95% CI 39.8, 214.0]; I² 50.0%).

Small-Study Effects

Tests for small-study effects were statistically significant for preterm birth (Peters p = 0.026) and approached statistical significance for SGA/IUGR (Peters p = 0.072) (data not shown). This suggests that the pooled effects for these outcomes may overestimate the effectiveness of aspirin due to differences in the way smaller studies were conducted or the absence of small null or negative studies in the published literature. Visual inspection of funnel plots also revealed an imbalance of the distribution of study findings and standard errors consistent with the statistically significant test for small-study effects. Because random effects meta-analysis modeling weights smaller studies more heavily, they can exacerbate bias due to small-study effects.⁸³ The fixed effect sensitivity analysis (data not shown) suggests that the random effect estimates for SGA/IUGR and preterm birth could be slightly overestimated due to small-study effects, but that the random effects confidence intervals provide good coverage for the range of likely effects.

KQ1a. Does the Effectiveness of Aspirin for Reducing Adverse Health Outcomes Vary by Subpopulations Defined by Personal Characteristics or Preeclampsia Risk Factors?

Valid tests for subgroup differences were available from the three largest trials (**Appendix F Table 1**).^{94, 98, 111} We conducted subgroup comparisons at the aggregate level using stratified forest plots and meta-regression. Subgroup differences were assessed for the outcomes of preterm birth (<37 weeks of gestation) and SGA/IUGR since these outcomes had enough studies and precision to reasonably support statistical comparisons between groups (**Appendix F Figure 3 and Figure 4**). These study level comparisons should be interpreted with reservations given

the evidence of small-study effects for these outcomes and others, and as the two largest trials started aspirin use later in pregnancy (≥ 16 weeks of gestation) and used very low aspirin dosages (60mg).

Aspirin Timing and Dosage

The CLASP trial compared the effect of aspirin (60 mg) initiated at ≤ 20 weeks versus >20 weeks of gestation for preventing preterm birth, SGA/IUGR, and perinatal mortality and found no statistically significant differences (**Appendix F Table 1**).⁹⁴ Similarly, no differences were found in the MFMU trial for comparisons of participants initiating aspirin before or after 16 weeks of gestation.⁹⁸

In across study comparisons of subgroup study effects, a comparison of the pooled relative risk of IUGR/SGA between studies with aspirin started before 16 weeks of gestation (pooled RR, 0.59 [95% CI, 0.41, 0.86], I^2 35%, $k=6$, $n=2,351$) and those where the majority of participants began daily aspirin use later (pooled RR, 0.95 [95% CI, 0.80, 1.13], I^2 16%, $k=10$, $n=12,034$) was statistically significant ($p=0.03$) (**Figure 5, Appendix F Figure 3**). For prevention of preterm birth, there was further evidence that aspirin is more effective if started before 16 weeks of gestation, but only four studies reporting the outcome were in the earlier initiation subgroup; there was a statistically significant difference between the pooled relative risk of preterm birth associated with initiation of aspirin before 16 weeks of gestation (pooled RR, 0.49 [95% CI, 0.26, 0.95], I^2 28%, $k=4$, $n=2,103$) and the estimate from studies where the majority of participants began aspirin later (pooled RR, 0.91 [95% CI, 0.85, 0.96], I^2 0%, $k=9$, $n=11,516$) ($p=0.02$) (**Figure 6, Appendix F Figure 4**).

Personal Characteristics and Clinical History

The CLASP trial found no significant difference in the effectiveness of aspirin or preventing perinatal mortality, IUGR, and preterm birth (<37 weeks of gestation) between multiparous and nulliparous study participants. The MFMU trial compared preterm birth and SGA outcomes by BMI and smoking status. Only one of these contrasts was statistically significant—the effectiveness of aspirin (60 mg/day) to prevent preterm birth among nonsmokers (RR 0.91, 95% CI 0.84, 0.99) was greater than for smokers (RR 1.17, 95% CI 0.99, 1.39) ($p=0.01$) (**Appendix F Table 1**). The ASPRE trial tested differences in aspirin effectiveness for preventing preeclampsia with preterm birth for several personal characteristic and clinical history factors including maternal age, BMI, racial origin (Afro Caribbean, Caucasian, Other), assisted versus natural conception, smoking status, family history of preeclampsia, parity, and diagnosed chronic hypertension, but did not identify any statistically significant differences. Several of these post-hoc comparisons were limited by relatively small numbers of participants in certain subgroups, and low event rates for this primary outcome resulted in wide confidence intervals.

In study level comparisons, there was a significantly stronger effect of aspirin on prevention of preterm birth in studies in which a majority (more than 50%) of participants were nulliparous than in those in which nulliparous participants were a minority (**Appendix F Figure 4**). Again, however, few studies were available for comparisons, limiting interpretation of the association.

Preeclampsia Risk Assessment Approach

Differences among trials in their criteria for identifying populations at increased preeclampsia risk were not associated with differences in the effectiveness of aspirin for prevention of preterm birth or SGA/IUGR. Trials that used risk assessments based on clinical history rather than clinical tests or prediction models were not found to be less effective at identifying a population that benefited from aspirin (**Appendix F Figure 3 and Figure 4**). There was also no significant difference between effects of aspirin use on prevention of SGA/IUGR or preterm birth in studies that used abnormal Doppler ultrasound as part of the risk assessment with those that did not. There were also no significant differences in the estimated effectiveness of aspirin in studies where more or less than 12 percent of participants in the placebo group developed preeclampsia, although absolute risk reductions were greater when the incidence of preeclampsia in the control group was higher.

KQ2. Does Aspirin Prevent Preeclampsia in Pregnant Persons at Increased Risk for Preeclampsia?

Summary of Results

Aspirin use was started as early as 11 weeks of gestation and as late as 32 weeks of gestation at dosages of 50 to 150mg per day in the 16 RCTs (N=15,767) reporting preeclampsia outcomes for pregnant individuals at increased risk. The meta-analysis estimated a statistically significant 15 percent reduced risk of preeclampsia for aspirin treatment relative to placebo. The possibility that this risk reduction is overestimated could not be ruled out. We found evidence of small-study effects that could be due to the absence of smaller studies with null or negative findings in the published literature or differences in the ways smaller studies were conducted (e.g., aspirin protocols, populations enrolled, implementation, study quality). The two largest trials found modest, nonsignificant within study effects of 10 to 12 percent reduced risk of preeclampsia, however, these trials also used lower doses of aspirin than many of the other trials. Across the trials, approaches for identifying individuals at increased risk varied, as did the inclusion and exclusion criteria, generating a broad range of absolute risk reductions. We found no evidence of statistically significant differences in the magnitude of preeclampsia risk reductions related to the timing of treatment initiation; the dosage of aspirin used; or personal characteristics such as smoking history, parity, and BMI, nor were the approaches to assessing preeclampsia risk or the incidence of preeclampsia observed in the study population related to differences in effectiveness.

Detailed Results

Sixteen included RCTs (N = 15,767) reported on the effectiveness of aspirin to prevent preeclampsia (**Table 5, Appendix E Table 8**). Across these studies, the incidence of preeclampsia in the placebo condition ranged from 4 to 30 percent, reflecting the broad range of criteria used for identifying increased-risk populations (**Table 4**). Aspirin was associated with a statistically significant reduction in the risk of preeclampsia compared to placebo (pooled RR,

0.85 [95% CI, 0.75 to 0.95], I^2 0%) (**Figure 7**). Effects were consistent across the included studies; all but two small studies reported effects in the direction of a treatment benefit.^{100, 105} Of the largest trials, two using 60mg aspirin reported approximately 10 percent risk reductions with confidence intervals that were not statistically significant^{57,58} and a more recent large trial using 150mg aspirin showed a statistically significant 28 percent reduction in the risk of preeclampsia.⁵⁸

A test for small-study effects was statistically significant (Peters $p = 0.035$), suggesting that the pooled effect may overestimate the effectiveness of aspirin for this outcome due to publication bias or differences in the ways smaller studies were conducted. Visual inspection of a funnel plot also showed a tendency for larger studies to report smaller effect sizes. Because random effects meta-analysis models weight smaller studies more heavily, they can exacerbate bias due to small-study effects. Therefore, we conducted sensitivity analyses for the preeclampsia outcome using fixed effects meta-analysis. Unlike the findings for health outcomes, the fixed effects result for preeclampsia was less conservative, estimating 17 percent risk reduction with greater precision than the random effects model. Thus, we did not find evidence that the random effects modeling approach resulted in overestimation of the effect for this outcome, possibly due to the lack of statistical heterogeneity seen for the outcome.

Other Hypertensive Disorders of Pregnancy

Half of the studies reporting preeclampsia also reported the effectiveness of aspirin for prevention of gestational hypertension ($k=9$, $N=2,789$). Meta-analysis of this outcome showed inconsistency of effects, imprecision, and higher statistical heterogeneity than for more commonly reported outcomes, and could not include the two largest trials; it did not find a significant effect of aspirin use on gestational hypertension (pooled RR 0.74, [95%CI, 0.46, 1.18], I^2 51%, $k=9$, $n=2,591$) (**Appendix E Table 9, Appendix F Figure 2**). An analysis of 8 studies where it was possible to combine cases of gestational hypertension with cases of preeclampsia to create a composite outcome measure of hypertensive disorders of pregnancy showed a statistically significant effect of aspirin use (RR 0.65; [95%CI 0.44, 0.96]; I^2 59.4%) (data not shown), but the result cannot support strong inferences given the exclusion of the two largest trials from the analysis and substantial clinical and statistical heterogeneity.

KQ2a. Does the Effectiveness of Aspirin for Reducing Preeclampsia Vary by Subpopulations Defined by Personal Characteristics or Preeclampsia Risk Factors?

Valid tests for subgroup differences were available from the two largest trials.^{94, 98} We also conducted subgroup comparisons at the aggregate level using stratified plots and meta-regression.

Aspirin Timing and Dosage

Both of the largest trials (CLASP and MFMU) reported tests for differences in aspirin

effectiveness according to the timing of initiation; however, none of the comparisons were statistically significant (the CLASP trial required a 99 percent confidence range for tests of subgroup effects to account for multiple comparisons) (**Appendix F Table 1**).

In aggregate, in subgroup comparisons with the included studies in this review, we did not find an effect of aspirin administration timing on preeclampsia risk. The pooled risk reduction in studies initiating aspirin before 16 weeks of gestation was lower (RR, 0.68 [95% CI, 0.53 to 0.89], I^2 0%, $k=5$, $n=2,346$) than for studies with later initiation (RR, 0.88 [95% CI, 0.77 to 1.00], I^2 0%, $k=11$, $n=11,747$) but the difference was not statistically significant ($p=0.08$) (**Figure 8**).

There was also not a statistically significant difference in aspirin effectiveness by dosage in aggregate stratified analyses. Preeclampsia risk reduction with ≥ 75 mg of aspirin (RR, 0.72 [95% CI, 0.56, 0.93]) was slightly greater compared with < 75 mg (RR, 0.89 [95% CI, 0.74, 1.07]), but the test for an interaction was not statistically significant ($p=0.10$) (**Appendix F Figure 5**). Analyses comparing dosages at or above 100mg to lower dosages also did not yield statistically significant interactions. We also did not find any indication of a dose-response relationship in forest plots sorted by aspirin dosage.

Personal Characteristics and Clinical History

The CLASP trial found no evidence of differences in effectiveness of aspirin use (60mg) for reducing preeclampsia between nulliparous versus multiparous study participants. Comparisons from the MFMU trial found no evidence of differences in the effectiveness of aspirin for preeclampsia according to BMI (< 30 versus ≥ 30 kg/m²) or reported smoking during pregnancy (**Appendix F Table 1**). Stratified estimates of the effectiveness of aspirin by broad racial categories were conducted in the MFMU trial, but statistical tests for interaction were not conducted; a smaller magnitude risk reduction was reported for “non-White” participants but the confidence intervals overlapped for the two strata (data not shown).

Meta-regression comparisons of subgroups of the included studies for this outcome did show a significant difference in aspirin effectiveness associated with the proportion of participants who were nulliparous. The effect size was estimated to be somewhat greater in the studies where more than half of participants were nullipara at increased preeclampsia risk (RR, 0.64 [95% CI, 0.46, 0.89]) compared to those with fewer nullipara (RR, 0.89 [95% CI, 0.80, 0.99]) ($p=0.02$). (**Appendix F Figure 5**). Differences in other characteristics of these studies could account for this trend, but there were too few studies included in the review to adjust for potential confounders by including additional covariates in meta-regression.

Preeclampsia Risk Assessment Approach

We did not observe significant differences in the estimated effectiveness of aspirin for prevention of preeclampsia based on how studies identified an increased-risk population, nor between studies that identified populations where more or less than 12 percent of participants developed preeclampsia (**Appendix F Figure 5**).

KQ3. What Are the Harms of Aspirin Use to Prevent Preeclampsia During Pregnancy?

Summary of Results

Data from 21 trials (N= 26,757) provided evidence on potential harms of daily low dosages of aspirin. The most consistently reported harms outcomes were placental abruption, postpartum hemorrhage, and fetal intracranial bleeding. Other reported perinatal outcomes included cephalohematoma, congenital malformations and anomalies, and respiratory distress syndrome. We did not identify greater risks of placental abruption, postpartum hemorrhage, or fetal intracranial bleeding in participants treated with aspirin than those who were not in pooled analyses, nor were there differences within studies for less commonly reported and more rare perinatal harms. There was no evidence to indicate differences in these risks by the dosage or timing of aspirin used in the studies. Longer-term followup from one large multicenter trial was also reassuring, finding no difference in child developmental harms associated with aspirin for preeclampsia prevention taken during pregnancy. Finally, there were no significant differences found between treatment and control groups for other reported side effects or complications such as NICU admissions, rates of Cesarean deliveries, petechiae, purpura, and other instances of singular adverse events; and aspirin was generally well tolerated.

Detailed Results

We analyzed data from 21 trials on potential harms of aspirin use throughout pregnancy for the prevention of preeclampsia. Sixteen of the included trials reporting harms outcomes were conducted among pregnant individuals at increased preeclampsia risk and five were among average-risk populations. Daily aspirin doses ranged from 50mg to 150mg per day and start times ranged from 11 to 32 weeks of gestation. Pooled analyses were possible for three reported outcomes related to potential bleeding harms of aspirin use (**Table 5**).

Postpartum Hemorrhage

There was not a statistically significant difference in the risk of postpartum hemorrhage (most commonly defined as ≥ 500 mL blood loss) in 11 trials (N=23,583) randomized to aspirin or placebo (pooled RR 1.03, [95% CI, 0.94, 1.12] I^2 0%, k=9, n=23,133). Most large trials consistently reported postpartum hemorrhage as a trial outcome, with the exception of the recent ASPRE trial. There was not a statistically significant difference in the risk of postpartum hemorrhage for average-risk versus increased-risk study populations ($p = 0.12$) (**Figure 10, Table 5, Appendix E Table 10**).

Placental Abruption

Thirteen trials (N=25,761), seven conducted among increased-risk populations and five in average-risk populations, reported placental abruption. Three of the trials reported no cases in either study arm,^{92, 93, 97} and three additional trials reported 2 or fewer cases in one or both study arms.^{90, 95, 104} There was not a statistically significant difference between groups in the risk of

abruption in meta-analysis (pooled RR, 1.15 [95% CI, 0.76, 1.72] I^2 25%, $k=10$, $n=24,970$) and this effect did not differ between studies of average-risk and increased-risk populations ($p = 0.41$) (**Figure 9, Table 5, Appendix E Table 11**).

Infant Bleeding Outcomes

Aspirin was not associated with a difference in the risk of fetal intracranial bleeding (cerebral or intraventricular hemorrhage) in a meta-analysis of nine trials ($N=23,959$) using daily doses ranging from 60mg to 150mg (pooled RR, 0.90 [95% CI, 0.51, 1.57]; I^2 19%, $k=6$, $n=23,719$) (**Figure 11, Table 5**). The outcome is rare—three trials reported one or fewer cases (**Appendix E Table 12**). Two trials reported cases of cephalohematoma; one small trial reported a single case in the control group⁸⁸ and a larger trial reported more cases in the intervention condition (68/1480, 4.6%) compared to placebo (55/1505, 3.7%); however, the relative risk in the larger trial was imprecise because of the low event rate (RR, 1.26, [95% CI, 0.89, 1.78]).⁹¹ The same trial reported a composite outcome defined as any infant bleeding disorder, again finding slightly more cases in the aspirin group than the placebo group (7.0% vs. 6.5%) but not a statistically significant difference (RR, 1.08, [96% CI, 0.83, 1.41]).

Congenital Malformations and Other Infant Outcomes

Four trials reported on congenital anomalies or malformations ($N= 2,283$), which were very rare (**Appendix E Table 12**). Three of the trials reported more cases in the placebo group^{58, 96, 104} and one small trial reported a single case of anencephaly that occurred in the aspirin group.⁹² One trial reported cases of respiratory distress syndrome, with more occurring in the placebo group, but there was statistical imprecision in the estimated odds based on the study reported confidence interval estimated for this secondary outcome (aOR 0.53, [99% CI, 0.20, 1.40]).

Longer-Term Infant Harms

The CLASP trial conducted followup surveys and reported on longer-term child health outcomes for participants from trial sites in the United Kingdom and Canada. This included developmental outcomes at age 12 months ($n=4,168$ – United Kingdom followup data) and 18 months ($n=4,365$ – United Kingdom and Ottawa followup data). The trial found no differences in hospital visits, gross motor development, or height and weight measurements between treatment groups (**Appendix E Table 13**).

Other Maternal and Infant Complications

In addition to the primary, more commonly reported potential harms of aspirin evaluated above, some trials also reported on other adverse events (**Appendix E Table 12**). For the most part, comparisons between study groups for these outcomes provide assurance of the safety of aspirin during pregnancy. A large trial conducted with an average-risk population using a 75mg/day aspirin dose ($n=3647$) reported no differences in duration of hospital stay, prenatal hospital admission, blood transfusions, or rates of induced labor between participants taking aspirin than those taking placebo.⁹⁹ Another large trial in an average-risk population using a 60mg/day dose ($n=2985$) reported similar results, with no significant differences in newborn rates of petechiae,

purpura, excessive bleeding with circumcision, or any bleeding disorders between the two study groups.⁹¹ Further, there were no significant differences in the need for transfusions between those taking aspirin and those taking placebo, and no cases of bleeding complications among women who received aspirin and epidural anesthesia. The MFMU trial (n=2503) reported no differences in a composite measure of any adverse events for mothers or neonates, but provided few details.⁹⁸ In the CLASP trial there were fewer cases of infants admitted to special care units in the treatment group, and the duration of stay was similar between treated and untreated infants. In the aspirin-treated group, however, significantly more women needed a blood transfusion after delivery (4.0% vs. 3.2%, significance test not reported). The higher rate of transfusion was not associated with different rates or severity of postpartum hemorrhage. A few studies reported other potential maternal harms, including premature rupture of membranes and pulmonary embolism, and did not find differences by study group (**Appendix E Table 14**). However, these studies were underpowered for comparing rare outcomes. Nine other trials^{86, 87, 90, 92, 93, 95, 101, 102, 104} reported comparisons between study groups in the rates of induction of labor, Cesarean deliveries, transfusions, or neonatal admissions, finding no significant differences, with the exception of one study⁸⁶ finding a greater number of Cesarean Sections in the placebo group compared with those taking aspirin, and two studies finding significantly more neonatal admissions in the placebo groups.^{92, 102}

Treatment Adherence and Reasons for Trial Withdrawals

Participants withdrew from treatment for a variety of reasons. It was common for participants to withdraw due to nonmedical reasons, including relocating, changing their minds about trial participation, or noncompliance with treatment. Adherence, when reported, was generally high, although was reported in a variety of ways. A common reporting measure of adherence in the trials was the number of women who were found to be taking at least 80 percent of their treatment medication; this number ranged from 72 to 88 percent. The mean compliance index, as well as the median number of pills taken based on pill counts was also reported in select trials and ranged from 94 to 97 percent. Medical reasons for withdrawal included concerns regarding asthma attacks, increased bleeding time, increased activity of aspartate amino transferase in serum, urticaria, and epigastric pain. Women also withdrew from trials after miscarriage or the termination of pregnancy.

KQ3a. Do the Harms of Aspirin Use to Prevent Preeclampsia Vary by Subpopulations Defined by Personal Characteristics or Preeclampsia Risk Factors?

We tested for differences in placental abruption or postpartum hemorrhage by aspirin dosage and timing among the included trials using meta-regression. No differences in harms related to the dosage of aspirin were seen in comparisons that included all trials included for these harms (data not shown).

When comparing the effect estimates between studies of average-risk and increased-risk study populations, there were no statistically significant differences in estimates of placental abruption (p=0.41), postpartum hemorrhage (p=0.12), or intracranial fetal bleeding (p=0.34), although the

effect estimates in the increased-risk population studies were closer to 1 in all cases (**Figure 9, Figure 10, Figure 11**).

Chapter 4. Discussion

Summary of Evidence

We identified three new trials eligible for this updated systematic review. The summaries of evidence and evaluations of the overall strength of evidence for each Key Question are provided in **Table 6**. There was reasonably consistent and precise evidence of effectiveness of daily low dose aspirin for prevention of fetal growth restriction and preterm birth among individuals at increased risk of preeclampsia. Preventing these outcomes has long-term implications for child and adult health. Newer evidence from a large trial contributed to greater precision for estimating a statistically significant association between aspirin and the prevention of perinatal mortality than in the 2014 review. Data from meta-analyses suggest that use of prophylactic low dose aspirin in women at increased risk for preeclampsia results in a 15 to 20 percent reduced risk of the disease and also for complications that lead to perinatal morbidity and mortality. The incidence of preeclampsia for participants assigned to placebo groups ranged widely, and this contributes to the range of absolute risk reductions related to disease incidence. The two largest trials, CLASP and MFMU, reported similarly modest, statistically insignificant risk reductions of approximately 10 percent with absolute risk reductions near 2 percent. The more recent ASPRE trial obtained larger and statistically significant risk reductions, with absolute reductions for the incidence of preeclampsia closer to 2 percent and for preterm birth over 3 percent.

We estimated the number needed to treat (NNT) for preeclampsia, SGA/IUGR, and preterm birth using a range of values of incidence reported for the outcomes in the included evidence. The NNTs calculated from the midrange of study reported incidence rates for preterm birth and SGA/IUGR (17%) were similar, both around 30, but with wide confidence intervals (SGA/IUGR NNT = 33, 95% CI 18, 588 and preterm NNT= 29, 95% CI 18, 118) (**Table 7**). The NNT was estimated to be 20 or fewer at higher assumed levels of disease incidence for these outcomes, while at the lowest levels of risk, the NNT was 79 for SGA/IUGR and 50 for preterm birth. We estimated the NNT for preeclampsia prevention to be 74 (95% CI 44, 222) when the incidence of preeclampsia is nine percent, over twice the population average, and 51 (95% CI 31, 154) when the incidence is thirteen percent, approximately three times the population average.

Because aspirin use is known to affect pathways involving the inhibition of platelet aggregation, maternal and perinatal bleeding outcomes are important potential harms to consider. The addition of three new trials in the analysis of potential aspirin harms did not alter the previous null findings for the most consistently reported bleeding outcomes. Notably, however, the largest new trial, ASPRE, did not report postpartum hemorrhage outcomes, which limits the information available for assessing potential bleeding harms with a higher dosage than most trials (150mg). Bleeding outcomes, maternal hemorrhage and fetal intracranial bleeding (intraventricular and cerebral hemorrhage) were reported most consistently across all of the included trials. Given the lack of any statistically significant evidence of harms associated with daily low-dose aspirin use in pregnancy, we did not compute numbers needed to harm.

Maternal hemorrhage is the leading cause of severe maternal morbidity in the United States.¹¹⁷ Data from the U.S.-based National inpatient sample indicate the incidence of postpartum

hemorrhage has been increasing.¹¹⁸ From 1993 to 2014, the number of delivery hospitalizations affected by postpartum hemorrhage requiring blood transfusion increased from 7.9/10,000 to 39.7/10,000. In 2009, postpartum hemorrhage accounted for slightly more than 10 percent of maternal deaths (1.7 per 100,000 live births).^{7, 117, 119, 120} Having preeclampsia may increase a patient's risk for postpartum hemorrhage by various mechanisms, including prolonged induction of labor or coagulation abnormalities in patients with HELLP syndrome.^{117, 121} The effect of aspirin prophylaxis during pregnancy on postpartum hemorrhage risk, defined in most studies as blood loss of 500 mL or more, was reported in eleven of the trials in this review, comprising results from over 23,000 participants. There was no increased risk of postpartum hemorrhage associated with aspirin use in trial populations at average or increased risk for preeclampsia. This null finding was accompanied by a reasonably precise confidence interval ranging from three percent reduced risk to nine percent increased risk, and no evidence of statistical heterogeneity. Postpartum blood loss may not be accurately measured in clinical practice, and the clinical significance of the 500mL threshold has been questioned, particularly for patients who receive cesarean section procedures.¹²² A direct consequence of increased blood loss, such as incidence of maternal transfusion, would be a more informative outcome. The absence of differences in the available bleeding related outcomes, however, is reassuring for the daily regimen of 150mg or less of aspirin in pregnancy. The possibility of clinically significant differences for more rare, serious bleeding complications cannot be entirely ruled out, but because preeclampsia increases hemorrhage risk, effective aspirin prophylaxis could also reduce the occurrence of this harm.

Placental abruption, defined as premature separation of the placenta before delivery¹²³ is a condition which may be accompanied by maternal hemorrhage, disruption of blood flow to the fetus, need for emergent delivery, and/or stillbirth. The diagnosis of abruption is primarily made clinically and is less common than either preeclampsia or preterm birth, with national and international population-based studies suggesting abruption occurs in about 0.5 to 1.4 percent of pregnancies.¹⁰⁷⁻¹¹¹ Placental abruption was a reported outcome in 13 trials, including 5 among average-risk individuals, comprising data on over 25,000 trial participants. The null finding for this intermediate outcome was less precisely estimated, and less consistent, than for postpartum hemorrhage, however, the lack of significant effect is consistent with the null findings for health outcomes that may result from placental abruption, including maternal and fetal hemorrhage and death. For less consistently reported harms such as congenital malformations, there was no evidence of an association with aspirin use, but the power to evaluate group differences for these outcomes was limited by their rarity.

Intraventricular hemorrhage (fetal intracranial bleeding) is the most common neurologic complication of prematurity, with risk for this outcome increasing with the degree of fetal prematurity at birth.¹²⁴ In a meta-analysis of 9 trials, the incidence of fetal intracranial bleeding did not differ between aspirin and placebo groups. The pooled effect confidence interval, which ranged from a 40 percent risk reduction to a 35 percent increase, was imprecise because the outcome was rare, however, the effect was in the direction of a reduced risk of fetal intracranial bleeding associated with prophylactic low dose aspirin, which parallels the observed reduction in risk of preterm birth.

Overall, meta-regression analyses did not support notable differences in effects for different study designs or population characteristics. Some benefit was observed across a range of aspirin

dosages and timing, risk assessment approaches, and enrolled populations. Although there is some evidence for a greater benefit for perinatal health outcomes when aspirin is started earlier in pregnancy, results were inconsistent across outcomes, and comparisons may be confounded by other shared study features. There were too few studies included in the review to include multiple covariates in meta-regression, as would be needed to identify independent associations adjusted for other factors. In addition, some aspirin protocol features were too uncommon to compare across studies.

Comparison of Our Results With Findings From Other Systematic Reviews

The results of this review are consistent with findings from other systematic reviews including a recent Cochrane Collaboration review¹²² and an older individual participant meta-analysis conducted by the Paris Collaborative Group (Paris IPD-MA) with trials published before the year 2006.^{60, 125-128} Ten of the RCTs included in this review^{88, 90, 91, 94, 96-99, 102, 106} were included in the Paris IPD-MA.⁶⁰ The Cochrane review included 77 trials conducted with 40,249 participants and their babies and incorporated available Paris IPD-MA data when available, pooling it with study level trial results that were more recently published. The Paris IPD-MA and Cochrane review both included trials of other anti-platelet medication, as well as aspirin combined with other medications. Despite differences in scope and approach, the Cochrane review results were mostly consistent with the findings of our review, with statistically significant benefits for prevention of preeclampsia, perinatal mortality, preterm birth, and small for gestational age, and null, less precisely estimated effects for risks of harm. There also were not statistically significant differences in aspirin effectiveness at different levels of preeclampsia risk in interaction tests conducted in the IPD-MA or the Cochrane review. The pooled effect magnitudes differed for some outcomes compared with our review findings. The most disparate result was for preterm birth, with the Cochrane analyses estimating a more modest statistically significant 9 percent risk reduction compared to the 19 percent risk reduction we estimated. The Cochrane review differed from ours in terms of included study design characteristics (e.g., included settings and medication regimens), preeclampsia risk levels (e.g., the Cochrane review included lower risk populations), risk of bias exclusions, and statistical methods (i.e., Cochrane review used fixed effects meta-analysis and incorporated individual level trial data). Likely a combination of these differences contributed to the differences in effect magnitudes, but the consistency of the finding of a beneficial effect of aspirin across important outcomes among all three reviews speaks to the strength of the evidence of aspirin for preeclampsia prevention in the large body of studies that has accrued to date.

The Cochrane review and Paris IPD-MA reported findings that were consistent with our review with respect to risks of harms associated with daily low dose aspirin use in pregnancy. None of the three reviews found statistically significant differences between groups in rates of postpartum hemorrhage ($\geq 500\text{mL}$), placental abruption, fetal intraventricular hemorrhage, or other neonatal bleeding harms. In the Paris IDP-MA, there was a trend on the margin of statistical significance suggesting the possibility of higher risk of postpartum hemorrhage, based on evidence from 16 trials (RR 1.06, 95%CI 1.00, 1.13); the Cochrane review, which added results from 3 additional trials, showed a similar trend that was also not statistically significant. These results are difficult

to interpret given the problems with the outcome definition and clinical significance of postpartum hemorrhage already discussed. Additionally, differences in the review scope, such as the inclusion of other antiplatelet medications and combined antithrombotic therapies in the other two systematic reviews could be associated with different risks for hemorrhage than aspirin alone. Overall, the large number of participants in trials evaluated in these reviews and ours do not provide a clear signal suggesting bleeding harms.

Consistent with our results, neither the Cochrane review nor the Paris IPD-MA found any definitive subgroup differences. As noted, the Paris IPD-MA results do not include evidence from more recent trials and was scoped more broadly than our current review.⁶⁰ Nevertheless, due to the strengths of an individual participant level of analysis for defining outcomes and assessing subgroup differences, the detailed findings warrant additional consideration. There was no evidence of subgroup differences tested with the Paris IPD-MA in effects by aspirin dosage, the timing of initiation of aspirin therapy, or according to characteristics and preeclampsia risks for study participants. The study found no difference in effectiveness for participants starting before 20 weeks of gestation compared to those starting treatment 20 weeks or later in pregnancy ($p=0.24$). Similarly, a secondary analysis of the data found no difference in effectiveness for preeclampsia prevention or for prevention of perinatal mortality when comparing treatment starting at less than 16 weeks of gestation to later initiation ($p=0.98$ and $p=0.08$, respectively).¹²⁶ The study also tested differences in intervention effects on preeclampsia prevention by aspirin dosage (<75 mg versus >75 mg) among trials that used aspirin only (excluding studies with other anti-platelet medications). The interaction test was not statistically significant ($p=0.23$). A secondary analysis comparing effects for dosages of ≤ 81 mg compared with >81 mg doses suggest a possible trend toward greater risk reduction at higher aspirin doses (<81 mg: RR = 0.92, 95%CI 0.85, 0.99; >81 mg: RR = 0.74, 95%CI 0.60, 0.92), but the estimated effect confidence intervals overlap and a test for interaction was not reported.¹²⁸

Overall, there is consistency across all major reviews of the accumulated evidence in finding that aspirin is effective for preventing preeclampsia and related outcomes, with no notable harms, and little indication of differences in effectiveness between particular regimens or risk populations. In general, there are no clear differences in effectiveness for different protocols or study populations supported by our review or other well-conducted comprehensive reviews on the topic.

Limitations

Exclusions for Publication Language and Study Setting

Our search was limited to English language literature, and only trials conducted in settings with very high Human Development Index scores were included. Large trials conducted in other settings could provide additional relevant information, but other reviews without this exclusion have not found substantively different results.^{60, 122} A large, recently published multicountry trial enrolled average-risk nulliparous participants from seven community sites in the Democratic Republic of Congo, Guatemala, Kenya, Pakistan, and Zambia from March 2016 to June 2018 and randomized 11,976 pregnant individuals to 81mg of aspirin or placebo between 6/0 weeks

and 13/6 weeks of gestation.⁶² Had we not limited our review by setting, this trial would have contributed to our analysis of Key Question 3 (evaluating harms) and would not have changed our main conclusions: the trial did not find any study group differences in a large set of reported adverse events and potential harms, including postpartum hemorrhage and congenital malformations.

Small-Study Effects

We found evidence of small-study effects for several outcomes. This suggests that smaller null or negative studies may not be present in the published literature or that smaller trials have systematic differences in conduct and methods than larger trials, skewing pooled results toward an overestimation of effectiveness. We conducted sensitivity analyses using fixed effects models, given the propensity for random effects models to amplify small-study effects. These analyses demonstrated small differences in the magnitude of effects for various outcomes, though all effects remained statistically significant. Different statistical approaches for estimating pooled effects and testing subgroup comparisons, however, cannot fully address some limitations in the data. For example, the two largest studies used the same low aspirin dosage (60mg) and enrolled the majority of their study participants after 16 weeks of gestation. The MFMU trial additionally instructed participants to cease aspirin use if preeclampsia developed. Thus, the small-study effects seen in the evidence cannot be entirely disentangled from other features of the included studies. It is not possible to determine which factors – those external to the evidence such as publication bias, or those specific to the design of the included studies – are responsible for small-study effects, and both types of factors could contribute.

Confounding of Aspirin Initiation Timing Effects

The most recent Cochrane review¹²² and the Paris IPD-MA⁶⁰ did not find evidence of differences in aspirin effectiveness associated with the timing of aspirin use during pregnancy when analyzing effects with initiation before 20 weeks of gestation compared to later start times using study level data and individual participant data. Our subgroup findings agree with and were similarly interpreted. Other systematic reviews focused on selected subsets of the available trial evidence have reported significant differences in effectiveness related to the timing of aspirin initiation using a cutpoint of 16 weeks of gestation. This could be due to problems with subgroup comparisons at the study level, which are subject to more serious bias and confounding than individual level comparisons. A secondary analysis of the IPD-MA data examining effects for aspirin initiated before 16 weeks of gestation or started later also did not show a statistically significant differences in the effects.¹²⁶ Given the inconsistency in evidence for timing effects across outcomes and the potential for confounding to influence subgroup comparisons, there is not clear evidence that aspirin is more effective when aspirin is started before 16 weeks of gestation. This was true in our current review of aspirin effectiveness focused on increased-risk populations, and in the Cochrane review that includes a broader literature along with results from the Paris IPD-MA.¹²² Further research is needed to examine the possibility that earlier initiation may lead to greater benefit for some individuals, but it appears that even those presenting for prenatal care beyond 16 weeks can derive a benefit from daily low-dose aspirin.

Observational Evidence Exclusions

The 2014 USPSTF review included two observational cohort studies that were not included in this update. These studies examined any aspirin use during pregnancy and reported null findings for the potentially harmful outcomes considered, miscarriage and cryptorchidism.^{107, 108} The current review adopted a narrower approach to assessing observational studies for inclusion, given the size and maturity of the trial evidence on the topic, and the unique aspirin protocol used in trials that is not found in large observational studies examining outcomes associated with aspirin use at any time or dosage during pregnancy for a range of possible indications. We would have included any observational cohort studies that reported harms outcomes among women taking daily aspirin for preeclampsia prevention, but none were identified in our literature search.

A large body of observational evidence has examined whether aspirin and nonsteroidal anti-inflammatory drugs increase the risk of birth defects. Findings are somewhat mixed; most studies report no association or unclear results,¹²⁹⁻¹³² but there are important limitations to the literature. In studies of aspirin for preeclampsia prevention, a low dosage is initiated after the first trimester, when embryogenesis is complete. Therefore, some types of birth defects would not be expected to occur. Most observational studies cannot adjust for the timing, dosage, or indication for aspirin exposure. Ascertainment bias and unmeasured confounding also limit conclusions that can be drawn even from large, well conducted case-control studies.¹³¹ From the observational evidence, there are not clear or consistent associations found between aspirin and birth defects. If present, such effects would be very small in order to have escaped detection in the observational and trial data available on this topic.

Adherence

Our review reported and meta-analyzed results only from intention to treat analyses, in accordance with methodological guidance for conducting systematic reviews and best practices for interpreting trial evidence.^{83, 133} There is evidence in the literature from secondary analyses of trials that differences in adherence to daily aspirin use may contribute to the differences in effectiveness observed in the trial literature, and that efficacy of aspirin use is considerably higher than effectiveness.^{58, 134} We did not analyze data from trials based on adherence to treatment, and from a pragmatic standpoint, the estimates we provide in this review are more representative of the effects that might be expected in clinical populations with varying levels of adherence to a recommended preventive intervention. The MFMU trial, which reported modest null effects of prophylaxis, reported that 80 percent of participants reported taking at least 80 percent of their aspirin tablets, and found no differences in effectiveness based on self-reported adherence.⁹⁸ However, there is evidence from a multicenter observational cohort study that adherence to aspirin (100mg or 150mg) greater than 90 percent (assessed with blood test) compared with any lower level of adherence resulted in statistically significant differences in its effectiveness for preeclampsia and perinatal health outcomes.¹³⁵ An accompanying editorial noted that adherence could be affected by the methods a trial takes to determine risk and to assess medication adherence, and these adherence differences could account for differences in effect magnitudes seen across aspirin trials.¹³⁶ Similarly, there is evidence that adherence to aspirin prophylaxis may be relatively low in pregnancy, with somewhat higher levels for those

concerned about their risk for preeclampsia.¹³⁷ These studies highlight the potential importance of further research to understand the importance of adherence for obtaining preventive benefits.

Identifying Patients at Increased Risk for Preeclampsia

Currently, there is limited evidence about the best approach for identifying women at risk of preeclampsia.^{138, 139} Trials varied in their approaches for identifying participants and in the exclusion criteria they used. A little over half of the included trials used multiple risk factors to identify women at risk for preeclampsia, and of the trials that relied on a single risk factor, only two relied on the same one (abnormal doppler ultrasound readings).^{89, 102} Consequently, baseline characteristics of the study populations varied significantly, resulting in a broad range of preeclampsia incidence rates in the control groups (4 - 30%) (**Table 4**). Some of the tests used to identify women at risk of developing preeclampsia in the older trials were conducted in the second trimester of pregnancy (some as late as 26-28 weeks of gestation) and are not currently used in clinical practice (i.e., hemoglobin >13.2 g/dL, maternal roll over test resulting in rise of diastolic blood pressure >15 mg Hg above baseline, and increased blood pressure response to angiotensin II infusion).^{86, 88, 95}

Study level subgroup comparisons tested in this review did not find differences in aspirin effectiveness based on the approach taken for identifying women at increased risk of preeclampsia. While studies that used clinical risk factors in conjunction with clinical tests or imaging consistently recruited populations with high incidence of preeclampsia (18 and 23%), several studies^{88, 103} using clinical risk factors alone achieved similarly high levels of preeclampsia incidence ranging from 8 percent to 20 percent (**Table 4**).^{56, 92-94, 97, 98, 100} Subgroup analyses did not show any differences in the effectiveness of aspirin between studies that used clinical history risk factors alone to identify women at increased risk compared to those that incorporated clinical tests or imaging, nor did the effectiveness of aspirin vary by the incidence of preeclampsia in the study control group ($\leq 12\%$ vs $>12\%$) (**Table 4**). However, subgroup comparisons were limited by the diversity of approaches taken and the inability to account for potential confounding from other study characteristics (e.g., aspirin protocol) with the available trials.

The most recently published RCT of aspirin for prevention of preeclampsia, the ASPRE trial, was the only trial to use a previously developed, externally validated risk prediction model for predicting preeclampsia.¹⁴⁰ The model was used to assess risk of preeclampsia leading to preterm birth in participants with singleton pregnancies at 11 to 12 weeks of gestation. The prediction algorithm draws on Bayesian statistics to first estimate risk based on several clinical history characteristics; weight, height, maternal age, race/ethnicity, previous preeclampsia, family history of preeclampsia, chronic hypertension, diabetes type 1 or 2, systemic lupus erythematosus or antiphospholipid syndrome, and conception by *in vitro* fertilization. This was then supplemented with serum biomarker tests (pregnancy associated plasma protein-A and placental growth factor), mean arterial pressure, and Doppler ultrasound readings of the uterine artery pulsatility index. Using this algorithm, the trial identified a study population where 11 percent of control group participants developed preeclampsia, about 3 times the incidence in the general population. It is not clear whether the greater effectiveness of aspirin seen in the ASPRE

trial relative to other large trials can be attributed to the risk assessment approach, and how much is due to other differences, such as the higher aspirin dosage, or the early initiation of aspirin at 12 to 13 weeks of gestation and continuation until 36 weeks or onset of labor. Whether the algorithm identifies a population particularly responsive to aspirin cannot be determined without further comparative research.¹⁴⁰

The MFMU trial conducted stratified enrollment and analyses to test whether aspirin was more effective for certain risk groups relative to others. The trial defined four risk categories: those at risk of diabetes, chronic hypertension, and multifetal gestation, and those with a previous pregnancy affected by preeclampsia. There were no statistically significant effects of aspirin for preventing preeclampsia or other health outcomes in the trial overall or within risk category subgroups. All risk groups except the chronic hypertension category had similarly modest reductions in preeclampsia risk (approximately 10%). In the chronic hypertension risk category, more participants in the aspirin treatment group developed preeclampsia than in the placebo group, but the estimate was imprecise and comparisons between groups were statistically underpowered.

Despite ongoing efforts to develop a screening test or prediction algorithm that can be used in early pregnancy to identify individuals who are likely to develop preeclampsia, clinical history taking remains a practical and commonly-used approach that can select populations with higher incidence of preeclampsia than the population average. Aspirin was effective across a range of populations at different levels of risk who were selected based on a variety of approaches. Risk prediction models that include uterine artery Doppler ultrasound readings and biochemical markers from serum testing alongside patient characteristics are more resource-intensive and difficult to clinically implement but may hold potential. The most widely tested prediction models have yet to be externally validated, and require further evaluation to determine their clinical effectiveness and value to support widespread use across a range of populations.^{1, 39, 65}

Some researchers have advocated for broader clinical uptake of the algorithm used in the ASPRE trial for selecting patients to aspirin prophylaxis,¹⁴¹ including a recommendation from the International Federation of Gynecology and Obstetrics (FIGO) (**Table 1**) (co-authored by the model developers). Others have questioned this based on practical shortcomings and knowledge gaps, highlighting its limited suitability to general clinical practice for now.^{1, 39, 65, 138, 142} Many pregnant individuals would not be eligible for screening, including those entering prenatal care beyond 12 weeks of gestation or who are pregnant with twins or more. It is also unclear what level and attributes of test performance are most important assigning aspirin prophylaxis, given the low risk of the intervention. For low dose aspirin prophylaxis, optimizing specificity rather than sensitivity (higher false positive rate) might be reasonable. Validated prediction tools with strong test performance attributes would be very valuable for improving prenatal care and early intervention if they were able to identify those who are very likely to develop preeclampsia, but may be less important for identifying a clinical population in routine care that might benefit from aspirin prophylaxis. A review comparing the test performance attributes of development and validation models for the predictive model found significant differences in accuracy depending on the cohort used, leading the authors to conclude that more effort should be dedicated to refinement and validation of models as well as studies evaluating whether model use improves clinical care and outcomes.¹⁴³

Comparisons of clinical risk assessment approaches recommended by ACOG, NICE, and USPSTF to formal prediction algorithms for identifying populations at increased risk that could benefit from aspirin to prevent preeclampsia have highlighted differences in test performance characteristics. Not surprisingly, these comparisons show that prediction models designed to optimize accuracy have greater predictive value. As others have noted,¹⁴² however, while accurate prediction of preterm preeclampsia in singleton pregnancy is important for many purposes, lengthy clinical risk assessment procedures may not represent the most efficient, practical, or effective way to ensure that recommendations to use aspirin during pregnancy reach the broad population of individuals that could potentially benefit from use. Further research is needed to determine whether there is a discernable difference between aspirin effectiveness for individuals identified using a prediction model and those identified using clinical history risk factors in routine prenatal. Implementation trials comparing these strategies in real-world settings would be particularly valuable, especially if designed to address questions related to aspirin timing as well. It would be especially important to further strengthen approaches for identifying individuals that are most likely to manifest severe features of preeclampsia that lead to more serious morbidity and mortality.

In addition to the trial evidence included in this review, several reviews and large cohort studies provide evidence on important risk factors for preeclampsia. These were cited in the previous USPSTF recommendation statement⁶⁸ and were referenced in the recent ACOG recommendation for aspirin to prevent preeclampsia.^{65, 144} A similar set of risk factors were cited in the updated National Institute for Health and Clinical Excellence recommendation for the United Kingdom (**Table 1**). These include several risk factors known to be independently associated with the highest likelihoods of developing preeclampsia: history of preeclampsia, multifetal gestation, chronic hypertension, and type 1 or type 2 diabetes. Other, more moderate risk factors are associated with an increased risk, such as maternal age over 35 and nulliparity, but are inconsistently or only modestly independently associated with preeclampsia risk. More than one moderate risk factor, occurring in the same individual, however, would be expected to identify increased risk for preeclampsia similar to levels observed in the control groups of the trials that established intervention effectiveness.

There has been no change in the main clinical risk factors associated with preeclampsia that would be reasonable to consider for identifying similar-risk populations to those included in the 2014 USPSTF review. Clinical risk factors included in the risk prediction model used in the ASPRE trial were similar to those recommended by NICE, ACOG, and the USPSTF, with the exception of *in vitro* conception. This clinical history could be incorporated into current clinical assessment practices, although only a small fraction of participants in aspirin trials had this risk factor. There are also other risk factors not included in the current guidelines that may also be associated with a moderate risk for preeclampsia and warrant further research both in terms of establishing the potential association and evaluating whether aspirin prophylaxis would be beneficial. These include high altitude residence, exposure to air pollution,¹⁴⁵ sleep disordered breathing during pregnancy,^{146, 147} and migraine headaches.¹⁴⁸ **Table 8** provides a list of established clinical history risk factors that can be used alongside clinical judgment of risk to identify increased-risk populations that are likely to approximate the population included in the trials on aspirin prophylaxis reviewed here.

Having had a previous pregnancy complicated by a hypertensive disorder is a consistent, well-established risk factor for preeclampsia, and there is some evidence that this risk factor identifies a population that may be particularly responsive to aspirin prophylaxis.⁶⁰ Additionally, a history of preeclampsia may be associated with an increased risk of having an SGA infant, even if preeclampsia does not occur, highlighting the importance of this risk factor for identifying individuals that could benefit from aspirin prophylaxis.¹⁴⁹ Of course, for nulliparous individuals, this important pregnancy history risk factor is unavailable. First pregnancies are at somewhat greater risk for preeclampsia than later pregnancies; as a result, the population-attributable risk is high for nulliparous patients.³⁷ Studies we reviewed that enrolled general nulliparous populations without additional risk factors had incidence of preeclampsia below or just above the population average in their control groups.^{90, 91, 99, 101, 104} Efforts to develop prediction algorithms focused on nulliparous populations have had limited success,^{138, 150-152} and generally have not performed better than recommended approaches to clinical risk assessment. The addition of serum markers and imaging findings may improve performance,^{38, 153} but further validation in nulliparous populations and comparison to other risk assessment approaches is needed. Research to inform risk assessment for those having a first pregnancy is needed to ensure that those who would benefit from aspirin prophylaxis are identified.

Finally, there is a developing discussion in the literature¹⁵⁴⁻¹⁵⁶ as to whether low dose aspirin prophylaxis should be considered for all pregnant women given challenges predicting who will develop preeclampsia, particularly among nulliparous individuals, and the absence of any clear evidence of harms.^{155, 157} The inclusion of low-risk populations in systematic reviews tends to result in more modest effect estimates compared to reviews focused on higher risk populations, but in the most recent Cochrane review, there was not a statistically significant difference in aspirin effectiveness between different levels of risk.¹²² We also did not find differences in the main harms outcomes we evaluated when comparing the effects between studies conducted among average-risk (mainly nulliparous) populations and studies with increased-risk study populations. With additional research, it may become reasonable to consider a shift away from identifying increased-risk populations toward identifying low-risk populations that are unlikely to derive any benefit from aspirin prophylaxis for preeclampsia prevention. This will require more evidence on the long-term safety and benefits of aspirin prophylaxis to support intervention involving larger numbers needed to treat.

Evidence Gaps and Future Research

In addition to the gaps around identifying patients at increased risk for preeclampsia and most likely to have more severe complications (detailed above), there are a number of other evidence gaps that call for future research. The literature we reviewed is limited for the purposes of identifying the specific aspirin protocol (dosage, timing, continuation, time of day) that is likely to have the greatest benefit, and the risk population that is most likely to benefit. In addition, the long-term risks and benefits of low dose aspirin for the mother or her child are uncertain, as the included trials have not reported long-term outcomes. The studies we reviewed varied considerably on these characteristics, in addition to having differences in excluded populations and sample size. No studies reported head-to-head comparisons of different protocols or risk assessment strategies, and only one trial was designed to compare effectiveness for different risk

populations, and it was underpowered for drawing conclusions. Interaction tests for subgroup differences within trials were uncommon, and, when available, did not generally identify differences in effectiveness by the factors examined. Meta-analysis results for some outcomes, and especially for making subgroup comparisons, were limited by low numbers of included studies and low event rates for some outcomes. Comparisons between studies in meta-regression were generally not statistically significant. When differences were seen, as for the timing of aspirin initiation on preterm birth prevention, the limited number of studies compared and potential for confounding prevented us from drawing strong conclusions. For example, only one study evaluated differences in the action of aspirin depending on the time of day it was taken,⁵⁶ providing limited evidence that effectiveness may be greater when aspirin is taken at least 8 hours after waking or at bedtime. Further research is needed to explore this possibility. Overall, the low to moderate statistical heterogeneity and consistency of effects across a range of outcomes supports the conclusion of this review, that low dose aspirin is effective for preventing preeclampsia and related perinatal morbidity and mortality for individuals at increased risk for preeclampsia. More research will be needed to determine the optimal aspirin protocol and identify key populations most likely to obtain benefits.

Applicability of Evidence and Health Equity

The included studies in this review ranged from explanatory to pragmatic in their design features, but were all randomized trials and therefore represent a subset of the larger population of pregnant individuals who were willing to undergo randomization to daily medication use during pregnancy. The effectiveness of the intervention, due to differences in adherence or the health status between participants and the general population, may not match what has been found in clinical trials.

The majority of participants in the included trials were White, limiting the generalizability of results to racially and ethnically diverse populations at risk for preeclampsia. Only the two United States trials, the MFMU trial,⁹⁸ which recruited women at increased risk for developing preeclampsia, and the Sibai trial,⁹¹ in average-risk nulliparous individuals, enrolled substantial numbers of Black and Hispanic study participants; none of the included studies enrolled significant numbers of other minority racial or ethnic groups. Thus, the evidence is limited for assessing whether there could be racial or ethnic differences in the effectiveness or potential harms of aspirin in pregnancy. This is a troubling in light of the greater morbidity from preeclampsia experienced by Black women in the United States compared to other racial and ethnic groups.¹¹ More than half the participants enrolled in MFMU were Black, and the trial did not find a difference in aspirin effectiveness by race/ethnicity (coded as nonWhite versus White), but the low magnitude of effects in this trial limited the comparison. Racial/ethnic differences in aspirin effectiveness or harms could exist, owing to the prevalence of different risk factors in different racial groups, environment/gene interactions, or other factors. Such differences could contribute to heterogeneity in the effect magnitudes observed in the evidence we reviewed. Given the troubling disparities in maternal health and birth outcomes documented among Black women in the United States,^{11, 158, 159} research is urgently needed to address this evidence gap.

A retrospective cohort study of births among Hispanic and African-American individuals with a history of preeclampsia provides evidence that benefits seen in the trials included in this review can be obtained in routine care settings. The study compared preeclampsia recurrence before and after the release of the 2014 USPSTF recommendation for aspirin prophylaxis.⁶³ Adjusted analyses indicated 30 percent reduced recurrence of preeclampsia, despite an increase in rates of pregestational diabetes over the study period (a significant risk factor for preeclampsia). Although the observational study design is subject to risk of bias from historical trends and other unmeasured confounding, it supports the applicability of trial evidence synthesized in this review on to routine care provided in clinical settings serving diverse populations. Nevertheless, further research on implementation will be needed to ensure that effective interventions are equitably distributed, as health disparities in preeclampsia may be further exacerbated by unequal access high quality antenatal health care where effective interventions are implemented in practice.

Chronic and Gestational Hypertension

A recent study of over 44 million deliveries in the Nationwide Inpatient Sample dataset found that the incidence of hypertensive disorders in pregnancy increased from 8.4 percent in 2004 to 10.9 percent in 2014, with the largest increases seen in the categories of gestational hypertension and chronic hypertension. Of note, women with chronic hypertension without preeclampsia had higher risks of cardiovascular events (myocardial infarction and arrhythmia) than women with preeclampsia without hypertension. Women with chronic hypertension as well as preeclampsia had the highest risk of adverse outcomes (including much higher risks of preterm birth, stillbirth, and stroke).¹⁶⁰

The trials included in our review also showed higher preeclampsia incidence for participants entering the study with elevated blood pressure than those without elevated blood pressure, when this was reported. Participants with chronic hypertension enrolled in the MFMU trial had the highest rates of preeclampsia, and nearly the same proportion of participants developed preeclampsia in the aspirin group as in the placebo group, 25 and 26 percent, respectively. A subgroup analysis in the ASPRE trial suggested a higher risk of preterm birth with preeclampsia among women with chronic hypertension in the aspirin group than in similar women assigned to placebo:⁵⁸ a test for an interaction between hypertension status and treatment group was on the margin of statistical significance. These two trials suggest that aspirin may not be effective for individuals with hypertension diagnosed before pregnancy, but the results are inconclusive. In addition, the trial literature we reviewed was limited for determining whether a diagnosis of chronic hypertension using the AHA's revised lower systolic and diastolic blood pressure cut points of 130 and 80, respectively, should be incorporated when identifying women at risk of preeclampsia.¹⁶¹ In a systematic review of cohort studies evaluating risk factors for preeclampsia, a systolic blood pressure at the initial visit of ≥ 130 mm Hg vs < 130 was associated with an unadjusted relative risk of preeclampsia of 2.37 (1.78 to 3.15) and a diastolic blood pressure of ≥ 80 versus < 80 was associated with a unadjusted relative risk of preeclampsia of 1.38 (1.01 to 1.87).¹⁶²

There is limited, but suggestive evidence from secondary analysis of data from the MFMU trial that for individuals with stage 1 hypertension at baseline (less than 26 weeks of gestation) that were not previously hypertensive, aspirin may be more beneficial for certain risk groups.¹⁶³ Though the analysis had limitations, it suggested that the presence of stage 1 hypertension before 26 weeks of gestation was associated with later preeclampsia incidence, and that aspirin was more effective for preeclampsia prevention among individuals with this level of hypertension than among those without. A similar analysis of the entire study population reported in the primary MFMU trial publication, however, found similar aspirin effectiveness for participants who were not hypertensive at baseline but had systolic blood pressure of 120 to 134 mmHg and those who were normotensive at study entry (before 26 weeks of gestation).⁹⁸ A trial included for harms in this review, also conducted in racially and ethnically diverse patients recruited from MFMU network⁹¹ sites, found that aspirin may be effective in healthy nullipara with SBP 120-134 mm Hg at baseline, whereas no difference between aspirin and placebo was found for lower baseline blood pressure levels.⁹⁸ Among the risk group in MFMU with chronic hypertension at study entry, who were excluded from the secondary analysis described above, there were more cases of preeclampsia in the aspirin group than the control group.

Taken together, these findings suggest that those with chronic hypertension (now considered stage 2 chronic hypertension) may be less responsive to aspirin prophylaxis, while individuals without a history of chronic hypertension who are newly identified as having stage 1 hypertension early in pregnancy may be more likely to benefit than individuals with no hypertension. Future research is needed to determine whether there is a subgroup of women with pre-existing hypertension, perhaps those with a recent diagnosis at a lower diagnostic cut point, who may benefit from aspirin therapy. The limited data leave it unclear whether the risk of superimposed preeclampsia may be mitigated by aspirin prophylaxis, especially among women with stage 2 hypertension, but no studies have been adequately powered to examine this hypothesis.

Another area of uncertainty is the benefit of aspirin for the prevention of gestational hypertension in the absence of preeclampsia. We attempted to address this question by pooling all cases of gestational hypertension from studies that provided the outcome. Our meta-analysis found that the direction of risk reduction was the same for gestational hypertension as for preeclampsia, but even though effect magnitude was greater, the estimate did not achieve statistical significance. One challenge in assessing gestational hypertension from the available RCT data is that the presence of this condition is not commonly reported in a way that would allow us to identify participants who were diagnosed with preeclampsia after an initial diagnosis of gestational hypertension. Thus, if aspirin reduces the likelihood that gestational hypertension will transition to preeclampsia, there could be more cases of gestational hypertension (rather than preeclampsia) reported in the intervention group. Aspirin might also reduce the risk for gestational hypertension. Given that gestational hypertension and preeclampsia are now treated the same at term (i.e. delivery is recommended at 37 weeks of gestation for both conditions),¹ we conducted an additional exploratory analysis assessing whether aspirin was effective for reducing the risk of hypertensive disorders of pregnancy as a whole in trials reporting adequate data. We found a significant risk reduction in the aspirin condition, supporting the possibility that aspirin may be useful for prevention of both gestational hypertension and preeclampsia. Further confirmatory research is needed.

Obesity

One question for which current trial data are limited is whether the dosing of aspirin for the prevention of preeclampsia should differ for women with a higher body mass index. Only two of the included trials performed analyses stratified by BMI to examine the differential impact of LDA for prevention of preeclampsia (MFMU, 60 mg) and prevention of preeclampsia with preterm birth (ASPRE, 150 mg) and found no significant differences by BMI category. One hypothesis for why the dose of aspirin may need to be different for women with obesity is that the degree of thromboxane inhibition achieved at a given dose may vary by BMI. To examine whether aspirin is associated with less thromboxane inhibition among women with obesity, a subsequent secondary analysis was conducted in a subgroup of 1002 of the original 2539 ppts the MFMU trial.¹⁶⁴ Women in the cohort were stratified by BMI (obese ≥ 30 kg/m² vs nonobese < 30 kg/m²). The primary outcome was rate of complete thromboxane B2 (TXB2) inhibition, defined as undetectable TXB2 levels < 0.01 ng/mL and the secondary outcomes were rates of preeclampsia at any gestational age and preeclampsia with preterm delivery < 37 weeks in women with and without undetectable TXB2 levels. Women with obesity assigned to LDA had statistically significant lower rates of undetectable TXB2 levels in the second trimester (39.4% vs 30.2%; $P = .04$), but not during the third trimester (41.4% vs 32.9%; $P = .12$), and were substantially less likely to have undetectable levels of TXB2 at both time points (30.6% vs 17.7%; $P = .010$). However, there were no differences in rates of preeclampsia in women receiving LDA with undetectable TXB2 levels in the second trimester (17.2% vs 20.8%; $P = .34$), third trimester (16.4% vs 19.7%; $P = .46$), or at both time points (16.5% vs 17.5%; $P = .83$), and there were no differences in rates of preeclampsia with delivery less than 37 weeks in women with undetectable TXB2 levels in the second trimester (9.9% vs 9.2%; $P = .71$), third trimester (4.5% vs 7.4%; $P = .27$), or at both time points (5.1% vs 6.3%; $P = .83$).

In summary, while there appears to be a differential effect of aspirin on thromboxane inhibition among women with obesity, this does not appear to result in a differential impact on outcomes, although trial data examining the effect of BMI are limited.

Identifying Serious Harms

The challenge in identifying potentially serious harms is apparent when considering the outcome of clinically significant postpartum hemorrhage. As noted above, the trial evidence most commonly used a measure of 500mL or more of blood loss to define postpartum hemorrhage, but this number is of questionable clinical significance.¹²² In 2014, after all but two of the studies included in this review were published, the national reVITALize initiative¹⁶⁵ worked to standardize obstetric data definitions and redefined postpartum hemorrhage as a cumulative blood loss greater than or equal to 1,000 mL, or blood loss accompanied by signs or symptoms of hypovolemia within 24 hours after the birth process (including intrapartum loss) regardless of route of delivery.¹⁶⁶ This new definition is believed to be more clinically meaningful and has been adopted by the American College of Obstetrics & Gynecology.¹¹⁷ Future trials may benefit from adoption of an outcome measure consistent with this definition to better assess whether there are meaningful differences in postpartum hemorrhage rates associated with aspirin to

prevent preeclampsia. In the meantime, the possibility of clinically significant differences for more rare, serious implications cannot be entirely ruled out.

Longer-Term Health Outcomes

It is possible that there are longer-term child and adult health implications associated with preeclampsia and/or its prevention with aspirin prophylaxis. Only one included trial reported followup beyond the perinatal period:⁹⁴ a subset of the children of trial participants were assessed at 18 months of age.¹¹² While this study was reassuring with regard to the lack of developmental harms in toddlerhood, there remains a need for longer-term followup, to assess potential harms and potential benefits of intervention. A growing body of research suggests that exposure to preeclampsia in utero may be harmful to future health. Longer-term trial followup could evaluate the possibility that prevention of preeclampsia with aspirin may reduce offspring risk of cardiovascular disease^{167, 168} as well as for neurocognitive conditions such as attention deficit hyperactivity disorder,^{169, 170} autism spectrum disorder, epilepsy, intellectual disability, and vision or hearing loss.¹⁷¹

Followup on trial participants would similarly have the potential to shed light on whether prevention of preeclampsia during pregnancy has long-term health implications for mothers. A number of studies have found associations between preeclampsia and long-term cardiovascular health outcomes.^{172, 173} Estimates suggest a possible doubling or tripling of cardiovascular disease risk in women who have had preeclampsia during any pregnancy,^{174, 175} as well as an increased risk for early stroke (before age 60).¹⁷⁶ Risks may be considerably higher in women who had early-onset preeclampsia.¹⁷⁷ Subclinical brain and endothelial damage may occur from the cardiovascular strain and metabolic effects of preeclampsia.^{173, 178-180} It is unclear whether these changes cause the observed increased long-term cardiovascular disease and mortality risk associated with a history of preeclampsia, or if differences arise from underlying risk profiles among women that potentiate the development of both of these short- and long-term outcomes.¹⁸¹ Further research, including extended followup on aspirin trials, could shed light on the mechanisms and causal pathways between preeclampsia prevention and longer-term health.¹⁷⁸

Conclusions

Few new studies have been published since the previous review conducted for the USPSTF. The addition of a large trial strengthened the evidence for the effectiveness of low dose aspirin for preventing preeclampsia and reducing the risk of SGA/IUGR, preterm birth, and perinatal mortality, across a range of aspirin protocols and increased-risk populations. Data from over 25,000 individuals randomized to daily aspirin use for preeclampsia prevention do not suggest that there are serious harms associated with low dose aspirin use for this purpose during pregnancy. The evidence included in this review cannot rule out rare harms from low dose aspirin use during pregnancy, although findings are reassuringly consistent with the findings of large registry and cohort studies (not included in this review) examining potential harms of aspirin exposure at any gestation, at any dose, and for any indication. The effectiveness of aspirin

is supported across a range of risk assessment strategies, aspirin dosages, and protocols regarding when during pregnancy to begin taking aspirin. Further research is needed to address gaps in the research related to potential variability in the effectiveness of aspirin that could not be resolved from subgroup comparisons in the existing trial evidence on this topic. Even if aspirin has only modest effectiveness for improving pregnancy outcomes, their importance suggests that the intervention could have important implications for population health and well-being across the lifespan.

References

1. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 202: Gestational Hypertension and Preeclampsia. *Obstetrics and gynecology*. 2019;133(1):e1-e25. PMID: 30575675. 10.1097/aog.0000000000003018
2. Hutcheon JA, Lisonkova S, Joseph KS. Epidemiology of pre-eclampsia and the other hypertensive disorders of pregnancy. *Best practice & research Clinical obstetrics & gynaecology*. 2011;25(4):391-403.
3. Trogestad L, Magnus P, Stoltenberg C. Pre-eclampsia: Risk factors and causal models. *Best practice & research Clinical obstetrics & gynaecology*. 2011;25(3):329-42.
4. Sibai BM. The HELLP syndrome (hemolysis, elevated liver enzymes, and low platelets): much ado about nothing? *Am J Obstet Gynecol*. 1990;162(2):311-6.
5. Ghulmiyyah L, Sibai B. Maternal mortality from preeclampsia/eclampsia. *Semin Perinatol*. 2012;36(1):56-9.
6. Mayrink J, Costa ML, Cecatti JG. Preeclampsia in 2018: Revisiting Concepts, Physiopathology, and Prediction. *ScientificWorldJournal*. 2018;2018:6268276. 10.1155/2018/6268276
7. Say L, Chou D, Gemmill A, et al. Global causes of maternal death: a WHO systematic analysis. *The Lancet Global Health*. 2014;2(6):e323-e33. 10.1016/S2214-109X(14)70227-X
8. Khan KS, Wojdyla D, Say L, et al. WHO analysis of causes of maternal death: a systematic review. *Lancet*. 2006;367(9516):1066-74. 10.1016/S0140-6736(06)68397-9
9. Booker WA, Ananth CV, Wright JD, et al. Trends in comorbidity, acuity, and maternal risk associated with preeclampsia across obstetric volume settings. *J Matern Fetal Neonatal Med*. 2019;32(16):2680-7. 10.1080/14767058.2018.1446077
10. Fingar KR M-HI, Ngo-Metzger Q, et al. Delivery Hospitalizations Involving Preeclampsia and Eclampsia, 2005–2014: Statistical Brief #222. Rockville, MD: AHRQ; 2017.
11. Gyamfi-Bannerman C, Pandita A, Miller EC, et al. Preeclampsia outcomes at delivery and race. *J Matern Fetal Neonatal Med*. 2019;1-8. PMID: 30786794. 10.1080/14767058.2019.1581522
12. Petersen EE, Davis NL, Goodman D, et al. Racial/Ethnic Disparities in Pregnancy-Related Deaths - United States, 2007-2016. *MMWR Morb Mortal Wkly Rep*. 2019;68(35):762-5. PMID: 31487273. 10.15585/mmwr.mm6835a3
13. Creanga AA, Syverson C, Seed K, et al. Pregnancy-Related Mortality in the United States, 2011-2013. *Obstetrics and gynecology*. 2017;130(2):366-73. PMID: 28697109. 10.1097/aog.0000000000002114
14. Howell EA. Reducing Disparities in Severe Maternal Morbidity and Mortality. *Clin Obstet Gynecol*. 2018;61(2):387-99. PMID: 29346121. 10.1097/GRF.0000000000000349
15. Centers for Disease Control and Prevention. Causes of pregnancy-related death in the United States: 2011-2016. <https://www.cdc.gov/reproductivehealth/maternal-mortality/pregnancy-mortality-surveillance-system.htm>. Accessed: 2020.
16. Steegers EA, von DP, Duvekot JJ, et al. Pre-eclampsia. *Lancet*. 2010;376(9741):631-44.

17. Stevens W, Shih T, Incerti D, et al. Short-term costs of preeclampsia to the United States health care system. *Am J Obstet Gynecol.* 2017;217(3):237-48 e16. 10.1016/j.ajog.2017.04.032
18. Grobman WA, Bailit JL, Rice MM, et al. Frequency of and factors associated with severe maternal morbidity. *Obstetrics and gynecology.* 2014;123(4):804-10. 10.1097/AOG.0000000000000173
19. Lisonkova S, Joseph KS. Incidence of preeclampsia: risk factors and outcomes associated with early- versus late-onset disease. *Am J Obstet Gynecol.* 2013;209(6):544 e1- e12. 10.1016/j.ajog.2013.08.019
20. Ananth CV, Vintzileos AM. Maternal-fetal conditions necessitating a medical intervention resulting in preterm birth. *Am J Obstet Gynecol.* 2006;195(6):1557-63. 10.1016/j.ajog.2006.05.021
21. Honigberg MC, Zekavat SM, Aragam K, et al. Long-Term Cardiovascular Risk in Women With Hypertension During Pregnancy. *J Am Coll Cardiol.* 2019;74(22):2743-54. 10.1016/j.jacc.2019.09.052
22. Melamed N, Ray JG, Hladunewich M, et al. Gestational hypertension and preeclampsia: are they the same disease? *J Obstet Gynaecol Can.* 2014;36(7):642-7. 10.1016/S1701-2163(15)30545-4
23. Brouwers L, van der Meiden-van Roest AJ, Savelkoul C, et al. Recurrence of pre-eclampsia and the risk of future hypertension and cardiovascular disease: a systematic review and meta-analysis. *BJOG : an international journal of obstetrics and gynaecology.* 2018;125(13):1642-54. 10.1111/1471-0528.15394
24. American College of Obstetricians and Gynecologists T Fo Hi P. Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstetrics and gynecology.* 2013;122(5):1122-31. 10.1097/01.Aog.0000437382.03963.88
25. Behrens I, Basit S, Melbye M, et al. Risk of post-pregnancy hypertension in women with a history of hypertensive disorders of pregnancy: nationwide cohort study. *BMJ (Clinical research ed).* 2017;358:j3078. 10.1136/bmj.j3078
26. Williams D. Long-term complications of preeclampsia. *Semin Nephrol.* 2011;31(1):111-22. 10.1016/j.semnephrol.2010.10.010
27. Roberts JM. The perplexing pregnancy disorder preeclampsia: what next? *Physiol Genomics.* 2018;50(6):459-67. PMID: 29602298. 10.1152/physiolgenomics.00017.2018
28. Redman CW, Sargent IL, Staff AC. IFPA Senior Award Lecture: making sense of pre-eclampsia - two placental causes of preeclampsia? *Placenta.* 2014;35 Suppl:S20-5. 10.1016/j.placenta.2013.12.008
29. Redman CW, Staff AC. Preeclampsia, biomarkers, syncytiotrophoblast stress, and placental capacity. *Am J Obstet Gynecol.* 2015;213(4 Suppl):S9.e1, S9-11. 10.1016/j.ajog.2015.08.003
30. Rana S, Lemoine E, Granger JP, et al. Preeclampsia: Pathophysiology, Challenges, and Perspectives. *Circ Res.* 2019;124(7):1094-112. PMID: 30920918. 10.1161/CIRCRESAHA.118.313276
31. Preeclampsia and eclampsia. https://www.clinicalkey.com/#!/content/clinical_overview/67-s2.0-dcd12356-2aec-47a9-8277-9ed314ce1866. Accessed.

32. Phipps E, Prasanna D, Brima W, et al. Preeclampsia: Updates in Pathogenesis, Definitions, and Guidelines. *Clinical Journal of the American Society of Nephrology*. 2016;11(6):1102-13. 10.2215/cjn.12081115
33. Ayala DE, Hermida RC. Sex differences in the administration-time-dependent effects of low-dose aspirin on ambulatory blood pressure in hypertensive subjects. *Chronobiol Int*. 2010;27:345-62.
34. Esplin MS, Fausett MB, Fraser A, et al. Paternal and maternal components of the predisposition to preeclampsia. *New England Journal of Medicine*. 2001;344(12):867-72.
35. Gabbe SG. *Obstetrics : normal and problem pregnancies*. 2017.
36. Ghosh G, Grewal J, Männistö T, et al. Racial/ethnic differences in pregnancy-related hypertensive disease in nulliparous women. *Ethn Dis*. 2014;24(3):283-9.
37. Bartsch E, Medcalf KE, Park AL, et al. Clinical risk factors for pre-eclampsia determined in early pregnancy: systematic review and meta-analysis of large cohort studies. *BMJ (Clinical research ed)*. 2016;353:i1753. PMID: 27094586.
<https://dx.doi.org/10.1136/bmj.i1753>
38. Wright D, Tan MY, O'Gorman N, et al. Predictive performance of the competing risk model in screening for preeclampsia. *Am J Obstet Gynecol*. 2019;220(2):199.e1-e13. 10.1016/j.ajog.2018.11.1087
39. Henderson JT, Thompson JH, Burda BU, et al. Preeclampsia Screening: Evidence Report and Systematic Review for the US Preventive Services Task Force. *Jama*. 2017;317(16):1668-83. 10.1001/jama.2016.18315
40. Altman D, Carroli G, Duley L, et al. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? The Magpie Trial: a randomised placebo-controlled trial. *Lancet*. 2002;359(9321):1877-90.
41. Magpie Trial Follow-Up Study Collaborative Group. The Magpie Trial: a randomised trial comparing magnesium sulphate with placebo for pre-eclampsia. Outcome for children at 18 months. *BJOG : an international journal of obstetrics and gynaecology*. 2007;114(3):289-99. 10.1111/j.1471-0528.2006.01165.x
42. Group MTF-USC. The Magpie Trial: a randomised trial comparing magnesium sulphate with placebo for pre-eclampsia. Outcome for women at 2 years. *BJOG : an international journal of obstetrics and gynaecology*. 2007;114(3):300-9. 10.1111/j.1471-0528.2006.01166.x
43. Koopmans CM, Bijlenga D, Groen H, et al. Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre, open-label randomised controlled trial. *Lancet*. 2009;374(9694):979-88. 10.1016/s0140-6736(09)60736-4
44. Haapsamo M, Martikainen H, R.,s.,nen J. Low-dose aspirin reduces uteroplacental vascular impedance in early and mid gestation in IVF and ICSI patients: a randomized, placebo-controlled double-blind study. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2008;32:687-93.
45. Hofmeyr GJ, Lawrie TA, Atallah Á N, et al. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *The Cochrane database of systematic reviews*. 2018;10(10):Cd001059. 10.1002/14651858.CD001059.pub5

46. World Health Organization. WHO recommendation: calcium supplementation during pregnancy for prevention of pre-eclampsia and its complications: World Health Organization; 2018.
47. Rumbold A, Duley L, Crowther CA, et al. Antioxidants for preventing pre-eclampsia. The Cochrane database of systematic reviews. 2008(1):Cd004227. 10.1002/14651858.CD004227.pub3
48. Zhou SJ, Yelland L, McPhee AJ, et al. Fish-oil supplementation in pregnancy does not reduce the risk of gestational diabetes or preeclampsia. *Am J Clin Nutr.* 2012;95(6):1378-84. 10.3945/ajcn.111.033217
49. Meher S, Duley L. Garlic for preventing pre-eclampsia and its complications. The Cochrane database of systematic reviews. 2006(3):Cd006065. 10.1002/14651858.Cd006065
50. Bodnar LM, Catov JM, Simhan HN, et al. Maternal vitamin D deficiency increases the risk of preeclampsia. *The Journal of clinical endocrinology and metabolism.* 2007;92(9):3517-22. 10.1210/jc.2007-0718
51. Wen SW, White RR, Rybak N, et al. Effect of high dose folic acid supplementation in pregnancy on pre-eclampsia (FACT): double blind, phase III, randomised controlled, international, multicentre trial. *BMJ (Clinical research ed).* 2018;362:k3478. 10.1136/bmj.k3478
52. Duley L, Henderson-Smart D. Reduced salt intake compared to normal dietary salt, or high intake, in pregnancy. The Cochrane database of systematic reviews. 2000(2):Cd001687. 10.1002/14651858.Cd001687
53. Baschat AA, Magder LS, Doyle LE, et al. Prediction of preeclampsia utilizing the first trimester screening examination. *Am J Obstet Gynecol.* 2014;211(5):514 e1-7. 10.1016/j.ajog.2014.04.018
54. Vane JR, Botting RM. The mechanism of action of aspirin. *Thromb Res.* 2003;110(5-6):255-8. 10.1016/s0049-3848(03)00379-7
55. Wertaschnigg D, Reddy M, Mol BWJ, et al. Evidence-Based Prevention of Preeclampsia: Commonly Asked Questions in Clinical Practice. *J Pregnancy.* 2019;2019:2675101. 10.1155/2019/2675101
56. Ayala DE, Ucieda R, Hermida RC. Chronotherapy with low-dose aspirin for prevention of complications in pregnancy. *Chronobiol Int.* 2013;30(1-2):260-79. 10.3109/07420528.2012.717455
57. Baschat AA, Dewberry D, Seravalli V, et al. Maternal blood-pressure trends throughout pregnancy and development of pre-eclampsia in women receiving first-trimester aspirin prophylaxis. *Ultrasound in Obstetrics & Gynecology.* 2018;52(6):728-33. PMID: 29266502. <https://dx.doi.org/10.1002/uog.18992>
58. Rolnik DL, Wright D, Poon LC, et al. Aspirin versus Placebo in Pregnancies at High Risk for Preterm Preeclampsia. *N Engl J Med.* 2017;377(7):613-22. 10.1056/NEJMoa1704559
59. Duley L, Henderson-Smart DJ, Meher S, et al. Antiplatelet agents for preventing pre-eclampsia and its complications. The Cochrane database of systematic reviews. 2007(2):CD004659.
60. Askie LM, Duley L, Henderson-Smart DJ, et al. Antiplatelet agents for prevention of pre-eclampsia: a meta-analysis of individual patient data. *Lancet.* 2007;369(9575):1791-8. 10.1016/S0140-6736(07)60712-0

61. Voelker R. USPSTF: Low-dose aspirin may help reduce risk of preeclampsia. *JAMA - Journal of the American Medical Association*. 2014;311(20):2055. 10.1001/jama.2014.5390
62. Hoffman MK, Goudar SS, Kodkany BS, et al. Low-dose aspirin for the prevention of preterm delivery in nulliparous women with a singleton pregnancy (ASPIRIN): a randomised, double-blind, placebo-controlled trial. *The Lancet*. 2020;395(10220):285-93. 10.1016/S0140-6736(19)32973-3
63. Tolcher MC, Chu DM, Hollier LM, et al. Impact of USPSTF recommendations for aspirin for prevention of recurrent preeclampsia. *Am J Obstet Gynecol*. 2017;217(3):365.e1-.e8. 10.1016/j.ajog.2017.04.035
64. World Health O. WHO Recommendations for prevention and treatment of pre-eclampsia and eclampsia. World Health Organization; 2011.
65. American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 743: Low-Dose Aspirin Use During Pregnancy. *Obstetrics and gynecology*. 2018;132(1):e44-e52. 10.1097/aog.0000000000002708
66. National Institute for Health and Clinical Excellence. Hypertension in Pregnancy. Manchester, UK: National Institute for Health and Clinical Excellence; 2010.
67. Bushnell C, McCullough LD, Awad IA, et al. Guidelines for the prevention of stroke in women: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45(5):1545-88. 10.1161/01.str.0000442009.06663.48
68. LeFevre ML, Force USPST. Low-dose aspirin use for the prevention of morbidity and mortality from preeclampsia: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2014;161(11):819-26. PMID: 25200125. <https://dx.doi.org/10.7326/M14-1884>
69. Henderson JT, Whitlock EP, O'Connor E, et al. Low-dose aspirin for prevention of morbidity and mortality from preeclampsia: a systematic evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2014;160(10):695-703. 10.7326/M13-2844
70. Henderson JT, Whitlock EP, O'Connor E, et al. Low-dose aspirin for prevention of morbidity and mortality from preeclampsia: a systematic evidence review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2014;160(10):695-703. PMID: 24711050. <https://dx.doi.org/10.7326/M13-2844>
71. U.S. Preventive Services Task Force. Final Recommendation Statement: Low-Dose Aspirin Use for the Prevention of Morbidity and Mortality From Preeclampsia: Preventive Medication. <https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/low-dose-aspirin-use-for-the-prevention-of-morbidity-and-mortality-from-preeclampsia-preventive-medication>. Accessed.
72. United Nations Development Programme. Human Development Report. New York, NY: United Nations; 2016.
73. U.S. Preventive Services Task Force. U.S. Preventive Services Task Force Procedure Manual. Rockville, MD: U.S. Preventive Services Task Force; 2015.
74. Raudenbush SW. Analyzing effect sizes: Random-effects models. In: Cooper H, Hedges LV, Valentine JC, editors. *The Handbook of Research Synthesis and Meta-Analysis*. 2nd ed. New York, New York: Russell Sage Foundation; 2009. p. 296-314.

75. Knapp G, Hartung J. Improved tests for a random effects meta-regression with a single covariate. *Stat Med.* 2003;22(17):2693-710. PMID: 12939780. 10.1002/sim.1482
76. Veroniki AA, Jackson D, Bender R, et al. Methods to calculate uncertainty in the estimated overall effect size from a random-effects meta-analysis. *Res Synth Methods.* 2019;10(1):23-43. 10.1002/jrsm.1319
77. Yusuf S, Peto R, Lewis J, et al. Beta blockade during and after myocardial infarction: An overview of the randomized trials. *Progress in Cardiovascular Diseases.* 1985;27(5):335-71.
78. Cheng J, Pullenayegum E, Marshall JK, et al. Impact of including or excluding both-armed zero-event studies on using standard meta-analysis methods for rare event outcome: a simulation study. *BMJ open.* 2016;6(8):1-10. 10.1136/bmjopen-2015-010983
79. Peters JL, Sutton AJ, Jones DR, et al. Comparison of two methods to detect publication bias in meta-analysis. *Jama.* 2006;295(6):676-80. PMID: 16467236. 10.1001/jama.295.6.676
80. Sterne JAC, Harbord RM. Funnel plots in meta-analysis. *The Stata Journal.* 2004;4(2):127-41. <https://doi.org/10.1177/1536867X0400400204>
81. Jin Z-C, Zhou X-H, He J. Statistical methods for dealing with publication bias in meta-analysis. *Stat Med.* 2015;34(2):343-60. 10.1002/sim.6342
82. Whitlock EP, Eder M, Thompson JH, et al. An approach to addressing subpopulation considerations in systematic reviews: the experience of reviewers supporting the U.S. Preventive Services Task Force. *Systematic Reviews.* 2017;6(1):41. 10.1186/s13643-017-0437-3
83. Higgins JPTT, J.; Cumpston, M.; Li, T.; Page, M. J.; Welch, V. A. (editors). *Cochrane Handbook for Systematic Reviews of Interventions version 6.0.* www.training.cochrane.org/handbook. Accessed.
84. Berkman ND, Lohr KN, Ansari M, et al. Grading the Strength of a Body of Evidence When Assessing Health Care Interventions for the Effective Health Care Program of the Agency for Healthcare Research and Quality: An Update. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews.* AHRQ Publication No. 10(14)-EHC063-EF. Rockville (MD): Agency for Healthcare Research and Quality; 2014. p. 314-49. PMID: None.
85. Atkins D, Eccles M, Flottorp S, et al. Systems for grading the quality of evidence and the strength of recommendations I: critical appraisal of existing approaches The GRADE Working Group. *BMC Health Serv Res.* 2004;4(1):38. PMID: 15615589. <https://doi.org/10.1186/1472-6963-4-38>
86. Wallenburg HC, Dekker GA, Makovitz JW, et al. Low-dose aspirin prevents pregnancy-induced hypertension and pre-eclampsia in angiotensin-sensitive primigravidae. *Lancet.* 1986;1(8471):1-3.
87. Benigni A, Gregorini G, Frusca T, et al. Effect of low-dose aspirin on fetal and maternal generation of thromboxane by platelets in women at risk for pregnancy-induced hypertension. *New England Journal of Medicine.* 1989;321(6):357-62.
88. Schiff E, Peleg E, Goldenberg M, et al. The use of aspirin to prevent pregnancy-induced hypertension and lower the ratio of thromboxane A2 to prostacyclin in relatively high risk pregnancies. *New England Journal of Medicine.* 1989;321(6):351-6.
89. McParland P, Pearce JM, Chamberlain GV. Doppler ultrasound and aspirin in recognition and prevention of pregnancy-induced hypertension. *Lancet.* 1990;335(8705):1552-5.

90. Hauth JC, Goldenberg RL, Parker CR, Jr., et al. Low-dose aspirin therapy to prevent preeclampsia. *Am J Obstet Gynecol.* 1993;168(4):1083-91.
91. Sibai BM, Caritis SN, Thom E, et al. Prevention of preeclampsia with low-dose aspirin in healthy, nulliparous pregnant women. The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *New England Journal of Medicine.* 1993;329(17):1213-8.
92. Viinikka L, Hartikainen-Sorri AL, Lumme R, et al. Low dose aspirin in hypertensive pregnant women: effect on pregnancy outcome and prostacyclin-thromboxane balance in mother and newborn. *Br J Obstet Gynaecol.* 1993;100(9):809-15.
93. Caspi E, Raziell A, Sherman D, et al. Prevention of pregnancy-induced hypertension in twins by early administration of low-dose aspirin: a preliminary report. *American journal of reproductive immunology (New York, NY : 1989).* 1994;31(1):19-24.
94. CLASP: A randomised trial of low-dose aspirin for the prevention and treatment of pre-eclampsia. *Lancet.* 1994;343(8898):619.
95. Davies NJ, Gazvani MR, Farquharson RG, et al. Low-Dose Aspirin in the Prevention of Hypertensive Disorders of Pregnancy in Relatively Low-Risk Nulliparous Women. *Hypertens.* 1995;14(1):49-55.
96. Gallery EDM, Ross MR, Hawkins M, et al. Low-Dose Aspirin in High-Risk Pregnancy? *Hypertens.* 1997;16(2):229-38. 10.3109/10641959709031640
97. Hermida RC, Ayala DE, Iglesias M, et al. Time-dependent effects of low-dose aspirin administration on blood pressure in pregnant women. *Hypertension (Dallas, Tex : 1979).* 1997;30(3 Pt 2):589-95.
98. Caritis S, Sibai B, Hauth J, et al. Low-Dose Aspirin to Prevent Preeclampsia in Women at High Risk. *New England Journal of Medicine.* 1998;338(11):701-5. 10.1056/NEJM199803123381101
99. Rotchell YE, Cruickshank JK, Gay MP, et al. Barbados Low Dose Aspirin Study in Pregnancy (BLASP): a randomised trial for the prevention of pre-eclampsia and its complications. *Br J Obstet Gynaecol.* 1998;105(3):286-92.
100. Grab D, Paulus WE, Erdmann M, et al. Effects of low-dose aspirin on uterine and fetal blood flow during pregnancy: results of a randomized, placebo-controlled, double-blind trial. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology.* 2000;15(1):19-27.
101. Subtil D, Goeusse P, Puech F, et al. Aspirin (100 mg) used for prevention of pre-eclampsia in nulliparous women: the Essai Regional Aspirine Mere-Enfant study (Part 1). *BJOG : an international journal of obstetrics and gynaecology.* 2003;110(5):475-84.
102. Yu CK, Papageorghiou AT, Parra M, et al. Randomized controlled trial using low-dose aspirin in the prevention of pre-eclampsia in women with abnormal uterine artery Doppler at 23 weeks' gestation. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology.* 2003;22(3):233-9.
103. Villa P, Kajantie E, Raikonen K, et al. Aspirin in the prevention of pre-eclampsia in high-risk women: a randomised placebo-controlled PREDO Trial and a meta-analysis of randomised trials. *BJOG : an international journal of obstetrics and gynaecology.* 2013;120(1):64-74.
104. Mone F, Mulcahy C, McParland P, et al. Trial of feasibility and acceptability of routine low-dose aspirin versus Early Screening Test indicated aspirin for pre-eclampsia

- prevention (TEST study): a multicentre randomised controlled trial. *BMJ open*. 2018;8(7):e022056. PMID: 30056389. 10.1136/bmjopen-2018-022056
105. Scazzocchio E, Oros D, Diaz D, et al. Impact of aspirin on trophoblastic invasion in women with abnormal uterine artery Doppler at 11-14 weeks: a randomized controlled study. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2017;49(4):435-41. 10.1002/uog.17351
 106. Morris JM, Fay RA, Ellwood DA, et al. A randomized controlled trial of aspirin in patients with abnormal uterine artery blood flow. *Obstetrics and gynecology*. 1996;87(1):74-8. 10.1016/0029-7844(95)00340-1
 107. Jensen MS, Rebordosa C, Thulstrup AM, et al. Maternal use of acetaminophen, ibuprofen, and acetylsalicylic acid during pregnancy and risk of cryptorchidism. *Epidemiology*. 2010;21(6):779-85.
 108. Keim SA, Klebanoff MA. Aspirin use and miscarriage risk. *Epidemiology*. 2006;17(4):435-9.
 109. Newnham JP, Godfrey M, Walters BJ, et al. Low dose aspirin for the treatment of fetal growth restriction: a randomized controlled trial. *The Australian & New Zealand journal of obstetrics & gynaecology*. 1995;35(4):370-4.
 110. Vainio M, Kujansuu E, Iso-Mustajarvi M, et al. Low dose acetylsalicylic acid in prevention of pregnancy-induced hypertension and intrauterine growth retardation in women with bilateral uterine artery notches. *BJOG : an international journal of obstetrics and gynaecology*. 2002;109(2):161-7.
 111. Poon LC, Wright D, Rolnik DL, et al. Aspirin for Evidence-Based Preeclampsia Prevention trial: effect of aspirin in prevention of preterm preeclampsia in subgroups of women according to their characteristics and medical and obstetrical history. *Am J Obstet Gynecol*. 2017;217(5):585.e1-.e5. 10.1016/j.ajog.2017.07.038
 112. Clasp Collaborative Group. Low dose aspirin in pregnancy and early childhood development: follow up of the collaborative low dose aspirin study in pregnancy. CLASP collaborative group 50. *Br J Obstet Gynaecol*. 1995;102(11):861-8.
 113. Adkins K, Allshouse AA, Metz TD, et al. Impact of aspirin on fetal growth in diabetic pregnancies according to White classification. *American Journal of Obstetrics & Gynecology*. 2017;217(4):465.e1-.e5. PMID: 28599894. <https://dx.doi.org/10.1016/j.ajog.2017.05.062>
 114. Abramovici A, Jauk V, Wetta L, et al. Low-dose aspirin, smoking status, and the risk of spontaneous preterm birth. *American journal of perinatology*. 2015;32(5):445-50. PMID: 25261702. <https://dx.doi.org/10.1055/s-0034-1390352>
 115. Cantu JA, Jauk VR, Owen J, et al. Is low-dose aspirin therapy to prevent preeclampsia more efficacious in non-obese women or when initiated early in pregnancy? *Journal of Maternal-Fetal & Neonatal Medicine*. 2015;28(10):1128-32. PMID: 25048750. <https://dx.doi.org/10.3109/14767058.2014.947258>
 116. Tolcher MC, Sangi-Haghpeykar H, Aagaard KM. Low-dose aspirin (ASA) for preeclampsia prevention among high-risk women by ethnicity and race. *American Journal of Obstetrics and Gynecology*. 2018;218(1):S564.
 117. American College of Obstetricians and Gynecologists. Practice Bulletin No. 183: Postpartum Hemorrhage. *Obstetrics & Gynecology*. 2017;130(4):e168-e86. PMID: 00006250-201710000-00056. 10.1097/aog.0000000000002351

118. Centers for Disease Control and Prevention. Data on Selected Pregnancy Complications in the United States.
<https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-complications-data.htm#post>. Accessed.
119. Creanga AA, Berg CJ, Ko JY, et al. Maternal mortality and morbidity in the United States: where are we now? *J Womens Health (Larchmt)*. 2014;23(1):3-9. PMID: 24383493. 10.1089/jwh.2013.4617
120. Callaghan WM, Mackay AP, Berg CJ. Identification of severe maternal morbidity during delivery hospitalizations, United States, 1991-2003. *Am J Obstet Gynecol*. 2008;199(2):133 e1-8. PMID: 18279820. 10.1016/j.ajog.2007.12.020
121. Koopmans CM, van der Tuuk K, Groen H, et al. Prediction of postpartum hemorrhage in women with gestational hypertension or mild preeclampsia at term. *Acta obstetrica et gynecologica Scandinavica*. 2014;93(4):399-407. PMID: 24575790. 10.1111/aogs.12352
122. Duley L, Meher S, Hunter KE, et al. Antiplatelet agents for preventing pre-eclampsia and its complications. *Cochrane Database of Systematic Reviews*. 2019(10). 10.1002/14651858.CD004659.pub3
123. Landon M, Galan H, Jauniaux E, et al. Antepartum and Postpartum Hemorrhage. *Gabbe's Obstetrics: Normal and Problem Pregnancies*. 8 ed. Philadelphia, PA: Elsevier; 2021.
124. Landon M, Galan H, Jauniaux E, et al. The Neonate. *Gabbe's Obstetrics: Normal and Problem Pregnancies*. 8 ed. Philadelphia, PA: Elsevier; 2021.
125. Stewart GB, Altman DG, Askie LM, et al. Statistical analysis of individual participant data meta-analyses: a comparison of methods and recommendations for practice. *PLoS ONE [Electronic Resource]*. 2012;7(10):e46042.
126. Meher S, Duley L, Hunter K, et al. Antiplatelet therapy before or after 16 weeks' gestation for preventing preeclampsia: an individual participant data meta-analysis. *Am J Obstet Gynecol*. 2017;216(2):121-8.e2. 10.1016/j.ajog.2016.10.016
127. van Vliet EO, Askie LA, Mol BW, et al. Antiplatelet Agents and the Prevention of Spontaneous Preterm Birth: A Systematic Review and Meta-analysis. *Obstetrics and gynecology*. 2017;129(2):327-36. 10.1097/aog.0000000000001848
128. Seidler AL, Askie L, Ray JG. Optimal aspirin dosing for preeclampsia prevention. *American Journal of Obstetrics & Gynecology*. 2018;219(1):117-8. PMID: 29588190.
<https://dx.doi.org/10.1016/j.ajog.2018.03.018>
129. Nakhai-Pour HR, Berard A. Major malformations after first-trimester exposure to aspirin and NSAIDs. *Expert Review of Clinical Pharmacology*. 2008;1(5):605-16.
130. Coomarasamy A, Braunholtz D, Song F, et al. Individualising use of aspirin to prevent pre-eclampsia: a framework for clinical decision making. *BJOG : an international journal of obstetrics and gynaecology*. 2003;110(10):882-8.
131. Hernandez RK, Werler MM, Romitti P, et al. Nonsteroidal antiinflammatory drug use among women and the risk of birth defects. *American Journal of Obstetrics and Gynecology*. 2012;206(3):228.e1-.e8. 10.1016/j.ajog.2011.11.019
132. Given JE, Loane M, Garne E, et al. Gastroschisis in Europe – A Case-malformed-Control Study of Medication and Maternal Illness during Pregnancy as Risk Factors. *Paediatr Perinat Epidemiol*. 2017;31(6):549-59. 10.1111/ppe.12401
133. Schulz K, Grimes D. *Essential Concepts in Clinical Research: Randomised Controlled Trials and Observational Epidemiology*. 2 ed: Elsevier; 2019.

134. Wright D, Poon LC, Rolnik DL, et al. Aspirin for Evidence-Based Preeclampsia Prevention trial: influence of compliance on beneficial effect of aspirin in prevention of preterm preeclampsia. *American Journal of Obstetrics & Gynecology*. 2017;217(6):685.e1-.e5. PMID: 28888591. <https://dx.doi.org/10.1016/j.ajog.2017.08.110>
135. Shanmugalingam R, Wang X, Motum P, et al. Clinical Influence of Nonadherence With Prophylactic Aspirin in Preventing Preeclampsia in High-Risk Pregnancies: A Multicenter, Prospective, Observational Cohort Study. *Hypertension (Dallas, Tex : 1979)*. 2020;75(4):1125-32. PMID: 32114852. 10.1161/HYPERTENSIONAHA.119.14107
136. Young BC, Thadhani R, Karumanchi SA. Aspirin to Prevent Preterm Preeclampsia: More Progress. *Hypertension (Dallas, Tex : 1979)*. 2020;75(4):941-2. PMID: 32114845. 10.1161/HYPERTENSIONAHA.119.14277
137. van Montfort P, Scheepers HCJ, van Dooren IMA, et al. Low-dose-aspirin usage among women with an increased preeclampsia risk: A prospective cohort study. *Acta obstetrica et gynecologica Scandinavica*. 2020. PMID: 31953956. 10.1111/aogs.13808
138. Al-Rubaie Z, Askie LM, Ray JG, et al. The performance of risk prediction models for pre-eclampsia using routinely collected maternal characteristics and comparison with models that include specialised tests and with clinical guideline decision rules: a systematic review. *BJOG : an international journal of obstetrics and gynaecology*. 2016;123(9):1441-52. PMID: 10.1111/1471-0528.14029 [doi].
139. Tan MY, Wright D, Syngelaki A, et al. Comparison of diagnostic accuracy of early screening for pre-eclampsia by NICE guidelines and a method combining maternal factors and biomarkers: results of SPREE. *Ultrasound in Obstetrics & Gynecology*. 2018;51(6):743-50. 10.1002/uog.19039
140. Akolekar R, Syngelaki A, Poon L, et al. Competing risks model in early screening for preeclampsia by biophysical and biochemical markers. *Fetal Diagn Ther*. 2013;33(1):8-15. PMID: 22906914.
141. O'Gorman N, Wright D, Poon LC, et al. Multicenter screening for pre-eclampsia by maternal factors and biomarkers at 11-13 weeks' gestation: comparison with NICE guidelines and ACOG recommendations. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2017;49(6):756-60. PMID: 28295782. 10.1002/uog.17455
142. Roberts JM, Himes KP. Pre-eclampsia: Screening and aspirin therapy for prevention of pre-eclampsia. *Nature Reviews Nephrology*. 2017;13(10):602-4. 10.1038/nrneph.2017.121
143. Allen RE, Zamora J, Arroyo-Manzano D, et al. External validation of preexisting first trimester preeclampsia prediction models. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2017;217:119-25. 10.1016/j.ejogrb.2017.08.031
144. Bailey B, Euser AG, Bol KA, et al. High-altitude residence alters blood-pressure course and increases hypertensive disorders of pregnancy. *J Matern Fetal Neonatal Med*. 2020:1-8. PMID: 32228111. 10.1080/14767058.2020.1745181
145. Bai W, Li Y, Niu Y, et al. Association between ambient air pollution and pregnancy complications: A systematic review and meta-analysis of cohort studies. *Environ Res*. 2020;185:109471. PMID: 32276169. 10.1016/j.envres.2020.109471
146. Wilson DL, Howard ME, Fung AM, et al. The presence of coexisting sleep-disordered breathing among women with hypertensive disorders of pregnancy does not worsen

- perinatal outcome. *PloS one*. 2020;15(2):e0229568. PMID: 32101584. 10.1371/journal.pone.0229568
147. Li L, Zhao K, Hua J, et al. Association between Sleep-Disordered Breathing during Pregnancy and Maternal and Fetal Outcomes: An Updated Systematic Review and Meta-Analysis. *Front Neurol*. 2018;9:91. PMID: 29892255. 10.3389/fneur.2018.00091
 148. Aukes AM, Yurtsever FN, Boutin A, et al. Associations Between Migraine and Adverse Pregnancy Outcomes: Systematic Review and Meta-analysis. *Obstet Gynecol Surv*. 2019;74(12):738-48. PMID: 31881091. 10.1097/OGX.0000000000000738
 149. Palatnik A, Grobman WA, Miller ES. Is a history of preeclampsia associated with an increased risk of a small for gestational age infant in a future pregnancy? *American Journal of Obstetrics & Gynecology*. 2016;215(3):355.e1-6. PMID: 26994655. <https://dx.doi.org/10.1016/j.ajog.2016.03.011>
 150. North RA, McCowan LM, Dekker GA, et al. Clinical risk prediction for pre-eclampsia in nulliparous women: development of model in international prospective cohort. *BMJ (Clinical research ed)*. 2011;342:d1875.
 151. Myatt L, Clifton RG, Roberts JM, et al. First-trimester prediction of preeclampsia in nulliparous women at low risk. *Obstetrics and gynecology*. 2012;119(6):1234-42. PMID: 22617589.
 152. Chappell LC, Seed PT, Myers J, et al. Exploration and confirmation of factors associated with uncomplicated pregnancy in nulliparous women: prospective cohort study. *BMJ (Clinical research ed)*. 2013;347:f6398. PMID: 24270055.
 153. Skrastad R, Hov G, Blaas HG, et al. Risk assessment for preeclampsia in nulliparous women at 11-13 weeks gestational age: prospective evaluation of two algorithms. *BJOG : an international journal of obstetrics and gynaecology*. 2014. PMID: 25471057.
 154. Mone F, Mulcahy C, McParland P, et al. Should we recommend universal aspirin for all pregnant women? *American Journal of Obstetrics & Gynecology*. 2017;216(2):141.e1-.e5. PMID: 27659212. <https://dx.doi.org/10.1016/j.ajog.2016.09.086>
 155. Ayala NK, Rouse DJ. A Nudge Toward Universal Aspirin for Preeclampsia Prevention. *Obstetrics and gynecology*. 2019;133(4):725-8. PMID: 30870274. 10.1097/AOG.0000000000003167
 156. Mone F, O'Mahony JF, Tyrrell E, et al. Preeclampsia Prevention Using Routine Versus Screening Test-Indicated Aspirin in Low-Risk Women. *Hypertension (Dallas, Tex : 1979)*. 2018;72(6):1391-6. PMID: 30571234. <https://dx.doi.org/10.1161/HYPERTENSIONAHA.118.11718>
 157. Rolnik DL, O'Gorman N, Roberge S, et al. Early screening and prevention of preterm pre-eclampsia with aspirin: time for clinical implementation. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2017;50(5):551-6. PMID: 28887883. 10.1002/uog.18899
 158. Ross KM, Dunkel Schetter C, McLemore MR, et al. Socioeconomic Status, Preeclampsia Risk and Gestational Length in Black and White Women. *J Racial Ethn Health Disparities*. 2019;6(6):1182-91. PMID: 31368002. 10.1007/s40615-019-00619-3
 159. Shahul S, Tung A, Minhaj M, et al. Racial Disparities in Comorbidities, Complications, and Maternal and Fetal Outcomes in Women With Preeclampsia/eclampsia. *Hypertens Pregnancy*. 2015;34(4):506-15. PMID: 26636247. 10.3109/10641955.2015.1090581

160. Wu P, Chew-Graham CA, Maas AH, et al. Temporal Changes in Hypertensive Disorders of Pregnancy and Impact on Cardiovascular and Obstetric Outcomes. *Am J Cardiol*. 2020. PMID: 32273052. 10.1016/j.amjcard.2020.02.029
161. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension (Dallas, Tex : 1979)*. 2018;71(6):e13-e115. PMID: 29133356. 10.1161/hyp.0000000000000065
162. Duckitt K, Harrington D. Risk factors for pre-eclampsia at antenatal booking: systematic review of controlled studies. *BMJ (Clinical research ed)*. 2005;330(7491):565.
163. Hauspurg A, Sutton EF, Catov JM, et al. Aspirin Effect on Adverse Pregnancy Outcomes Associated With Stage 1 Hypertension in a High-Risk Cohort. *Hypertension (Dallas, Tex : 1979)*. 2018;72(1):202-7. PMID: 29802215. 10.1161/hypertensionaha.118.11196
164. Finneran MM, Gonzalez-Brown VM, Smith DD, et al. Obesity and laboratory aspirin resistance in high-risk pregnant women treated with low-dose aspirin. *American Journal of Obstetrics & Gynecology*. 2019;220(4):385.e1-.e6. PMID: 30786253.
<https://dx.doi.org/10.1016/j.ajog.2019.01.222>
165. Menard MK, Main EK, Currigan SM. Executive summary of the reVITALize initiative: standardizing obstetric data definitions. *Obstetrics and gynecology*. 2014;124(1):150-3. PMID: 24901267. 10.1097/AOG.0000000000000322
166. American College of Obstetricians and Gynecologists. Obstetric Data Definitions: reVITALize. <https://www.acog.org/-/media/project/acog/acogorg/files/pdfs/publications/revitalize-ob.pdf?la=en&hash=D9ECE1577C2A0B535F42C1B8DC492070>. Accessed.
167. Jayet PY, Rimoldi SF, Stuber T, et al. Pulmonary and systemic vascular dysfunction in young offspring of mothers with preeclampsia. *Circulation*. 2010;122(5):488-94. PMID: 20644018. 10.1161/CIRCULATIONAHA.110.941203
168. Nahum Sacks K, Friger M, Shoham-Vardi I, et al. Prenatal exposure to preeclampsia as an independent risk factor for long-term cardiovascular morbidity of the offspring. *Pregnancy Hypertens*. 2018;13:181-6. PMID: 30177050. 10.1016/j.preghy.2018.06.013
169. Maher GM, Dalman C, O'Keeffe GW, et al. Association between preeclampsia and attention-deficit hyperactivity disorder: a population-based and sibling-matched cohort study. *Acta Psychiatr Scand*. 2020. PMID: 32056200. 10.1111/acps.13162
170. Nahum Sacks K, Friger M, Shoham-Vardi I, et al. Long-term neuropsychiatric morbidity in children exposed prenatally to preeclampsia. *Early Hum Dev*. 2019;130:96-100. PMID: 30711915. 10.1016/j.earlhumdev.2019.01.016
171. Sun BZ, Moster D, Harmon QE, et al. Association of Preeclampsia in Term Births With Neurodevelopmental Disorders in Offspring. *JAMA Psychiatry*. 2020. 10.1001/jamapsychiatry.2020.0306
172. Davis EF, Lazdam M, Lewandowski AJ, et al. Cardiovascular risk factors in children and young adults born to preeclamptic pregnancies: a systematic review. *Pediatrics*. 2012;129(6):e1552-e61.
173. Valdiviezo C, Garovic VD, Ouyang P. Preeclampsia and hypertensive disease in pregnancy: their contributions to cardiovascular risk. *Clin Cardiol*. 2012;35(3):160-5.

174. Bellamy L, Casas JP, Hingorani AD, et al. Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and meta-analysis. *BMJ (Clinical research ed)*. 2007;335(7627):974.
175. Lykke JA, Langhoff-Roos J, Sibai BM, et al. Hypertensive pregnancy disorders and subsequent cardiovascular morbidity and type 2 diabetes mellitus in the mother. *Hypertension (Dallas, Tex : 1979)*. 2009;53(6):944-51.
176. Miller EC, Boehme AK, Moon YP, et al. Preeclampsia and early stroke incidence in The California teachers study. *Stroke*. 2018;49.
177. Mongraw-Chaffin ML, Cirillo PM, Cohn BA. Preeclampsia and cardiovascular disease death: prospective evidence from the child health and development studies cohort. *Hypertension (Dallas, Tex : 1979)*. 2010;56(1):166-71.
178. Bergman L, Torres-Vergara P, Penny J, et al. Investigating Maternal Brain Alterations in Preeclampsia: the Need for a Multidisciplinary Effort. *Curr Hypertens Rep*. 2019;21(9):72. PMID: 31375930. 10.1007/s11906-019-0977-0
179. Aukes AM, De Groot JC, Wiegman MJ, et al. Long-term cerebral imaging after pre-eclampsia. *BJOG : an international journal of obstetrics and gynaecology*. 2012;119(9):1117-22.
180. Wiegman MJ, De Groot JC, Jansonius NM, et al. Long-term visual functioning after eclampsia. *Obstetrics and gynecology*. 2012;119(5):959-66.
181. Lederer M, Wong A, Diego D, et al. Tracking the Development of Cerebrovascular Risk Factors Following Pregnancy With Preeclampsia. *J Stroke Cerebrovasc Dis*. 2020:104720. PMID: 32220554. 10.1016/j.jstrokecerebrovasdis.2020.104720
182. National Institute for Health and Clinical Excellence. Hypertension in pregnancy: diagnosis and management Manchester, UK: National Institute for Health and Clinical Excellence; 2020.
183. FIGO Working Group on Good Clinical Practice in Maternal-Fetal Medicine. Good clinical practice advice: First trimester screening and prevention of pre-eclampsia in singleton pregnancy. *Int J Gynaecol Obstet*. 2019;144(3):325-9. 10.1002/ijgo.12741

Figure 1. Analytic Framework





Figure 2. Perinatal Mortality, Sorted by Study Size




Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio

*Excluded 4 studies with both arms having zero events.


Figure 3. Preterm Birth Before 37 Weeks' Gestation, Sorted by Study Size



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio


Figure 4. Small for Gestational Age or Intrauterine Growth Restriction, Sorted by Study Size



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio


Figure 5. Small for Gestational Age and Intrauterine Growth Restriction Stratified by Timing of Aspirin Initiation



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio


Figure 6. Preterm Birth Stratified by Timing of Aspirin Initiation



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio


Figure 7. Preeclampsia, Sorted by Study Size



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio


Figure 8. Preeclampsia Stratified by Timing of Aspirin Initiation



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio

Figure 9. Placental Abruption Stratified by Risk, Sorted by Study Size




Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio

*Excluded 3 studies with both arms having zero events.

Figure 10. Postpartum Hemorrhage Stratified by Risk, Sorted by Study Size




Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio

*Excluded 2 studies with both arms having zero events.

Figure 11. Fetal Intracranial Bleeding Stratified by Risk, Sorted by Study Size



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PE = Preeclampsia; REML = Restricted maximum likelihood; RR = Risk ratio

*Excluded 3 studies with both arms having zero events.

Table 1. Existing Clinical Recommendations

Organization	Guideline	Definition of treatment population
<p><i>National Institute for Health and Clinical Excellence (NICE)</i>¹⁸²</p>	<p>Advise pregnant women at high risk or with more than 1 moderate risk factor for pre-eclampsia to take 75–150 mg of aspirin daily from 12 weeks until the birth of the baby.</p>	<p>High risk factors (any one): Hypertensive disease during a previous pregnancy Chronic kidney disease Autoimmune disease such as systemic lupus erythematosus or antiphospholipid syndrome Type 1 or type 2 diabetes Chronic hypertension</p> <p>Moderate risk factors (>1 required): First pregnancy Age 40 years or older Pregnancy interval of more than 10 years BMI of 35 kg/m² or more at first visit Family history of preeclampsia (mother, sister) Multi-fetal pregnancy</p>
<p><i>International Federation of Gynecology and Obstetrics (FIGO)</i>¹⁸³</p>	<p>Daily administration of aspirin starting at ≤ 16weeks and at a dose of ≥ 100 mg/d at night continuing until 36 weeks' gestation or presence of labor signs</p>	<p>Women with singleton pregnancy at gestational age 11-13⁶⁺ weeks based on risk assessment using a prediction algorithm using a combination of maternal factors with uterine artery pulsatility index, mean arterial pressure, and serum placental growth factor at 11-13 weeks' gestation</p>
<p><i>American College of Obstetricians and Gynecologists (ACOG)</i>⁶⁵</p>	<p>Clinical risk assessment using patient medical history to identify patients for whom to recommend initiation of 81 mg/day of aspirin between 12 and 28 weeks of gestation (optimally before 16 weeks), continued until delivery Recommend low-dose aspirin if the patient has one or more high-risk factors. Consider low-dose aspirin if the patient has more than one moderate risk factor. Do not recommend low-dose aspirin for low-risk individuals (e.g., previous uncomplicated full-term delivery)</p>	<p>High risk factors: History of preeclampsia, especially when accompanied by an adverse outcome Multifetal gestation Chronic hypertension Type 1 or 2 diabetes Renal disease Autoimmune disease (systemic lupus erythematosus, antiphospholipid syndrome)</p> <p>Moderate risk factors (>1): Nulliparity Obesity (body mass index > 30) Family history of preeclampsia (mother or sister) Sociodemographic characteristics (African American race, low socioeconomic status) Age 35 years or older Personal history factors (e.g., low birthweight or small for gestational age, previous adverse pregnancy outcome, more than 10-year pregnancy interval)</p>
<p><i>American Heart Association (AHA) and American Stroke Association (ASA)</i>⁶⁷</p>	<p>The AHA/ASA recommends that pregnant women with chronic primary or secondary hypertension or previous pregnancy-related hypertension should take low-dose aspirin from the 12th week of gestation until delivery to prevent preeclampsia.</p>	<p>Women with chronic primary or secondary hypertension or previous pregnancy-related hypertension</p>
<p><i>World Health Organization (WHO), 2011</i>⁶⁴</p>	<p>WHO recommends that women at high risk of preeclampsia take 75mg of aspirin daily, initiated before 20 weeks of pregnancy.</p>	<p>High risk: Previous preeclampsia Diabetes Chronic hypertension Renal disease Autoimmune disease Multiple pregnancy</p>

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
ASPRES, 2017 ⁵⁸	Spain, Italy, UK, Israel, Belgium, Greece	Good	1776	Increased risk, risk prediction algorithm at 11-13 weeks' gestation	Unconscious or severely ill status, learning difficulties or serious mental illness, major fetal abnormality, regular aspirin use within 28 days before screening, bleeding disorder, peptic ulceration, hypersensitivity to aspirin, long-term use of nonsteroidal antiinflammatory medication	150	Not specified 11 to 14 weeks; 36 weeks	Placebo	31.5*	IG: 68.5 CG: 66.1	White: 67.1 AA: 25.2 Asian: 6.4 Other: 1.3	IG: 7.1 CG: 7.2	IG: 6.1 CG: 7.4
Ayala, 2012 ⁵⁶	Spain	Good	350	Increased risk, receiving pregnancy care at high-risk obstetric unit due to range of factors	Multiple pregnancy, chronic HTN or any other condition requiring the use of BP-lowering medication, cardiovascular disorders, chronic liver disease, any disease requiring the use of anti-inflammatory medication, DM or any other endocrine disease such as hyperthyroidism	100	Randomly assigned (various) 12 to 16 weeks; delivery	Placebo	30.7	IG: 49.4 CG: 55.1	NR	NR	NR

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
Benigni, 1989 ⁸⁷	Italy	Fair	33	Increased risk, HTN or previous history of fetal death due to placental insufficiency, severe IUGR, early onset PreE [<32 weeks]	Women with antiphospholipid antibodies (lupus-like, anticoagulant, anticardiolipin antibodies)	60	Not specified 12 weeks; delivery	Placebo	31.5 (NR)	NR	NR	IG: 6.3 CG: 0	IG: 35.2 CG: 31.3
Caspi, 1994 ⁹³	Ireland	Good	48	Increased risk, uncomplicated twin pregnancies	Chronic renal, cardiovascular, pulmonary or hepatic disorders, coagulopathy or peptic ulcer, gestational DM, known hypersensitivity to aspirin	100	Morning Start of the second trimester; delivery	Placebo	28.3	IG: 41.6 CG: 30.4	NR	NR	IG: 1.0 CG: 3.0
CLASP, 1994 ⁹⁴	US, Australia, Canada, Germany, Spain, Hong Kong, Ireland, Netherlands, New Zealand, Sweden, UK,		9364	Increased risk, determined by clinician based on range of factors including obstetric history, family history, patient and pregnancy characteristics	Increased risk of bleeding, asthma, allergy to aspirin, or a high likelihood of immediate delivery	60	Not specified 12 to 32 weeks; delivery	Placebo	28.5	IG: 27.9 CG: 27.9	NR	IG: 20.7 CG: 20.1	IG: 19.9 CG: 20.2

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
Argentina, Belgium, Malaysia, Russia, UAE		Good											
Davies, 1995 ⁹⁵	UK	Fair	122	Increased risk, hemoglobin concentration greater than 13.2 g/dL	Multiple pregnancy, DM, recurrent spontaneous abortions, or any contraindication to aspirin therapy	75	Not specified 18 weeks; delivery	Placebo	25.0	IG: 100 CG: 100	White: 95.8	IG: 10.3 CG: 13.3	IG: 0 CG: 0
Gallery, 1997 ⁹⁶	Australia	Fair	108	Increased risk, chronic HTN or previous early, severe PreE	Women with history of aspirin allergy, aspirin-sensitive asthma, or preexisting bleeding diathesis	100	Not specified 17 to 19 weeks; two weeks before planned delivery	Placebo	28.5 (22-38)	IG: 42.0 CG: 43.0	White: 95.5	NR	IG: 57.0 CG: 52.0
Grab, 2000 ¹⁰⁰	Germany	Fair	43	Increased risk, early IUGR, impaired uteroplacental blood, chronic HTN or previous still birth, growth restriction, or PreE	DM, proteinuric HTN; fetal malformations or chromosome abnormalities	100	Not specified 18 weeks; 38 weeks	Placebo	NR	NR	NR	NR	IG: 50.0 CG: 38.1

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
Hermida, 1997 ⁹⁷ Spain Good			100	Increased risk, receiving pregnancy care at high-risk hospital unit due to broad range of PreE factors	Multiple gestation pregnancy, chronic HTN, chronic liver disease, disease requiring the use of anti-inflammatory medication, DM, hyperthyroidism	100	Randomly assigned (various) 12 to 16 weeks; delivery	Placebo	30.2	NR	White: 100 AA: 0 Asian: 0 NatAm: 0 Hispanic: 0 Other: 0	NR	NR
McParland, 1990 ⁸⁹ UK Fair			106	Increased risk, nulliparous with persistent abnormal Doppler flow-velocity waveforms at 24 weeks' gestation	Known aspirin allergy, maternal DM, bleeding disorders, peptic ulceration, systemic lupus erythematosus	75	Not specified 24 weeks; delivery	Placebo	26.1	IG: 100 CG: 100	White: 69.0 AA: 20.0 Asian: 7.0 Other: 4.0	IG: 16 CG: 23	NR
MFMU, 1998 ⁹⁸ US Good			2539	Increased risk, four risk groups: DM, chronic HTN, multifetal gestation, previous PreE	Multifetal gestation in combination with DM, HTN, or proteinuria, history of PreE, current proteinuria	60	Not specified 13 to 26 weeks; delivery or if PreE developed	Placebo	26.5	NR	White: 32.7 AA: 56.3 Hispanic: 10.8 Other: 0.5	IG: 16.8 CG: 16.4	NR
Morris, 1996 ¹⁰⁶			102	Nulliparous with abnormal uterine artery waveforms at 18 weeks' gestation	Pregnancies dated at less than 17 weeks' or more than 19 weeks' of gestation	100	Not specified 17-19 weeks; NR	Placebo	23.8	100	NR	NR	NR

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
Scazzocchio, 2017 ¹⁰⁵	Spain	Good	186	Increased risk, abnormal Doppler ultrasound at 11-14 weeks' gestation, maternal age > 18 years	Preexisting HTN, immune, renal, or cardiovascular disease, history of PreE in a previous pregnancy, history of gastric ulcer, known allergy or hypersensitivity to aspirin, hemorrhagic disease, fetal malformation, active treatment with heparin or aspirin before recruitment, multifetal gestation	150	Evening 11 to 14 weeks; delivery	Placebo	32.9	IG: 62.5 CG: 64.0	NR	IG: 11.3 CG: 12.0	IG: 0 CG: 0
Schiff, 1989 ⁸⁸	Israel	Good	65	Increased risk, nulliparity, twin gestation, history of PreE, or positive rollover test	Chronic HTN, pregnancy induced hypertension or proteinuria in current pregnancy, long-term treatment with nonsteroidal anti-inflammatory drugs or use in previous 6 weeks, thrombocytopenia, coagulation disorders, heart failure, chronic renal or pulmonary disease, hepatic or peptic ulcer	100	Not specified 28 or 29 weeks to 38 weeks	Placebo	27.4	NR	White: 100	NR	IG: 0 CG: 0

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
					disease, hypersensitivity to aspirin								
Viinikka, 1993 ⁹²	Finland	Fair	208	Increased risk, HTN or previous severe PreE	None	50	Not specified 16 weeks; delivery	Placebo	33.0	IG: 25.2 CG: 23.8	NR	NR	IG: 86.4 CG: 91.4
Villa, 2012 ¹⁰³	Finland	Fair	152	Increased risk, broad range of PreE risk factors and abnormal Doppler ultrasound at 12-14 weeks' gestation	Aspirin allergy, tobacco smoking in current pregnancy, multiple gestation pregnancy, asthma, peptic ulcer, placental ablation, inflammatory bowel diseases, rheumatoid arthritis, hemophilia or thrombophilia	100	Not specified 12 to 14 weeks; 35 weeks or delivery (whatever came first)	Placebo	30.9 (20-40)	IG: 26.2 CG: 15.0	NR	IG: 0 CG: 0	IG: 13.1 CG: 20.0
Wallenburg, 1986 ⁸⁶	Netherlands	Good	46	Increased risk, Angiotensin-II sensitivity determined by blood test	HTN or cardiovascular renal disease	60	Morning 28 weeks; delivery	Placebo	24 (17-38)	IG: 100 CG: 100	NR	IG: 21.7 CG: 17.4	IG: 0 CG: 0
Yu, 2003 ¹⁰²	UK, Chile, South Africa, Brazil	Good	560	Increased risk, abnormal Doppler ultrasound at 22-24 weeks' gestation	Chronic HTN, renal disease, cardiovascular disease, DM, bleeding disorders, systemic lupus erythematosus,	150	Not specified 22 to 24 weeks; 36 weeks	Placebo	29* (23-33)	IG: 26.8 CG: 23.4	White: 62.3 AA: 27.6 Other: 10.1	IG: 9.4 CG: 9.7	IG: 0 CG: 0

Table 2. Population, Study Design, and Intervention Characteristics, Increased-Risk Population

Author, year	Country	Study quality	N rand	Included Populations	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
					peptic ulceration, hypersensitivity to aspirin, fetal abnormality or fetal growth restriction at 23 weeks' gestation								

Abbreviations: AA = African American; ABPM = Ambulatory blood pressure monitoring; AIDS = Acquired immunodeficiency syndrome; CG = Control group; DM = Diabetes mellitus; Hisp = Hispanic; HTN = Hypertension; IG = Intervention group; NR = Not reported; PreE = Preeclampsia; UK = United Kingdom; US = United States

*Median

Table 3. Population, Study Design, and Intervention Characteristics, Average-Risk Population

Author, year	Country	Study quality	N rand	Included Population	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ethnicity, %	Smoking, %	Chronic HTN, %
Hauth, 1993 ⁹⁰	US	Good	606	Average risk, nulliparous	Any illness or condition known to increase the incidence of PreE or PIH (i.e. renal disease, collagen vascular disease, DM, multifetal gestation, chronic HTN)	60	Not specified No later than 22 weeks; delivery	Placebo	20.4	IG: 100 CG: 100	White: 28.5 AA: 71.5	NR	IG: 0 CG: 0
Mone, 2018 ¹⁰⁴	Ireland	Fair	362	Average risk, nulliparous	Fetal congenital anomaly at recruitment, aspirin hypersensitivity	75	Evening 11 to 13 weeks; 36 weeks	No treatment	33.5 (19-44)	IG: 100 CG: 100	White: 96.8 AA: 0.8 Asian: 2.4 Other: 0	IG: 9.2 CG: 5.9	IG: 0 CG: 0
Rotchell, 1998 ⁹⁹	Barbados	Good	3647	Average risk, nulliparous and multiparous	Increased risk of bleeding, known allergy or contraindication to aspirin, high likelihood of immediate delivery, previous placental abruption	75	Not specified 12 to 32 weeks; delivery	Placebo	NR	IG: 44.0 CG: 44.0	NR	NR	IG: 0.44 CG: 0.44
Sibai, 1993 ⁹¹	US	Good	3135	Average risk, nulliparous	Chronic HTN, renal disease, DM, or other [unspecified] medical illnesses	60	Not specified 13 to 25 weeks; delivery	Placebo	20.5	IG: 100 CG: 100	White: 17.9 AA: 49.8 Hisp: 31.6	IG: 11.5 CG: 10.7	IG: 0 CG: 0

Table 3. Population, Study Design, and Intervention Characteristics, Average-Risk Population

Author, year	N rand	Included Population	Main medical history exclusions	Aspirin dosage, mg	Treatment time of day, pregnancy duration	Control condition	Mean age (range)	Nulliparous, %	Race/ ethnicity, %	Smoking, %	Chronic HTN, %
Subtil, 2003 ¹⁰¹ France, Belgium Good	3294	Average risk, nulliparous	Chronic HTN, antiphospholipid antibodies, lupus, contraindication to aspirin or other anticoagulant use (e.g., allergy, frequent hematomas or bleeding, hemorrhage during medical procedures, recent ulcer, severe asthma)	100	Not specified 14 to 20 weeks; 34 weeks	Placebo	24.7	IG: 100 CG: 100	NR	IG: 25.0 CG: 24.9	IG: 0 CG: 0

Abbreviations: AA = African American; CG = Control group; DM = Diabetes mellitus; Hisp = Hispanic; HTN = Hypertension; IG = Intervention group; NR = Not reported; PreE = Preeclampsia; PIH = Pregnancy-induced hypertension

Table 4. Patient Risk Factors and Clinical Tests Used to Identify Study Populations at Increased Risk for Preeclampsia

Author, Year	%Preeclampsia incidence in control group	Hx of preeclampsia or hypertension*	Maternal Age	Nulliparity	Multifetal gestation	Renal Disease	Metabolic disease	Hx of Stillbirth	Hx spontaneous abortion	Hx of SGA/IUGR	Diabetes	Prediction model†	Hemoglobin concentration	Positive rollover test	Angiotensin-II sensitivity	Abnormal Doppler readings
Benigni, 1989 ⁸⁷	NR	x						x		x						
Gallery, 1997 ⁹⁶	NR	x				x										
Scazzocchio, 2017 ^{†105}	4															x
CLASP, 1994 ⁹⁴	8	x	x		x	x										
Caspi, 1994 ⁹³	9				x											
Grab, 2000 ¹⁰⁰	10	x						x		x						
ASPRE, 2017 ⁵⁸	11	x										x				x
Viinikka, 1993 ⁹²	11	x														
Davies, 1995 ⁹⁵	12			x [§]									x			
Ayala, 2012 ⁵⁶	13	x	x	x	x		x		x							
Hermida, 1997 ⁹⁷	14	x	x	x	x		x		x							
Villa, 2012 ¹⁰³	18	x							x	x	x					x
McParland, 1990 ⁸⁹	19			x [§]												x
Yu, 2003 ¹⁰²	19															x
MFMU, 1998 ⁹⁸	20	x			x						x					
Morris, 1996 ¹⁰⁶	14			x												x
Schiff, 1989 ⁸⁸	23	x		x	x									x		
Wallenburg, 1986 ⁸⁶	30			x [§]											x	

Abbreviations: Hx = History; SGA/IUGR = Small for gestational age/intrauterine growth restriction; NR = Not reported

* Includes personal history of gestational hypertension, chronic hypertension, or preeclampsia

† Prediction model includes maternal risk factors, serum biomarkers, Doppler ultrasound

Table 4. Patient Risk Factors and Clinical Tests Used to Identify Study Populations at Increased Risk for Preeclampsia

‡Excluded individuals with a multifetal gestation or history of preeclampsia, hypertension, renal disease, or cardiovascular disease.

§Population limited to nulliparous individuals

¶Excluded criteria included individuals with history of hypertensive disease, renal and cardiovascular disease, and diabetes.

Table 5. Summary of Meta-Analysis Results

	No. studies reporting outcome (N Randomized)	Pooled Analysis No. studies (n analyzed)*	Pooled RR random effects model† (95% CI)	I ² %	Tau ²	Effect Range	Effect Median (IQR)
Perinatal mortality	15 (15,527)	11 (13,860)	0.79 (0.66, 0.96)	0.0	0.0	0.31 to 5.15 ARD: -6.3 to 2.9	0.98 (0.78 to 1.13) ARD: -0.0 (-0.6 to 0.4)
Preterm birth	13 (15,213)	13 (13,619)	0.80 (0.67, 0.95)	48.7	0.02	0.12 to 1.03 ARD: -19.5 to 0	0.65 (0.35 to 0.90) ARD: -5.7 (-12.9 to -3.0)
SGA/IUGR	16 (15,757)	16 (14,385)	0.82 (0.68, 0.99)	41.2	0.04	0.30 to 1.22 ARD: -25.7 to 4.9	0.63 (0.48 to 0.97) ARD: -4.6 (-8.9 to -0.2)
Preeclampsia	16 (15,757)	16 (14,093)	0.85 (0.75, 0.95)	0.0	0.0	0.07 to 1.43 ARD: -30.4 to 4.1	0.74 (0.43 to 0.90) ARD: -3.1 (-6.3 to -0.9)
Postpartum hemorrhage	11 (23,583)	9 (23,133)	1.03 (0.94, 1.12)	0.0	0.0	0.38 to 2.84 ARD: -4.2 to 9.1	1.02 (0.94 to 1.23) ARD: 0.2 (-0.4 to 0.9)
Placental abruption	13 (25,761)	10 (24,970)	1.15 (0.76, 1.72)	25.3	0.07	0.64 to 5.56 ARD: -0.6 to 1.8	1.21 (0.96 to 2.07) ARD: 0.3 (0 to 0.6)
Fetal intracranial bleeding	9 (23,959)	6 (23,719)	0.90 (0.51, 1.57)	19.2	0.06	0.17 to 2.06 ARD: -0.3 to 0.6	0.94 (0.74 to 1.08) ARD: 0.0 (-0.1 to 0.0)

Abbreviations: ARD = Absolute risk difference; CI = Confidence interval; IQR = Interquartile range; k = Number of studies; N = Number of observations; RR = Risk ratio; SGA/IUGR = Small for gestational age / Intrauterine growth restriction

*Studies that reported no events in both study arms were excluded from the pooled analysis.

†Restricted maximum likelihood model with Knapp-Hartung confidence intervals

Table 6. Summary of Evidence

Key Question	No. of Studies (No. of observations) Study Designs	Summary of Findings	Consistency and Precision	Other Limitations	Strength of Evidence	Applicability
<p>KQ1. Does aspirin reduce adverse maternal, perinatal, child, or combined health outcomes in pregnant persons at increased risk of preeclampsia?</p> <p>KQ1a. Does the effectiveness of aspirin for reducing adverse health outcomes vary by subpopulations defined by personal characteristics or preeclampsia risk factors?</p>	<p>18 RCTs 10 good quality, 8 fair quality (n= 15,908)</p>	<p>Aspirin was associated with a reduced risk of perinatal mortality (pooled RR 0.79, 95% CI 0.66, 0.96; I^2 0%), preterm birth (pooled RR 0.80, 95%CI 0.67, 0.95; I^2 49%), and SGA/IUGR (pooled RR 0.82, 95%CI 0.68, 0.99; I^2 41%) Limited subpopulation data, but within trial comparison suggested greater effectiveness in nonsmokers</p>	<p>Reasonably consistent* Reasonably precise**</p>	<p>Small study effects could not be ruled out for SGA/IUGR and preterm delivery Rare maternal health outcomes, such as eclampsia and maternal mortality, occurred too infrequently to estimate preventive effectiveness Subgroup effects limited by modest number included studies and few within trial subgroup analyses reported</p>	<p>Moderate for perinatal health benefits</p>	<p>Studies in prenatal care settings in US or comparable settings, however mostly White participants Different criteria for identifying at risk populations Aspirin dose 50-150mg a day</p>

Table 6. Summary of Evidence

Key Question	No. of Studies (No. of observations) Study Designs	Summary of Findings	Consistency and Precision	Other Limitations	Strength of Evidence	Applicability
<p>KQ2. Does aspirin prevent preeclampsia in pregnant persons at increased risk for preeclampsia?</p> <p>KQ2a. Does the effectiveness of aspirin for reducing preeclampsia vary by subpopulations defined by personal characteristics or preeclampsia risk factors?</p>	<p>16 RCTs 10 good quality 6 fair quality</p> <p>(n= 15,767)</p>	<p>Aspirin was associated with a statistically significant reduction in the risk of preeclampsia compared to placebo (pooled RR, 0.85 [95% CI, 0.75 to 0.95], <i>I</i>² 0%)</p> <p>No evidence of statistical difference in the magnitude of preeclampsia risk reduction related to the timing of treatment initiation, the dosage of aspirin used, or personal characteristics such as smoking history, parity, and BMI</p>	<p>Reasonably consistent evidence for aspirin benefit</p> <p>Reasonably precise 15% reduced risk of preeclampsia associated with daily aspirin use</p>	<p>Small study effects could not be ruled out and might lead to some overestimation of pooled effect sizes</p> <p>Confounding of study size with other study and participant characteristics could influence subgroup comparisons</p> <p>Subgroup effects limited by modest number included studies and few within trial subgroup analyses reported</p>	<p>Moderate</p>	<p>Studies in prenatal care settings in US or comparable settings, however mostly White participants</p> <p>Different criteria for identifying at risk populations</p> <p>Aspirin dose 50-150mg a day</p>

Table 6. Summary of Evidence

Key Question	No. of Studies (No. of observations) Study Designs	Summary of Findings	Consistency and Precision	Other Limitations	Strength of Evidence	Applicability
<p>KQ3. What are the harms of aspirin use to prevent preeclampsia during pregnancy?</p> <p>KQ3a. Do the harms of aspirin use to prevent preeclampsia vary by subpopulations defined by personal characteristics or preeclampsia risk factors?</p>	<p>21 RCTs (16 increased risk and 5 average risk populations)</p> <p>14 good quality 7 fair quality (n = 26,757)</p>	<p>Studies conducted among average and increased risk populations did not find any clear evidence of harms associated with daily aspirin use (<150mg) taken during the second or third trimester of pregnancy. No difference in harms by the dosage or timing of aspirin or for specific populations were identified in limited subgroup comparisons.</p> <p>Bleeding harms were uncommon and showed null effects for differences in risk of postpartum hemorrhage (pooled RR 1.03, [95% CI, 0.94, 1.12] I² 0%, k=11), or fetal intracranial bleeding (pooled RR, 0.90 [95% CI, 0.51, 1.57]; I² 19%, k=9) were found. The result for placental abruption (pooled RR, 1.15 [95% CI, 0.76, 1.72] I² 25%, k=13) was also null. Longer term followup from one large trial found no difference in child developmental outcomes for aspirin compared with placebo exposed groups. No differences were found within a limited set of studies reporting other rare perinatal harms.</p>	<p>Reasonably consistent evidence of null effects for bleeding harms of daily aspirin, especially among pregnant individuals at increased preeclampsia risk.</p> <p>Reasonably precise evidence for null effects, but less precise for especially rare harms.</p>	<p>Reported harms were rare and not consistently reported across studies.</p>	<p>Moderate for no difference in bleeding harms between groups, low for very rare or inconsistently reported harms.***</p>	<p>Studies in prenatal care settings in US or comparable settings, however mostly White participants Different criteria for identifying at risk populations Aspirin dose 50-150mg a day Harms from trials in average risk and increased risk populations</p>

*Direction and magnitude of effects within and across for important perinatal health outcomes, with low to moderate statistical heterogeneity

** Estimates of approximately 20 percent reduced risk; similar effect magnitudes for the perinatal outcomes that could be quantitatively synthesized

*** Postpartum hemorrhage >1000 mL, longer term developmental harms, congenital malformations.

Table 7. Theoretical Number Needed to Treat (NNT) Across Different Levels of Outcome Incidence

Outcome	% Outcome incidence in population	Pooled RR estimate	% Absolute risk reduction	NNT (95% CI)
SGA/IUGR	7	0.82	-1	79 (45, 1429)
	17	0.82	-3	33 (18, 588)
	24	0.82	-4	23 (13, 417)
Preterm Birth	10	0.80	-2	50 (30, 200)
	17	0.80	-3	29 (18, 118)
	31	0.80	-6	16 (10, 65)
Perinatal mortality	1	0.79	-0.2	476 (294, 2500)
	6	0.79	-1	79 (49, 417)
Preeclampsia	9	0.85	-1	74 (44, 222)
	13	0.85	-2	51 (31, 154)
	19	0.85	-3	35 (21, 105)

Abbreviations: NNH = Number needed to harm; NNT= Number needed to treat; RR = Relative risk; SGA/IUGR = Small for gestational age/Intrauterine growth restriction

Table 8. Preeclampsia Risk Factors Based on Patient Medical History*

Level of Risk	Risk Factors
Highest risk[†]	Prior preeclampsia Multifetal gestation Chronic hypertension Pregestational type I or II diabetes
Increased risk[‡]	Nulliparous, having never given birth Obesity (i.e., BMI >30 kg/m ²) Renal disease Autoimmune disease (i.e., systemic lupus erythematosus, antiphospholipid syndrome) Family history of preeclampsia (i.e., mother, sister) Sociodemographic characteristics (i.e., Black race, low socioeconomic status) Age ≥35 Personal pregnancy history (e.g., born low birth weight or small for gestational age, previous adverse pregnancy outcome, greater than 10 year pregnancy interval) <i>In vitro</i> conception
Lower risk	Prior uncomplicated term delivery and absence of risk factors

*Includes only risk factors that can be obtained from the patient medical history. Clinical measures such as serum biomarkers and uterine artery Doppler ultrasound have also been used in some trials to identify an increased risk population. A risk prediction algorithm combining clinical risk factors and biochemical and imaging markers was used in one trial for risk assessment in the first trimester among individuals with singleton pregnancies.

[†]Risk factors consistently and independently associated with the greatest risk of preeclampsia; preeclampsia risk is at least twice as high as rates observed in an unselected population.

[‡] Risk factors independently associated with moderate preeclampsia risk, some more consistently than others.

Clinical judgment of risk for the specific patient and population or the presence of more than one risk factor could be used to identify a population at risk for preeclampsia at levels observed in the trial evidence on aspirin prophylaxis

Appendix A. Detailed Methods

Sources Searched	Number of items YEAR -present
MEDLINE	729
PUBMED	56
Cochrane Central Register of Controlled Clinical Trials	520
Embase	3192

Key:

/ = MeSH subject heading

\$ = truncation

ti = word in title

ab = word in abstract

pt = publication type

* = truncation

kw = keyword

Ink = subheading

de = index term

exp = explode

py = publication year

MEDLINE

Database: Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations, Daily and Versions(R) <1946 to July 22, 2019>

Search Strategy:

-
- 1 Pregnancy/
 - 2 Pregnant Women/
 - 3 pregnan\$.ti,ab.
 - 4 Pre-Eclampsia/
 - 5 Hypertension, Pregnancy-Induced/
 - 6 Eclampsia/
 - 7 HELLP Syndrome/
 - 8 preeclamp\$.ti,ab.
 - 9 eclamp\$.ti,ab.
 - 10 ((edema or oedema or proteinuria or hypertension or hypertensive) adj5 (gestosis or gestoses or antenatal or prenatal or gestation\$ or perinatal)).ti,ab.
 - 11 eph gestosis.ti,ab.
 - 12 (tox?emi\$ adj3 (eph or pregnan\$)).ti,ab.
 - 13 HELLP.ti,ab.
 - 14 or/1-13
 - 15 exp Aspirin/
 - 16 Platelet Aggregation Inhibitors/
 - 17 aspirin.ti,ab.
 - 18 acetylsalicylic acid.ti,ab.
 - 19 acetyl salicylic acid.ti,ab.
 - 20 antiplatelet\$.ti,ab.
 - 21 anti platelet\$.ti,ab.
 - 22 (platelet\$ adj3 (inhibit\$ or antiaggregant\$ or antagonist\$)).ti,ab.

Appendix A. Detailed Methods

- 23 or/15-22
- 24 (clinical trial or controlled clinical trial or meta analysis or randomized controlled trial or pragmatic clinical trial).pt.
- 25 clinical trials as topic/ or controlled clinical trials as topic/ or randomized controlled trials as topic/
- 26 Meta-Analysis as Topic/
- 27 control groups/ or double-blind method/ or single-blind method/
- 28 clinical trial\$.ti,ab.
- 29 controlled trial\$.ti,ab.
- 30 random\$.ti,ab.
- 31 (metaanaly\$ or meta analy\$).ti,ab.
- 32 trial.ti.
- 33 placebo.ti,ab.
- 34 or/24-33
- 35 14 and 23 and 34
- 36 Pregnancy/
- 37 (pregnan\$ or postpartum or post partum or antenatal or prenatal or gestation\$ or perinatal or maternal or f?etal or f?etus or neonat\$).ti,ab.
- 38 36 or 37
- 39 adverse effects.fs.
- 40 harm\$.ti,ab.
- 41 adverse.ti,ab.
- 42 Hemorrhage/
- 43 h?emorrhag\$.ti,ab.
- 44 bleed\$.ti,ab.
- 45 blood loss.ti,ab.
- 46 exp Gastrointestinal Diseases/
- 47 (gastro\$ or gastric or epigastric or instestinal).ti,ab.
- 48 (behavio* or development\$).ti,ab.
- 49 or/39-48
- 50 38 and 49
- 51 Pregnancy complications/
- 52 Maternal death/
- 53 Maternal mortality/
- 54 Maternal Exposure/
- 55 Fetal death/
- 56 Fetal mortality/
- 57 Congenital abnormalities/
- 58 Postpartum hemorrhage/
- 59 Abruptio Placentae/
- 60 Abortion, Spontaneous/
- 61 Ductus Arteriosus, Patent/
- 62 Chorioamnionitis/
- 63 ((pregnan\$ or maternal or f?etal or f?etus or neonat\$) adj3 complication\$).ti,ab.
- 64 ((congenital or birth or f?etal or f?etus) adj3 (defect\$ or abnormal\$ or anomal\$ or malform\$)).ti,ab.
- 65 ((maternal or f?etal or f?etus or perinatal) adj3 (death\$ or mortality)).ti,ab.
- 66 miscarr\$.ti,ab.
- 67 (spontaneous abortion\$ or spontaneous labo?r).ti,ab.
- 68 abruptio\$.ti,ab.

Appendix A. Detailed Methods

- 69 ductus arteriosus.ti,ab.
- 70 (chorioamnionitis or intraamniotic infection or intra-amniotic infection).ti,ab.
- 71 or/51-70
- 72 50 or 71
- 73 23 and 72
- 74 35 or 73
- 75 limit 74 to (english language and yr="2012 -Current")
- 76 75 not (animals/ not humans/)
- 77 remove duplicates from 76

PUBMED- [publisher supplied references only]

- #1 Search pregnan*[tiab] OR preeclamp*[tiab] OR eclamp*[tiab]
- #2 Search aspirin[tiab] OR acetylsalicylic acid[tiab] OR acetyl salicylic acid[tiab] OR antiplatelet\$[tiab] OR anti platelet\$[tiab] OR (platelet*[tiab] AND (inhibit*[tiab] OR antiaggregant*[tiab] OR antagonist*))
- #3 Search (#1 AND #2)
- #4 Search (#3 AND publisher[sb])
- #5 Search (#3 AND publisher[sb]) Filters: Publication date from 2012/01/01
- #6 Search (#3 AND publisher[sb]) Filters: Publication date from 2012/01/01; English

Cochrane Central Register of Controlled Clinical Trials (CENTRAL)

- #1 pregnan*:ti,ab,kw or preeclamp*:ti,ab,kw or eclamp*:ti,ab,kw
- #2 ((edema:ti,ab or oedema:ti,ab or proteinuria:ti,ab or hypertension:ti,ab or hypertensive:ti,ab) NEAR/5 (gestosis:ti,ab or gestoses:ti,ab or antenatal:ti,ab or prenatal:ti,ab or gestation*:ti,ab or perinatal:ti,ab))
- #3 eph gestosis:ti,ab
- #4 (tox*emi*:ti,ab NEAR/3 (eph:ti,ab or pregnan*:ti,ab))
- #5 #1 OR #2 OR #3 OR #4
- #6 aspirin:ti,ab,kw or "acetylsalicylic acid":ti,ab,kw or antiplatelet*:ti,ab,kw or anti next platelet*:ti,ab,kw
- #7 platelet*:ti,ab AND (inhibit*:ti,ab or antiaggregant*:ti,ab or antagonist*:ti,ab)
- #8 #6 OR #7
- #9 #5 AND #8 with Publication Year from 2012 to 2019, in Trials

Embase

- | No. | Query | Results |
|-----|--|---------|
| #1 | 'pregnancy'/de | |
| #2 | 'pregnant woman'/de | |
| #3 | pregnan*:ab,ti | |
| #4 | 'eclampsia and preeclampsia'/exp | |
| #5 | 'pregnancy toxemia'/de | |
| #6 | 'hellp syndrome'/de | |
| #7 | preeclamp*:ti,ab | |
| #8 | eclamp*:ti,ab | |
| #9 | ((edema OR oedema OR proteinuria OR hypertension OR hypertensive) NEAR/5 (gestosis OR gestoses OR antenatal OR prenatal OR gestation* OR perinatal)):ti,ab | |
| #10 | 'eph gestosis':ti,ab | |
| #11 | (tox\$emi* NEAR/3 (eph OR pregnan*)):ti,ab | |

Appendix A. Detailed Methods

- #12 hellp:ti,ab
- #13 #1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12
- #14 'acetylsalicylic acid'/de
- #15 'antithrombocytic agent'/de
- #16 aspirin:ti,ab
- #17 'acetylsalicylic acid':ti,ab
- #18 'acetyl salicylic acid':ti,ab
- #19 antiplatelet*:ti,ab
- #20 'anti platelet*':ti,ab
- #21 (platelet* NEAR/3 (inhibit* OR antiaggregant* OR antagonist*)):ti,ab
- #22 #14 OR #15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21
- #23 'clinical trial'/de
- #24 'controlled clinical trial'/de
- #25 'randomized controlled trial'/exp
- #26 'single blind procedure'/de
- #27 'double blind procedure'/de
- #28 'crossover procedure'/de
- #29 'placebo'/de
- #30 'randomized controlled trial (topic)'/de
- #31 'clinical trial (topic)'/de
- #32 'controlled clinical trial (topic)'/de
- #33 'meta analysis'/exp
- #34 'meta analysis (topic)'/de
- #35 'clinical trial*':ti,ab
- #36 'controlled trial*':ti,ab
- #37 random*:ti,ab
- #38 metaanaly*:ti,ab OR 'meta analy*':ti,ab
- #39 trial:ti
- #40 placebo:ti,ab
- #41 #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37 OR #38 OR #39 OR #40
- #42 #13 AND #22 AND #41
- #43 pregnan*:ti,ab OR postpartum:ti,ab OR 'post partum':ti,ab OR antenatal:ti,ab OR prenatal:ti,ab OR gestation*:ti,ab OR perinatal:ti,ab OR maternal:ti,ab OR f\$etal:ti,ab OR f\$etus:ti,ab OR neonat*:ti,ab
- #44 #1 OR #43
- #45 'adverse drug reaction'/lnk OR 'complication'/lnk OR 'drug toxicity'/lnk OR 'side effect'/lnk
- #46 harm*:ti,ab
- #47 adverse:ti,ab
- #48 'bleeding'/de
- #49 h\$emorrhag*:ti,ab
- #50 bleed*:ti,ab
- #51 blood AND loss:ti,ab
- #52 'gastrointestinal disease'/de
- #53 gastro*:ti,ab OR gastric:ti,ab OR epigastric:ti,ab OR instestinal:ti,ab
- #54 behavio*:ti,ab OR development*:ti,ab
- #55 #45 OR #46 OR #47 OR #48 OR #49 OR #50 OR #51 OR #52 OR #53 OR #54
- #56 #44 AND #55
- #57 'pregnancy disorder'/de

Appendix A. Detailed Methods

#58 'pregnancy complication'/de
#59 'maternal death'/de
#60 'maternal mortality'/de
#61 'maternal exposure'/de
#62 'fetus death'/de
#63 'fetus mortality'/de
#64 'congenital disorder'/de
#65 'postpartum hemorrhage'/de
#66 'solutio placentae'/de
#67 'spontaneous abortion'/de
#68 'patent ductus arteriosus'/de
#69 'chorioamnionitis'/exp
#70 ((pregnan* OR maternal OR f\$etal OR f\$etus OR neonat*) NEAR/3 complication*):ti,ab
#71 ((congenital OR birth OR f\$etal OR f\$etus) NEAR/3 (defect* OR abnormal* OR anomal* OR malform*)):ti,ab
#72 ((maternal OR f\$etal OR f\$etus OR perinatal) NEAR/3 (death* OR mortality)):ti,ab
#73 miscarr*:ti,ab
#74 'spontaneous abortion*':ti,ab OR 'spontaneous labo\$r':ti,ab
#75 abruptio*:ti,ab OR 'solutio placentae':ti,ab
#76 'ductus arteriosus':ti,ab
#77 chorioamnionitis:ti,ab OR 'intraamniotic infection':ti,ab OR 'intra-amniotic infection':ti,ab
#78 #57 OR #58 OR #59 OR #60 OR #61 OR #62 OR #63 OR #64 OR #65 OR #66 OR #67 OR #68 OR #69 OR #70 OR #71 OR #72 OR #73 OR #74 OR #75 OR #76 OR #77
#79 #56 OR #78
#80 #22 AND #79
#81 #42 OR #80
#82 (#42 OR #80) AND [english]/lim
#83 #82 AND (2012:py OR 2013:py OR 2014:py OR 2015:py OR 2016:py OR 2017:py OR 2018:py OR 2019:py) AND [embase]/lim
#84 #83 NOT (('animal'/exp OR 'nonhuman'/de) NOT 'human'/exp)

Appendix A Table 1. Inclusion and Exclusion Criteria

	Inclusion	Exclusion
Populations	<p>KQs 1, 2 (Efficacy): Pregnant persons at increased risk for preeclampsia based on:</p> <ul style="list-style-type: none"> • Personal sociodemographic characteristics • Medical history • Diagnostic measurements or assays (e.g., uterine artery Doppler, biomarkers) • Risk prediction model <p>KQ 3 (Harms): Pregnant persons, fetuses, infants, and children.</p>	<p>Nonhuman populations; nonpregnant persons; studies that only/exclusively include persons seeking fertility treatment; and other selected nongeneralizable populations</p>
Disease/condition	<p>Primary prevention of preeclampsia</p>	<p>Trials of aspirin aimed at preventing other complications of pregnancy (e.g., stillbirth)</p>
Setting	<p>Countries categorized as “very high” on the 2017 Human Development Index (as defined by the United Nations Development Programme)</p>	<p>Countries not categorized as “very high” on the 2017 Human Development Index, as there is concern for nutritional deficiencies in developing countries</p>
Interventions	<p>Aspirin (≥50 mg)</p>	<p>Nonaspirin antiplatelet medications or aspirin combined with other potentially active interventions</p>
Comparisons	<p>Placebo or no treatment</p>	<p>Any active substance or intervention (e.g., nonaspirin medication, dietary supplements, dietary change, bed rest, or weight loss)</p>
Outcomes	<p>Maternal outcomes:</p> <ul style="list-style-type: none"> • Preeclampsia; Preeclampsia with severe features • Hemolysis, elevated liver enzymes, and low platelet count (HELLP) syndrome • Eclampsia, puerperal cerebrovascular disorder, cerebrovascular hemorrhage, edema, or embolus • Renal or hepatic injury/failure • Pulmonary edema, adult respiratory distress syndrome • Disseminated intravascular coagulation • Mental health diagnoses or symptoms • Maternal mortality • Measures of well-being or quality of life <p>Potential treatment harms:</p> <ul style="list-style-type: none"> • Abruptio placentae • Postpartum hemorrhage • Gastrointestinal complications (e.g., bleeding ulcer) <p>Fetal/neonatal/child outcomes:</p> <ul style="list-style-type: none"> • Preterm birth (<37 weeks): late preterm birth (34-36 weeks), moderate perterm birth, (32-34 weeks), very preterm birth (<32 weeks), extremely preterm birth (<28 weeks) • Mean gestational age • Low birth weight • Intrauterine growth restriction/small for gestational age (<10th percentile weight for gestational age) • Stillbirth or neonatal mortality <p>Potential treatment harms:</p> <ul style="list-style-type: none"> • Intracranial fetal bleeding • Fetal malformations • Nonclosure of the ductus arteriosus • Chorioamnionitis • Child behavioral or developmental problems 	<ul style="list-style-type: none"> • Length of hospital stay (without indication) • Intensive care unit admission • Neonatal intensive care unit admission

Appendix A Table 1. Inclusion and Exclusion Criteria

	Inclusion	Exclusion
Study Designs	<p>KQs 1, 2 (Efficacy): Randomized, controlled trial, individual participant data meta-analysis of trials</p> <p>KQ 3 (Harms): Randomized, controlled trial or comparative cohort studies, individual participant data meta-analysis of trials</p>	<p>KQs 1, 2 (Efficacy): Any nonrandomized controlled trial</p> <p>KQ 3 (Harms): Editorials, narrative review, commentary, postmarketing surveillance, or case reports</p>
Study Quality	Good- and fair-quality studies	Poor-quality studies
Language	English	Languages other than English


Abbreviations: KQ = Key Question; USPSTF = U.S. Preventative Services Task Force

Appendix A Table 2. Study Design-Specific Quality Rating Criteria

Study Design	Adapted Quality Criteria
Randomized and non-randomized controlled trials, adapted from the U.S. Preventive Services Task Force methods ⁷³	<p>Bias arising in the randomization process or due to confounding</p> <ul style="list-style-type: none"> • Valid random assignment/random sequence generation method used • Allocation concealed • Balance in baseline characteristics <p>Bias in selecting participants into the study</p> <ul style="list-style-type: none"> • CCT only: No evidence of biased selection of sample <p>Bias due to departures from intended interventions</p> <ul style="list-style-type: none"> • Fidelity to the intervention protocol • Low risk of contamination between groups • Participants were analyzed as originally allocated <p>Bias from missing data</p> <ul style="list-style-type: none"> • No, or minimal, post-randomization exclusions • Outcome data are reasonably complete and comparable between groups • Reasons for missing data are similar across groups • Missing data are unlikely to bias results <p>Bias in measurement of outcomes</p> <ul style="list-style-type: none"> • Blinding of outcome assessors • Outcomes are measured using consistent and appropriate procedures and instruments across treatment groups • No evidence of inferential statistics <p>Bias in reporting results selectively</p> <ul style="list-style-type: none"> • No evidence that the measures, analyses, or subgroup analyses are selectively reported

* Good quality studies generally meet all quality criteria. Fair quality studies do not meet all the criteria but do not have critical limitations that could invalidate study findings. Poor quality studies have a single fatal flaw or multiple important limitations that could invalidate study findings. Critical appraisal of studies using *a priori* quality criteria are conducted independently by at least two reviewers. Disagreements in final quality assessment are resolved by consensus, and, if needed, consultation with a third independent reviewer

Appendix B Figure 1. Literature Flow Diagram



*Studies may appear in more than one Key Question

Appendix C. Included Studies

Below is a list of included studies and their ancillary publications (indented below main results publication):

Ayala, DE, Ucieda, R, et al. Chronotherapy With Low-Dose Aspirin for Prevention of Complications in Pregnancy. *Chronobiol.Int.* 30(1-2): 260-279. 2012.
<https://dx.doi.org/10.3109/07420528.2012.717455>.

Benigni, A, Gregorini, G, et al. Effect of low-dose aspirin on fetal and maternal generation of thromboxane by platelets in women at risk for pregnancy-induced hypertension. *N Engl J Med.* 321(6): 357-362. 1989. <https://dx.doi.org/>

Caritis, Steve, Sibai, Baha, et al. Low-Dose Aspirin to Prevent Preeclampsia in Women at High Risk. *N Engl J Med.* 338(11): 701-705. 1998.
<https://dx.doi.org/10.1056/NEJM199803123381101>

Abramovici, A, Jauk, V, et al. Low-dose aspirin, smoking status, and the risk of spontaneous preterm birth. *Am J Perinatol.* 32(5): 445-50. 2015. PMID: 25261702.
<https://dx.doi.org/10.1055/s-0034-1390352>

Adkins, K, Allshouse, AA, et al. Impact of aspirin on fetal growth in diabetic pregnancies according to White classification. *Am J Obstet Gynecol.* 217(4): 465.e1-465.e5. 2017. PMID: 28599894. <https://dx.doi.org/10.1016/j.ajog.2017.05.062>

Cantu, JA, Jauk, VR, et al. Is low-dose aspirin therapy to prevent preeclampsia more efficacious in non-obese women or when initiated early in pregnancy?. *Journal of Maternal-Fetal & Neonatal Medicine.* 28(10): 1128-32. 2015. PMID: 25048750.
<https://dx.doi.org/10.3109/14767058.2014.947258>

Tolcher, MC, Sangi-Haghpeykar, H, et al. Low-dose aspirin (ASA) for preeclampsia prevention among high-risk women by ethnicity and race. *Am J Obstet Gynecol.* 218(1): S564. 2018.

Caspi, E, Raziel, A, et al. Prevention of pregnancy-induced hypertension in twins by early administration of low-dose aspirin: a preliminary report. *Am J Reprod Immunol.* 31(1): 19-24. 1994.

CLASP: A randomised trial of low-dose aspirin for the prevention and treatment of pre-eclampsia. *Lancet.* 343(8898): 619. 1994.

Clasp Collaborative Group. Low dose aspirin in pregnancy and early childhood development: follow up of the collaborative low dose aspirin study in pregnancy. CLASP collaborative group 50. *Br J Obstet Gynaecol.* 102(11): 861-868. 1995.

Davies, NJ, Gazvani, MR, et al. Low-Dose Aspirin in the Prevention of Hypertensive Disorders of Pregnancy in Relatively Low-Risk Nulliparous Women. *Hypertension in Pregnancy.* 14(1): 49-55. 1995.

Gallery, EileenDM, Ross, MargaretR, et al. Low-Dose Aspirin in High-Risk Pregnancy?. *Hypertension in Pregnancy.* 16(2): 229-238. 1997.
<https://dx.doi.org/10.3109/10641959709031640>

Appendix C. Included Studies

- Leslie, GI, Gallery, ED, et al. Neonatal outcome in a randomized, controlled trial of low-dose aspirin in high-risk pregnancies. *J Paediatr Child Health*. 31(6): 549-552. 1995.
- Grab, D, Paulus, WE, et al. Effects of low-dose aspirin on uterine and fetal blood flow during pregnancy: results of a randomized, placebo-controlled, double-blind trial. *Ultrasound Obstet Gynecol*. 15(1): 19-27. 2000.
- Hauth, JC, Goldenberg, RL, et al. Low-dose aspirin therapy to prevent preeclampsia. *Am J Obstet Gynecol*. 168(4): 1083-1091. 1993.
- Hermida, RC, Ayala, DE, et al. Time-dependent effects of low-dose aspirin administration on blood pressure in pregnant women. *Hypertension*. 30(3 Pt 2): 589-595. 1997.
- McParland, P, Pearce, JM, et al. Doppler ultrasound and aspirin in recognition and prevention of pregnancy-induced hypertension. *Lancet*. 335(8705): 1552-1555. 1990.
- Mone, F, Mulcahy, C, et al. Trial of feasibility and acceptability of routine low-dose aspirin versus Early Screening Test indicated aspirin for pre-eclampsia prevention (TEST study): a multicentre randomised controlled trial. *BMJ Open*. 8(7): e022056. 2018. PMID: 30056389. <https://dx.doi.org/10.1136/bmjopen-2018-022056>
- Mone, F, Mulcahy, C, et al. Evaluation of the Effect of Low-Dose Aspirin on Biochemical and Biophysical Biomarkers for Placental Disease in Low-Risk Pregnancy: Secondary Analysis of a Multicenter RCT. *Am J Perinatol*. 2019. PMID: 30646422. <https://dx.doi.org/10.1055/s-0038-1677476>
- Morris, JM, Fay, RA, et al. A randomized controlled trial of aspirin in patients with abnormal uterine artery blood flow. *Obstet Gynecol*. 87(1): 74-8. 1996.
- Rolnik, Wright DL, Poon D, O'Gorman LC, et al. Aspirin Versus Placebo in Pregnancies at High Risk for Preterm Preeclampsia. *N Engl J Med*. 377(7): 613-622. 2017. <https://dx.doi.org/10.1056/NEJMoa1704559>
- Ling, HZ, Jara, PG, et al. Maternal cardiac function in women at high-risk for pre-eclampsia treated with 150 mg aspirin or placebo: an observational study. *Bjog*. 2020.
- Poon, LC, Wright, D, et al. Aspirin for Evidence-Based Preeclampsia Prevention trial: effect of aspirin in prevention of preterm preeclampsia in subgroups of women according to their characteristics and medical and obstetrical history. *Am J Obstet Gynecol*. 217(5): 585.e1-585.e5. 2017. <https://dx.doi.org/10.1016/j.ajog.2017.07.038>
- Rotchell, YE, Cruickshank, JK, et al. Barbados Low Dose Aspirin Study in Pregnancy (BLASP): a randomised trial for the prevention of pre-eclampsia and its complications. *Br J Obstet Gynaecol*. 105(3): 286-292. 1998.
- Scazzocchio, E, Oros, D, et al. Impact of aspirin on trophoblastic invasion in women with abnormal uterine artery Doppler at 11-14 weeks: a randomized controlled study. *Ultrasound Obstet Gynecol*. 49(4): 435-441. 2017. <https://dx.doi.org/10.1002/uog.17351>
- Schiff, E, Peleg, E, et al. The use of aspirin to prevent pregnancy-induced hypertension and lower the ratio of thromboxane A2 to prostacyclin in relatively high risk pregnancies. *N Engl J Med*. 321(6): 351-356. 1989.

Appendix C. Included Studies

Sibai, BM, Caritis, SN, et al. Prevention of preeclampsia with low-dose aspirin in healthy, nulliparous pregnant women. The National Institute of Child Health and Human Development Network of Maternal-Fetal Medicine Units. *N Engl J Med.* 329(17): 1213-1218. 1993.

Tolcher, MC, Sangi-Haghpeykar, H, et al. Race and ethnicity varies both efficacy and attributable risk of low-dose aspirin (ASA) for preeclampsia prevention among low-risk women. *Am J Obstet Gynecol.* 218(1): S203. 2018.

Subtil, D, Goeusse, P, et al. Aspirin (100 mg) used for prevention of pre-eclampsia in nulliparous women: the Essai Regional Aspirine Mere-Enfant study (Part 1). *BJOG.* 110(5): 475-484. 2003.

Viinikka, L, Hartikainen-Sorri, AL, et al. Low dose aspirin in hypertensive pregnant women: effect on pregnancy outcome and prostacyclin-thromboxane balance in mother and newborn. *Br J Obstet Gynaecol.* 100(9): 809-815. 1993.

Villa, P, Kajantie, E, et al. Aspirin in the prevention of pre-eclampsia in high-risk women: a randomised placebo-controlled PREDO Trial and a meta-analysis of randomised trials. *BJOG.* 120(1): 64-74. 2013.

Wallenburg, HC, Dekker, GA, et al. Low-dose aspirin prevents pregnancy-induced hypertension and pre-eclampsia in angiotensin-sensitive primigravidae. *Lancet.* 1(8471): 1-3. 1986.

Yu, CK, Papageorgiou, AT, et al. Randomized controlled trial using low-dose aspirin in the prevention of pre-eclampsia in women with abnormal uterine artery Doppler at 23 weeks' gestation. *Ultrasound Obstet Gynecol.* 22(3): 233-239. 2003.

Appendix D. Excluded Studies

Reason for Exclusion*
E1. Relevance E1a. Not primary data (e.g., editorial, narrative review)
E2. Study design E2a. Single arm cohort, case-control study
E3. Setting E3a. Study country not a “very high” HDI
E4. Population E4a. Population not at increased risk (e.g. nulliparous otherwise healthy women)
E5. Intervention E5a. Aspirin plus other medication E5b. Intervention other than aspirin
E6. Comparator E6a. Comparative effectiveness – active comparator
E7. No relevant outcomes E7a. Secondary analysis of included study with no relevant outcomes E7b. Preliminary reporting, incomplete information
E8. Poor study quality
E9. Non-English language

*Assigned at full-text phase

Appendix D. Excluded Studies

1. Abramovici, A, Jauk, V, et al. Analysis of low-dose aspirin by smoking status for select perinatal outcomes. *Am J Obstet Gynecol.* 208(1): S261. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.779> **KQ1E7b, KQ2E7b, KQ3E7b**
2. Abramovici, A, Jauk, V, et al. Low-dose aspirin is associated with a reduction in spontaneous preterm birth among non-smokers. *Am J Obstet Gynecol.* 208(1): S260-S261. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.778> **KQ1E7b, KQ2E7b, KQ3E7b**
3. Adkins, K, Allshouse, A, et al. Impact of low-dose aspirin on fetal growth in diabetic pregnancies: the importance of white classification. *Am J Obstet Gynecol.* 216(1): S292-S293. 2017. **KQ1E7b, KQ2E7b, KQ3E7b**
4. Allshouse, AA, Jessel, RH, et al. The impact of low-dose aspirin on preterm birth: secondary analysis of a randomized controlled trial. *J Perinatol.* 36(6): 427-31. 2016. PMID: 26890552. <https://dx.doi.org/10.1038/jp.2016.3> **KQ1E2, KQ2E2, KQ3E2**
5. Ananth, C, Coletta-Lucas, J, et al. Timing of adverse neonatal outcomes in pregnancies complicated by preeclampsia and small for gestational age births in a cohort of high-risk pregnancies. *Am J Obstet Gynecol.* 208(1): S264. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.786> **KQ1E7b, KQ2E7b, KQ3E7b**
6. Anca-Daniela, S, Banica, R, et al. Low dose aspirin for preventing fetal growth restriction: A randomised trial. *J Perinat Med.* 43. 2015. <https://dx.doi.org/10.1515/jpm-2015-2002> **KQ1E7b, KQ2E7b, KQ3E7b**
7. Anderson, S, Jauk, V, et al. Mid-trimester blood pressures below 140/90 and risk of small-for-gestational age (SGA) in pregnancies complicated by chronic hypertension. *Am J Obstet Gynecol.* 212(1): S345-S346. 2015. <https://dx.doi.org/10.1016/j.ajog.2014.10.914> **KQ1E7b, KQ2E7b, KQ3E7b**
8. Andrikopoulou, M, Purisch, SE, et al. Beyond preeclampsia: Low dose aspirin reduces spontaneous preterm birth. *Am J Obstet Gynecol.* 218(1): S10. 2018. **KQ1E4a, KQ2E4a, KQ3E7a**
9. Andrikopoulou, M, Purisch, SE, et al. Low-dose aspirin is associated with reduced spontaneous preterm birth in nulliparous women. *Am J Obstet Gynecol.* 219(4): 399.e1-399.e6. 2018. PMID: 29913174. <https://dx.doi.org/10.1016/j.ajog.2018.06.011> **KQ1E4a, KQ2E4a, KQ3E7a**
10. Ankumah, NA, Cantu, J, et al. Risk of adverse pregnancy outcomes in women with mild chronic hypertension before 20 weeks of gestation. *Obstet Gynecol.* 123(5): 966-72. 2014. PMID: 24785847. <https://dx.doi.org/10.1097/AOG.0000000000000205> **KQ1E2, KQ2E2, KQ3E2**
11. Ankumah, NA, Tita, A, et al. Pregnancy outcome vary by blood pressure level in women with mild-range chronic hypertension. *Am J Obstet Gynecol.* 208(1): S261-S262. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.780> **KQ1E7b, KQ2E7b, KQ3E7b**
12. Askie, L, Duley, L. Associations between the timing and dosing of aspirin prophylaxis and term and preterm preeclampsia. *BMJ Evid Based Med.* 2018. PMID: 29880699. <https://dx.doi.org/10.1136/bmjebm-2018-110931> **KQ1E1a, KQ2E1a, KQ3E1a**
13. Askie, LM, Duley, L, et al. Antiplatelet agents for prevention of pre-eclampsia: a meta-analysis of individual patient data. *Lancet.* 369(9575): 1791-1798. 2007. [https://dx.doi.org/10.1016/S0140-6736\(07\)60712-0](https://dx.doi.org/10.1016/S0140-6736(07)60712-0) **KQ1E5, KQ2E5, KQ3E5**
14. August, P, Helseth, G, et al. Sustained release, low-dose aspirin ameliorates but does not prevent preeclampsia (PE)

Appendix D. Excluded Studies

- in a high risk population. Proceedings of the 9th International Congress, International Society for the Study of Hypertension. 72. 1994. **KQ1E7b, KQ2E7b, KQ3E7b**
15. Ayala, DE, Hermida, RC. Chronotherapy with low-dose aspirin for prevention of complications in pregnancy. *Cardiology (Switzerland)*. 125. 410. 2013. <https://dx.doi.org/10.1159/000354059> **KQ1E7b, KQ2E7b, KQ3E7b**
16. Baschat, AA, Dewberry, D, et al. Maternal blood-pressure trends throughout pregnancy and development of pre-eclampsia in women receiving first-trimester aspirin prophylaxis. *Ultrasound Obstet Gynecol*. 52(6): 728-733. 2018. PMID: 29266502. <https://dx.doi.org/10.1002/uog.18992> **KQ1E2a, KQ2E2a, KQ3E2a**
17. Becker, R, Keller, T, et al. Individual risk assessment of adverse pregnancy outcome by multivariate regression analysis may serve as basis for drug intervention studies: retrospective analysis of 426 high-risk patients including ethical aspects. *Arch Gynecol Obstet*. 288(1): 41-8. 2013. PMID: 23389246. <https://dx.doi.org/10.1007/s00404-013-2723-1> **KQ1E2, KQ2E2, KQ3E2**
18. Block-Abraham, DM, Turan, OM, et al. First-trimester risk factors for preeclampsia development in women initiating aspirin by 16 weeks of gestation. *Obstet Gynecol*. 123(3): 611-617. 2014. PMID: 24513777. **KQ1E2a, KQ2E2a, KQ3E2a**
19. Cantu, J, Biggio, J, et al. Timing of initiation of low-dose aspirin therapy and preeclampsia prevention in high-risk women. *Am J Obstet Gynecol*. 208(1): S266. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.793> **KQ1E7b, KQ2E7b, KQ3E7b**
20. Carbillon, L. Prevention of early-onset pre-eclampsia. *The Lancet*. 389(10078): 1514-1515. 2017. [https://dx.doi.org/10.1016/S0140-6736\(17\)30950-9](https://dx.doi.org/10.1016/S0140-6736(17)30950-9) **KQ1E1a, KQ2E1a, KQ3E1a**
21. Carpentier, C, Bujold, E, et al. P08.03: Low-dose aspirin for prevention of fetal growth restriction and pre-eclampsia in twins: the GAP pilot randomised. *Ultrasound in Obstetrics & Gynecology*. 50. 178-178. 2017. <https://dx.doi.org/10.1002/UOG.18072> **KQ1E7b, KQ2E7b, KQ3E7b**
22. Carson, MP, Chen, KK. Hypertension in a woman planning pregnancy. *CMAJ*. 186(2): 129-130. 2014. <https://dx.doi.org/10.1503/cmaj.130433> **KQ1E1a, KQ2E1a, KQ3E1a**
23. Chen, Y, Zhao, D, et al. Association of intrauterine exposure to aspirin and blood pressure at 7 years of age: a secondary analysis. *BJOG: An International Journal of Obstetrics & Gynaecology*. 126(5): 599-607. 2019. PMID: 30471180. <https://dx.doi.org/10.1111/1471-0528.15561> **KQ1E2, KQ2E2, KQ3E2**
24. Chu, S, Huang, L, et al. In utero exposure to aspirin and risk of asthma in childhood. *Epidemiology*. 27(5): 726-731. 2016. <https://dx.doi.org/10.1097/EDE.0000000000000519> **KQ1E2a, KQ2E2a, KQ3E2a**
25. Connealy, B, Carreno, C, et al. A history of prior preeclampsia is a major risk factor for preterm birth. *Am J Obstet Gynecol*. 208(1): S264. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.785> **KQ1E2, KQ2E2, KQ3E2**
26. Connell, MT, Sjaarda, LA, et al. The Effects of Aspirin in Gestation and Reproduction (EAGeR) Trial: A Story of Discovery. *Semin Reprod Med*. 35(4): 344-352. 2017. PMID: 29036741. <https://dx.doi.org/10.1055/s-0037-1606384> **KQ1E1a, KQ2E1a, KQ3E1a**
27. Day, RO, Graham, GG. Therapeutics: Non-steroidal anti-inflammatory drugs (NSAIDs). *BMJ (Online)*. 347(7917).

Appendix D. Excluded Studies

2013.
<https://dx.doi.org/10.1136/bmj.f3195>
KQ1E1a, KQ2E1a, KQ3E1a
28. de Vries, JI, van Pampus, MG, et al. Low-molecular-weight heparin added to aspirin in the prevention of recurrent early-onset pre-eclampsia in women with inheritable thrombophilia: the FRUIT-RCT. *Journal of Thrombosis & Haemostasis*. 10(1): 64-72. 2012.
KQ1E6a, KQ2E6a, KQ3E6a
29. Demers, S, Bujold, E, et al. Enoxaparin and Aspirin Compared With Aspirin Alone to Prevent Placenta-Mediated Pregnancy Complications: A Randomized Controlled Trial [Comment on Haddad "Enoxaparin and Aspirin Compared With Aspirin Alone to Prevent Placenta-Mediated Pregnancy Complications: A Randomized Controlled Trial"]. *Obstet Gynecol*. 129(2): 387-388. 2017. PMID: 28121820.
<https://dx.doi.org/10.1097/aog.0000000000001882> **KQ1E1a, KQ2E1a, KQ3E1a**
30. Demers, S, Roberge, S, et al. Low-dose aspirin for the prevention of adverse pregnancy outcomes in women with elevated alpha-fetoprotein. *Journal of Maternal-Fetal & Neonatal Medicine*. 28(6): 726. 2015. PMID: 24893560.
<https://dx.doi.org/10.3109/14767058.2014.930428> **KQ1E1a, KQ2E1a, KQ3E1a**
31. Demers, S, Roberge, S, et al. The use of aspirin during pregnancy. *American Journal of Obstetrics & Gynecology*. 208(2):161-2, 2013 Feb. (2): 161-2, 2013. 2013. **KQ1E1a, KQ2E1a, KQ3E1a**
32. Di Renzo, GC, Gratacos, E, et al. Good clinical practice advice: First trimester screening and prevention of pre-eclampsia in singleton pregnancy. *International Journal of Gynecology and Obstetrics*. 144(3): 325-329. 2019.
<https://dx.doi.org/10.1002/ijgo.12741>
KQ1E1a, KQ2E1a, KQ3E1a
33. Dixon, CL, Marrs, C, et al. Effect of Low-Dose Aspirin on the Time of Onset of Preeclampsia and Time of Delivery. *Am J Perinatol*. 34(12): 1219-1226. 2017. PMID: 28445915.
<https://dx.doi.org/10.1055/s-0037-1602421> **KQ1E7a, KQ2E7a, KQ3E7a**
34. El-Achi, V, Park, F, et al. Does low dose aspirin prescribed for risk of early onset preeclampsia reduce the prevalence of preterm prelabor rupture of membranes?. *J Matern Fetal Neonatal Med*. 1-6. 2019. PMID: 31018725.
<https://dx.doi.org/10.1080/14767058.2019.1611768> **KQ1E2, KQ2E2, KQ3E8**
35. Erratum: Aspirin for Evidence-Based Preeclampsia Prevention trial: effect of aspirin in prevention of preterm preeclampsia in subgroups of women according to their characteristics and medical and obstetrical history (*American Journal of Obstetrics and Gynecology* (2017) 217(5) (585.e1–585.e5) (S0002937817309298) (10.1016/j.ajog.2017.07.038)). *Am J Obstet Gynecol*. 218(4): 454. 2018.
<https://dx.doi.org/10.1016/j.ajog.2018.01.002> **KQ1E1a, KQ2E1a, KQ3E1a**
36. Euser, AG, Metz, TD, et al. Low-dose aspirin for pre-eclampsia prevention in twins with elevated human chorionic gonadotropin. *J Perinatol*. 36(8): 601-5. 2016.
<https://dx.doi.org/10.1038/jp.2016.55>
KQ1E7a, KQ2E7a, KQ3E7a
37. Euser, AG, Metz, TD, et al. Low-dose aspirin reduces the incidence of preeclampsia in women with multiple gestations and elevated human chorionic gonadotropin. *Reproductive Sciences*. 22. 70A-71A. 2015. **KQ1E2, KQ2E2, KQ3E2**
38. Ferrier, C, North, R, et al. Low dose aspirin delays the onset of pre-eclampsia in pregnancies with abnormal uteroplacental circulation. *Proceedings of the 10th World Congress of the ISSHP*. 151. 1996. **KQ1E7b, KQ2E7b, KQ3E7b**

Appendix D. Excluded Studies

39. Finneran, MM, Gonzalez-Brown, VM, et al. Obesity and laboratory aspirin resistance in high-risk pregnant women treated with low-dose aspirin. *Am J Obstet Gynecol.* 220(4): 385.e1-385.e6. 2019. PMID: 30786253. <https://dx.doi.org/10.1016/j.ajog.2019.01.222> **KQ1E7a, KQ2E7a, KQ3E7a**
40. Fratto, VM, Ananth, CV, et al. Late preterm neonatal morbidity in hypertensive versus normotensive women. *Hypertension in Pregnancy.* 35(2): 242-9. 2016. PMID: 26930156. <https://dx.doi.org/10.3109/10641955.2016.1139720> **KQ1E2, KQ2E2, KQ3E2**
41. Gee, Se, Ma'ayeh, M, et al. 1156: glycemic control and aspirin resistance in patients taking low-dose aspirin for preeclampsia prevention. *Am J Obstet Gynecol.* 222(1): S712-. 2020. **KQ1E7, KQ2E7, KQ3E7**
42. Gray, PH. In infants born extremely preterm, aspirin or NSAID use during pregnancy are associated with increased risk of quadriparetic cerebral palsy. *Evid Based Nurs.* 17(1): 16-7. 2014. PMID: 23635955. <https://dx.doi.org/10.1136/eb-2012-101101> **KQ1E1a, KQ2E1a, KQ3E1a**
43. Greene, MF, Solomon, CG. Aspirin to prevent preeclampsia. *N Engl J Med.* 377(7): 690-691. 2017. <https://dx.doi.org/10.1056/NEJMe1708920> **KQ1E1a, KQ2E1a, KQ3E1a**
44. Greer, IA. Antithrombotic therapy for pregnancy complications: let's not throw the baby out with the bath water. *Evidence Based Medicine.* 20(1): 10. 2015. PMID: 25225334. <https://dx.doi.org/10.1136/ebmed-2014-110078> **KQ1E1a, KQ2E1a, KQ3E1a**
45. Gyamfi-Bannerman, C, Fratto, V, et al. Late preterm neonatal morbidity in high-risk hypertensive versus normotensive women. *Am J Obstet Gynecol.* 208(1): S207. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.645> **KQ1E7b, KQ2E7b, KQ3E7b**
46. Gyamfi-Bannerman, C, Fratto, V, et al. Risk of neonatal morbidity at late preterm gestations in relation to low-risk hypertensive versus normotensive women. *Am J Obstet Gynecol.* 208(1): S206-S207. 2013. <https://dx.doi.org/10.1016/j.ajog.2012.10.644> **KQ1E4a, KQ2E4a, KQ3E7a**
47. Haddad, B, Winer, N, et al. Enoxaparin and Aspirin Compared With Aspirin Alone to Prevent Placenta-Mediated Pregnancy Complications: A Randomized Controlled Trial. *Obstet Gynecol.* 128(5): 1053-1063. 2016. PMID: 27741174. **KQ1E6a, KQ2E6a, KQ3E6a**
48. Handal-Orefice, R, Andrikopoulou, M, et al. Aspirin: A potential for reducing intrauterine growth restriction in low-risk women. *Reproductive Sciences.* 25(1): 100A. 2018. **KQ1E4a, KQ2E4a, KQ3E7a**
49. Handal-Orefice, R, Turitz, A, et al. 818: risk-reducing potential of aspirin among low-risk nulliparous, obese women. *Am J Obstet Gynecol.* 222(1): S515-. 2020. **KQ1E7a, KQ2E7a, KQ3E7a**
50. Harrington, K, Kurdi, W, et al. A prospective management study of slow-release aspirin in the palliation of uteroplacental insufficiency predicted by uterine artery Doppler at 20 weeks. *Ultrasound Obstet Gynecol.* 15(1): 13-8. 2000. **KQ1E5, KQ2E5, KQ3E5**
51. Hastie, R, Hesselman, S, et al. Prophylactic aspirin use during pregnancy is associated antenatal complications; a Swedish population register-based cohort study. *Pregnancy Hypertens.* 17. S16. 2019. **KQ1E2, KQ2E2, KQ3E2a**
52. Hauspurg, A, Sutton, EF, et al. Aspirin Effect on Adverse Pregnancy Outcomes Associated With Stage 1 Hypertension in a High-Risk Cohort. *Hypertension.* 72(1): 202-207. 2018. PMID: 29802215. <https://dx.doi.org/10.1161/hypertensiona>

Appendix D. Excluded Studies

- ha.118.11196 **KQ1E7a, KQ2E7a, KQ3E7a**
53. Helou, A, Walker, S, et al. Management of pregnancies complicated by hypertensive disorders of pregnancy: Could we do better?. *Aust N Z J Obstet Gynaecol.* 57(3): 253-259. 2017. PMID: 27396975. <https://dx.doi.org/10.1111/ajo.12499> **KQ1E2, KQ2E2, KQ3E2**
54. Hernandez, RK, Werler, MM, et al. Nonsteroidal antiinflammatory drug use among women and the risk of birth defects. *Am J Obstet Gynecol.* 206(3): 228.e1-228.e8. 2012. <https://dx.doi.org/10.1016/j.ajog.2011.11.019> **KQ1E2a, KQ2E2a, KQ3E2a**
55. Hofmeyr, GJ, Magee, LA. Aspirin and pre-eclampsia: the heart of the matter?. *Bjog.* 2020. **KQ1E1a, KQ2E1a, KQ3E1a**
56. Jensen, MS, Rebordosa, C, et al. Maternal use of acetaminophen, ibuprofen, and acetylsalicylic acid during pregnancy and risk of cryptorchidism. *Epidemiology.* 21(6): 779-785. 2010. **KQ1E2, KQ2E2, KQ3E2**
57. Jessel, R, Allshouse, A, et al. Does low-dose aspirin prevent preterm birth?. *Am J Obstet Gynecol.* 212(1): S342. 2015. <https://dx.doi.org/10.1016/j.ajog.2014.10.907> **KQ1E7b, KQ2E7b, KQ3E7b**
58. Jiang, N, Liu, Q, et al. Effect of calcium channel blockers plus low-dosage aspirin on hypertensive pregnancy outcomes. *Obstet Gynecol.* 123. 57S. 2014. <https://dx.doi.org/10.1097/01.AOG.0000447355.99952.06> **KQ1E7b, KQ2E7b, KQ3E7b**
59. Kalafat, E, Shirazi, A, et al. The role of aspirin in prevention of preeclampsia in twin pregnancies: does the dose matter?. *Am J Obstet Gynecol.* 2020. **KQ1E2, KQ2E2, KQ3E7**
60. Kane, SC, da Silva Costa, F, et al. Recent developments in early pregnancy screening: Are we getting closer to the Holy Grail?. *Med J Aust.* 200(3): 140-141. 2014. <https://dx.doi.org/10.5694/mja13.10955> **KQ1E1a, KQ2E1a, KQ3E1a**
61. Katsi, V, Kanellopoulou, T, et al. Aspirin vs Heparin for the Prevention of Preeclampsia. *Curr Hypertens Rep.* 18(7): 57. 2016. PMID: 27251704. <https://dx.doi.org/10.1007/s11906-016-0664-3> **KQ1E2, KQ2E2, KQ3E2**
62. Keim, SA, Klebanoff, MA. Aspirin use and miscarriage risk. *Epidemiology.* 17(4): 435-439. 2006. **KQ1E2, KQ2E2, KQ3E2**
63. Khazardoost, S, Mousavi, S, et al. Effect of aspirin in prevention of adverse pregnancy outcome in women with elevated alpha-fetoprotein. *Journal of Maternal-Fetal & Neonatal Medicine.* 27(6): 561-5. 2014. PMID: 23901798. <https://dx.doi.org/10.3109/14767058.2013.822483> **KQ1E3a, KQ2E3a, KQ3E3a**
64. Kumar, N, Das, V, et al. Pilot Interventional Study Comparing Fetomaternal Outcomes of 150 mg Versus 75 mg Aspirin Starting Between 11 and 14 Weeks of Pregnancy in Patients with High Risk of Preeclampsia: A Randomized Control Trial. *J Obstet Gynaecol India.* 70(1): 23-29. 2020. **KQ1E3a, KQ2E3a, KQ3E3a**
65. Lan, PG, Gillin, AG, et al. Effect of early use of low-dose aspirin therapy on late-onset preeclampsia. *Journal of Maternal-Fetal & Neonatal Medicine.* 32(13): 2137-2142. 2019. PMID: 29366353. <https://dx.doi.org/10.1080/14767058.2018.1427718> **KQ1E2, KQ2E2, KQ3E6**
66. Leeson, P. Updated review identifies no adverse impact on mother or offspring during the perinatal period of aspirin use for prevention of preeclampsia. *Evidence Based Medicine.* 20(1): 11. 2015. PMID: 25238770. <https://dx.doi.org/10.1136/ebmed-2014-110056> **KQ1E1a, KQ2E1a, KQ3E1a**

Appendix D. Excluded Studies

67. Leon, M, Moussa, H, et al. Rate of GDM and pregnancy outcomes in chronic hypertensive pregnancies with and without gestational diabetes mellitus. *Am J Obstet Gynecol.* 212(1): S403-S404. 2015. <https://dx.doi.org/10.1016/j.ajog.2014.10.1048> **KQ1E7b, KQ2E7b, KQ3E7b**
68. Leon, MG, Moussa, HN, et al. Rate of Gestational Diabetes Mellitus and Pregnancy Outcomes in Patients with Chronic Hypertension. *Am J Perinatol.* 33(8): 745-50. 2016. PMID: 26890438. <https://dx.doi.org/10.1055/s-0036-1571318> **KQ1E2, KQ2E2, KQ3E2**
69. Li, YY, Gu, QR, et al. Preventing preterm preeclampsia. *Am J Obstet Gynecol.* 219(6): 633-634. 2018. <https://dx.doi.org/10.1016/j.ajog.2018.08.027> **KQ1E1a, KQ2E1a, KQ3E1a**
70. Lind, DV, Main, KM, et al. Maternal use of mild analgesics during pregnancy associated with reduced anogenital distance in sons: a cohort study of 1027 mother-child pairs. *Hum Reprod.* 32(1): 223-231. 2017. PMID: 27852690. **KQ1E2a, KQ2E2a, KQ3E2a**
71. Liu, F, Yang, H, et al. Effect of a small dose of aspirin on quantitative test of 24-h urinary protein in patients with hypertension in pregnancy. *Exp Ther Med.* 13(1): 37-40. 2017. PMID: 28123464. <https://dx.doi.org/10.3892/etm.2016.3924> **KQ1E3a, KQ2E3a, KQ3E3a**
72. Liu, FM, Zhao, M, et al. Effect of regular oral intake of aspirin during pregnancy on pregnancy outcome of high-risk pregnancy-induced hypertension syndrome patients. *Eur Rev Med Pharmacol Sci.* 20(23): 5013-5016. 2016. PMID: 27981529. **KQ1E3a, KQ2E3a, KQ3E3a**
73. Low-dose aspirin for prevention of preeclampsia. *Med Lett Drugs Ther.* 56(1445): 49. 2014. **KQ1E1a, KQ2E1a, KQ3E1a**
74. Mason, R, Jones, HA. An audit of antenatal management and outcomes of pregnancies with low first-trimester serum pregnancy-associated plasma protein-A (PAPP-A). *BJOG: An International Journal of Obstetrics and Gynaecology.* 125. 68. 2018. <https://dx.doi.org/10.1111/1471-0528.15493> **KQ1E1, KQ2E1, KQ3E1**
75. Meher, S, Alfirevic, Z. Aspirin for pre-eclampsia: beware of subgroup meta-analysis. *Ultrasound Obstet Gynecol.* 41(5): 479-85. 2013. PMID: 23610032. <https://dx.doi.org/10.1002/uog.12470> **KQ1E1a, KQ2E1a, KQ3E1a**
76. Meher, S, Duley, L, et al. Antiplatelet therapy before or after 16 weeks' gestation for preventing preeclampsia: an individual participant data meta-analysis. *Am J Obstet Gynecol.* 216(2): 121-128.e2. 2017. <https://dx.doi.org/10.1016/j.ajog.2016.10.016> **KQ1E5, KQ2E5, KQ3E5**
77. Meiri, H, Sammar, M, et al. Prediction of preeclampsia by placental protein 13 and background risk factors and its prevention by aspirin. *J Perinat Med.* 42(5): 591-601. 2014. PMID: 24607918. <https://dx.doi.org/10.1515/jpm-2013-0298> **KQ1E2, KQ2E2, KQ3E2**
78. Mendel, A, Bernatsky, SB, et al. Low aspirin use and high prevalence of preeclampsia risk factors among pregnant women in a multinational SLE inception cohort. *Ann Rheum Dis.* 78(7): 1010-1012. 2019. **KQ1E2, KQ2E2, KQ3E7**
79. Mogra, R. Screening and prevention of pre-term pre-eclampsia – A prime time to act. *Australasian Journal of Ultrasound in Medicine.* 21(4): 187-190. 2018. <https://dx.doi.org/10.1002/ajum.12101> **KQ1E1a, KQ2E1a, KQ3E1a**
80. Mol, BWJ, Van Vliet, EOG, et al. Antiplatelet agents and the prevention of spontaneous preterm birth: A meta-analysis of individual participant data. *J Paediatr Child Health.* 51. 53. 2015.

Appendix D. Excluded Studies

- <https://dx.doi.org/10.1111/jpc.12884-4>
KQ1E7b, KQ2E7b, KQ3E7b
81. Mone, F, Mulcahy, C, et al. A randomised controlled trial and cost effectiveness analysis of low dose aspirin with an early screening test for pre-eclampsia in low risk women. *BJOG: An International Journal of Obstetrics and Gynaecology*. 124. 10-11. 2017.
<https://dx.doi.org/10.1111/1471-0528.14585> **KQ1E7a, KQ2E7a, KQ3E7a**
82. Mone, F, Mulcahy, C, et al. A randomised controlled trial and cost-effectiveness analysis of low dose aspirin with an early screening test for preeclampsia in low risk women. *Reproductive Sciences*. 24(1): 68A. 2017.
<https://dx.doi.org/10.1177/1933719117699773> **KQ1E7a, KQ2E7a, KQ3E7a**
83. Mone, F, Mulcahy, C, et al. Performance of the fetal medicine foundation preeclampsia screening test in nulliparous women: results of the TEST multicenter RCT. *BJOG: an international journal of obstetrics and gynaecology*. Conference: 19th annual conference of the british maternal and fetal medicine society, BMFMS 2017. Netherlands. 124. 10. 2017.
<https://dx.doi.org/10.1111/1471-0528.14585> **KQ1E7b, KQ2E7b, KQ3E7b**
84. Mone, F, Mulcahy, C, et al. The impact of low dose aspirin on markers of placental disease-results of the test multicenter RCT. *BJOG: An International Journal of Obstetrics and Gynaecology*. 124. 11. 2017.
<https://dx.doi.org/10.1111/1471-0528.14585> **KQ1E7b, KQ2E7b, KQ3E7b**
85. Mone, F, Mulcahy, C, et al. The impact of low dose aspirin on preeclampsia biomarkers and fetal growth in lowrisk women. *Am J Obstet Gynecol*. 218(1): S303. 2018. **KQ1E7a, KQ2E7a, KQ3E7a**
86. Moore, G, Allshouse, A, et al. Early aspirin decreases late-onset preeclampsia (≥ 34 weeks), particularly in women with chronic hypertension. *Am J Obstet Gynecol*. 210(1): S144-S145. 2014.
<https://dx.doi.org/10.1016/j.ajog.2013.10.305> **KQ1E2, KQ2E2, KQ3E2**
87. Moore, GS, Allshouse, AA, et al. Baseline placental growth factor levels for the prediction of benefit from early aspirin prophylaxis for preeclampsia prevention. *Pregnancy Hypertens*. 5(4): 280-6. 2015. PMID: 26597741.
<https://dx.doi.org/10.1016/j.pregphy.2015.06.001> **KQ1E2, KQ2E2, KQ3E2**
88. Moore, GS, Allshouse, AA, et al. Early initiation of low-dose aspirin for reduction in preeclampsia risk in high-risk women: a secondary analysis of the MFMU High-Risk Aspirin Study. *J Perinatol*. 35(5): 328-31. 2015.
<https://dx.doi.org/10.1038/jp.2014.214> **KQ1E7a, KQ2E7a, KQ3E7a**
89. Mosha, D, Mazuguni, F, et al. Medication exposure during pregnancy: A pilot pharmacovigilance system using health and demographic surveillance platform. *BMC Pregnancy Childbirth*. 14(1). 2014.
<https://dx.doi.org/10.1186/1471-2393-14-322> **KQ1E3a, KQ2E3a, KQ3E3a**
90. Mousavi, S. Effect of aspirin in prevention of adverse pregnancy outcome in women with elevated alpha-fetoprotein. *J Perinat Med*. 43. 2015.
<https://dx.doi.org/10.1515/jpm-2015-2003> **KQ1E3a, KQ2E3a, KQ3E3a**
91. Moussa, HN, Hosseini Nasab, S, et al. Adverse pregnancy outcomes in deliveries prior to, at and beyond 39 weeks; low- and high-risk women. *Journal of Maternal-Fetal & Neonatal Medicine*. 31(19): 2545-2549. 2018. PMID: 28651448.

Appendix D. Excluded Studies

- <https://dx.doi.org/10.1080/14767058.2017.1347624> **KQ1E7a, KQ2E7a, KQ3E7a**
92. Moussa, HN, Leon, MG, et al. Pregnancy outcomes between superimposed preeclampsia with and without severe features-evaluation of the new ACOG task force guidelines. *Reproductive Sciences*. 22. 179A. 2015. **KQ1E7b, KQ2E7b, KQ3E7b**
93. Moussa, HN, Leon, MG, et al. Pregnancy Outcomes in Women with Preeclampsia Superimposed on Chronic Hypertension with and without Severe Features. *Am J Perinatol*. 34(4): 403-408. 2017. PMID: 27606778. <https://dx.doi.org/10.1055/s-0036-1592134> **KQ1E7, KQ2E7, KQ3E7**
94. Moussa, HN, Wu, ZH, et al. Customized versus Population Fetal Growth Norms and Adverse Outcomes Associated with Small for Gestational Age Infants in a High-Risk Cohort. *Am J Perinatol*. 32(7): 621-6. 2015. PMID: 25262454. <https://dx.doi.org/10.1055/s-0034-1390343> **KQ1E7a, KQ2E7a, KQ3E7a**
95. Mulcahy, C, Mone, F, et al. Performance of routine first trimester sonographic evaluation of fetal anatomy: Results of the multicentre test RCT. *Am J Obstet Gynecol*. 216(1): S83-S84. 2017. **KQ1E7a, KQ2E7a, KQ3E7a**
96. Mulcahy, C, Mone, F, et al. The Impact of Aspirin on Ultrasound Markers of Uteroplacental Flow in Low-Risk Pregnancy: Secondary Analysis of a Multicenter RCT. *Am J Perinatol*. 36(8): 855-863. 2019. PMID: 30396226. <https://dx.doi.org/10.1055/s-0038-1675208> **KQ1E7, KQ2E7, KQ3E7**
97. Murtoniemi, K, Vahlberg, T, et al. The effect of low-dose aspirin on serum placental growth factor levels in a high-risk PREDO cohort. *Pregnancy Hypertens*. 13. 51-57. 2018. PMID: 30177071. <https://dx.doi.org/10.1016/j.preghy.2018.04.003> **KQ1E7, KQ2E7, KQ3E7**
98. Navaratnam, K, Agarwal, U, et al. Aspirin nonresponsiveness in pregnant women at high risk of pre-eclampsia. *BJOG: An International Journal of Obstetrics and Gynaecology*. 126. 4. 2019. <https://dx.doi.org/10.1111/1471-0528.15629> **KQ1E1, KQ2E1, KQ3E1**
99. Newnham, JP, Godfrey, M, et al. Low dose aspirin for the treatment of fetal growth restriction: a randomized controlled trial. *Aust N Z J Obstet Gynaecol*. 35(4): 370-374. 1995. **KQ1E4, KQ2E4, KQ3E4**
100. Nicolaides, KH. Aspirin versus Placebo in Pregnancies at High Risk for Preterm Preeclampsia. *N Engl J Med*. 377(24): 2400. 2017. PMID: 29236633. <https://dx.doi.org/10.1056/NEJMc1713798> **KQ1E1a, KQ2E1a, KQ3E1a**
101. Nobles, CJ, Mendola, P, et al. Preconception and very early pregnancy blood pressure and development of preterm preeclampsia, term preeclampsia and gestational hypertension. *Fertil Steril*. 112(3): e301-e302. 2019. **KQ1E2, KQ2E2, KQ3E2**
102. Odibo, AO, Goetzinger, KR, et al. Early prediction and aspirin for prevention of pre-eclampsia (EPAPP) study: a randomized controlled trial. *Ultrasound Obstet Gynecol*. 46(4): 414-8. 2015. <https://dx.doi.org/10.1002/uog.14889> **KQ1E8, KQ2E8, KQ3E8**
103. Ohno, M, Girsan, A, et al. Does low-dose aspirin for preeclampsia prevention increase the risk of antepartum bleeding or placental abruption?. *Am J Obstet Gynecol*. 210(1): S189. 2014. <https://dx.doi.org/10.1016/j.ajog.2013.10.406> **KQ1E7a, KQ2E7a, KQ3E7a**
104. Otero Gonzalez, A, Uribe Moya, S, et al. Hyperbaric index in the primary prevention of hypertensive complications in high-risk pregnancy. *Nefrologia*. 35(6): 572-7. 2015. PMID: 26547790.

Appendix D. Excluded Studies

- <https://dx.doi.org/10.1016/j.nefro.2015.10.001> **KQ1E2a, KQ2E2a, KQ3E2a**
105. Papatheodorou, SI. Re: Early prediction and aspirin for prevention of pre-eclampsia (EPAPP) study: a randomized controlled trial. A. O. Odibo, K. R. Goetzinger, L. Odibo and M. G. Tuuli. *Ultrasound Obstet Gynecol* 2015; 46: 414-418. *Ultrasound Obstet Gynecol*. 46(4): 389. 2015. PMID: 26428714. <https://dx.doi.org/10.1002/uog.15736> **KQ1E1a, KQ2E1a, KQ3E1a**
106. Parchem, JG, Sibai, BM. An aspirin a day in pregnancy keeps childhood hypertension away? *BJOG: An International Journal of Obstetrics and Gynaecology*. 126(5): 608. 2019. <https://dx.doi.org/10.1111/1471-0528.15598> **KQ1E1a, KQ2E1a, KQ3E1a"**
107. Parikh, NI, Gonzalez, J. Preeclampsia and hypertension: Courting a long while: Time to make it official. *JAMA Intern Med*. 177(7): 917-918. 2017. <https://dx.doi.org/10.1001/jamainternmed.2017.1422> **KQ1E1a, KQ2E1a, KQ3E1a**
108. Park, F, Russo, K, et al. Prediction and Prevention of Early-Onset Preeclampsia: Impact of Aspirin after First-Trimester Screening. *Obstet Gynecol Surv*. 71(1): 11-13. 2016. <https://dx.doi.org/10.1097/01.ogx.0000476311.74309.ae> **KQ1E1a, KQ2E1a, KQ3E1a**
109. Park, F, Russo, K, et al. The impact of aspirin on the prevalence of early onset pre-eclampsia after first trimester screening. *Prenat Diagn*. 34. e4. 2014. <https://dx.doi.org/10.1111/pd.4477> **KQ1E2, KQ2E2, KQ3E2**
110. Pasquier, E, De Saint Martin, L. To the editor. *Ann Intern Med*. 163(6): 484-485. 2015. <https://dx.doi.org/10.7326/L1551372> **KQ1E1a, KQ2E1a, KQ3E1a**
111. Petersen, TG, Liew, Z, et al. Use of paracetamol, ibuprofen or aspirin in pregnancy and risk of cerebral palsy in the child. *Int J Epidemiol*. 47(1): 121-130. 2018. PMID: 29149272. <https://dx.doi.org/10.1093/ije/dyx235> **KQ1E2a, KQ2E2a, KQ3E2a**
112. Poon, LC, Rolnik, DL, et al. ASPRE Trial: Incidence of Preterm Preeclampsia in Patients Fulfilling ACOG and NICE Criteria According to Risk by FMF Algorithm. *Obstet Gynecol Surv*. 73(11): 623-625. 2018. <https://dx.doi.org/10.1097/01.ogx.0000547739.61320.71> **KQ1E7a, KQ2E7a, KQ3E7a**
113. Preconception blood pressure linked to pregnancy loss risk. *Clinical Pharmacist*. 10(7). 2018. <https://dx.doi.org/10.1211/CP.2018.20204909> **KQ1E1a, KQ2E1a, KQ3E1a**
114. Prefumo, F, Frusca, T, et al. Acetylsalicylic acid in pregnant women with chronic hypertension. *Am J Obstet Gynecol*. 218(4): 463-464. 2018. <https://dx.doi.org/10.1016/j.ajog.2018.01.020> **KQ1E1a, KQ2E1a, KQ3E1a**
115. Premru-Srsen, T, Kocic, Z, et al. Effect of current guidelines on prevention of pre-eclampsia with low-dose aspirin in primary settings: A population-based case-control study. *Int J Gynaecol Obstet*. 2020. **KQ1E2a, KQ2E2a, KQ3E2a**
116. Quinlivan, JA. Simple and safe: preventing preterm birth with aspirin. *Lancet*. 395(10220): 250-252. 2020. **KQ1E1a, KQ2E1a, KQ3E1a**
117. Rezk, M, Ellakwa, H, et al. Maternal and fetal morbidity following discontinuation of antihypertensive drugs in mild to moderate chronic hypertension: A 4-year observational study. *Pregnancy Hypertens*. 6(4): 291-294. 2016. PMID: 27939471. <https://dx.doi.org/10.1016/j.preghy.2016.05.002> **KQ1E3a, KQ2E3a, KQ3E3a**

Appendix D. Excluded Studies

118. Roberge, S, Demers, S, et al. Initiation of aspirin in early gestation for the prevention of pre-eclampsia. *BJOG: An International Journal of Obstetrics and Gynaecology*. 120(6): 773-774. 2013. <https://dx.doi.org/10.1111/1471-0528.12170> **KQ1E1a, KQ2E1a, KQ3E1a**
119. Roberge, S, Demers, S, et al. Low-dose aspirin for prevention of morbidity and mortality from preeclampsia. *Ann Intern Med*. 161(8): 613. 2014. <https://dx.doi.org/10.7326/L14-5020-4> **KQ1E1a, KQ2E1a, KQ3E1a**
120. Roberge, S, Giguère, Y, et al. Erratum: Early administration of low-dose aspirin for the prevention of severe and mild preeclampsia: A systematic review and meta-analysis (*American Journal of Perinatology* (2014) 31 (e3)). *Am J Perinatol*. 31(6): e3. 2014. <https://dx.doi.org/10.1055/s-0033-1343351> **KQ1E1a, KQ2E1a, KQ3E1a**
121. Roberge, S, Odibo, AO, et al. Aspirin for the Prevention of Preeclampsia and Intrauterine Growth Restriction. *Clin Lab Med*. 36(2): 319-29. 2016. <https://dx.doi.org/10.1016/j.cl.2016.01.013> **KQ1E2, KQ2E2, KQ3E2**
122. Roberts, JM, Himes, KP. Pre-eclampsia: Screening and aspirin therapy for prevention of pre-eclampsia. *Nature Reviews Nephrology*. 13(10): 602-604. 2017. <https://dx.doi.org/10.1038/nrneph.2017.121> **KQ1E1a, KQ2E1a, KQ3E1a**
123. Rodger, MA. Prevention of early-onset pre-eclampsia – Authors' reply. *The Lancet*. 389(10078): 1515. 2017. [https://dx.doi.org/10.1016/S0140-6736\(17\)30951-0](https://dx.doi.org/10.1016/S0140-6736(17)30951-0) **KQ1E1a, KQ2E1a, KQ3E1a**
124. Schiff, E, Barkai, G, et al. Low-dose aspirin does not influence the clinical course of women with mild pregnancy-induced hypertension. *Obstet Gynecol*. 76(5 Pt 1): 742-4. 1990. **KQ1E4, KQ2E4, KQ3E4**
125. Seidler, AL, Askie, L, et al. Optimal aspirin dosing for preeclampsia prevention. *Am J Obstet Gynecol*. 219(1): 117-118. 2018. PMID: 29588190. <https://dx.doi.org/10.1016/j.ajog.2018.03.018> **KQ1E5, KQ2E5, KQ3E5**
126. Sentilhes, L, Azria, E, et al. Aspirin versus placebo in pregnancies at high risk for preterm preeclampsia. *N Engl J Med*. 377(24): 2399-2400. 2017. <https://dx.doi.org/10.1056/NEJMc1713798> **KQ1E1a, KQ2E1a, KQ3E1a**
127. Sharma, AK, Bhatla, N. Aspirin for pregnancies at high risk for preterm pre-eclampsia. *Natl Med J India*. 31(1): 26-27. 2018. **KQ1E1a, KQ2E1a, KQ3E1a**
128. Sharma, N, Srinivasan, S, et al. Role of Aspirin in High Pulsatility Index of Uterine Artery: A Consort Study. *J Obstet Gynaecol India*. 68(5): 382-388. 2018. PMID: 30224843. <https://dx.doi.org/10.1007/s13224-017-1058-4> **KQ1E3a, KQ2E3a, KQ3E3a**
129. Sibai, BM. Therapy: Low-dose aspirin to reduce the risk of pre-eclampsia?. *Nature Reviews Endocrinology*. 11(1): 6-8. 2015. <https://dx.doi.org/10.1038/nrendo.2014.199> **KQ1E1a, KQ2E1a, KQ3E1a**
130. Sisti, G. Maternal Outcomes Associated with Lower Range Stage 1 Hypertension. *Obstet Gynecol*. 133(1): 190. 2019. <https://dx.doi.org/10.1097/AOG.00000000000003045> **KQ1E1a, KQ2E1a, KQ3E1a**
131. Sjaarda, LA, Radin, RG, et al. Preconception Low-Dose Aspirin Restores Diminished Pregnancy and Live Birth Rates in Women With Low-Grade Inflammation: A Secondary Analysis of a Randomized Trial. *J Clin Endocrinol Metab*. 102(5): 1495-1504. 2017. PMID: 28323989. <https://dx.doi.org/10.1210/jc.2016-2917> **KQ1E1, KQ2E1, KQ3E1**

Appendix D. Excluded Studies

132. Stanescu, AD, Banica, R, et al. Low dose aspirin for preventing fetal growth restriction: A randomised trial. *J Perinat Med.* 46(7): 776-779. 2018. <https://dx.doi.org/10.1515/jpm-2017-0184> **KQ1E8, KQ2E8, KQ3E8**
133. Stern, C, Karoline, MP, et al. The effect of low dose aspirin on pregnancy outcomes in women with recurrent preeclampsia. *Reproductive Sciences.* 21(3): 353A. 2014. <https://dx.doi.org/10.1177/1933719114528275> **KQ1E7b, KQ2E7b, KQ3E7b**
134. Stewart, GB, Altman, DG, et al. Statistical analysis of individual participant data meta-analyses: a comparison of methods and recommendations for practice. *PLoS ONE [Electronic Resource].* 7(10): e46042. 2012. **KQ1E5, KQ2E5, KQ3E5**
135. Sutton, EF, Hauspurg, A, et al. Maternal Outcomes Associated With Lower Range Stage 1 Hypertension Maternal Outcomes Associated With Lower Range Stage 1 Hypertension. *Obstet Gynecol.* 132(4): 843-849. 2018. <https://dx.doi.org/10.1097/aog.00000000000002870> **KQ1E7a, KQ2E7a, KQ3E7a**
136. Talari, H, Mesdaghinia, E, et al. Aspirin and preeclampsia prevention in patients with abnormal uterine artery blood flow. *Iran Red Crescent Med J.* 16(8): e17175. 2014. PMID: 25389483. <https://dx.doi.org/10.5812/ircmj.17175> **KQ1E3a, KQ2E3a, KQ3E3a**
137. Theilen, L, Campbell, H, et al. 613: platelet activation and placenta-mediated adverse pregnancy outcomes. *Am J Obstet Gynecol.* 222(1): S392-. 2020. **KQ1E7, KQ2E7, KQ3E7**
138. Thilaganathan, B. The ASPRE pre-eclampsia trial: Implications for basic research and clinical practice. *Cardiovasc Res.* 114(8): e60-e61. 2018. <https://dx.doi.org/10.1093/cvr/cvy107> **KQ1E1a, KQ2E1a, KQ3E1a**
139. Thompson, JM, Waldie, KE, et al. Associations between acetaminophen use during pregnancy and ADHD symptoms measured at ages 7 and 11 years. *PLoS ONE [Electronic Resource].* 9(9): e108210. 2014. PMID: 25251831. <https://dx.doi.org/10.1371/journal.pone.0108210> **KQ1E2a, KQ2E2a, KQ3E2a**
140. Thornton, J. Initiation of aspirin in early gestation for the prevention of pre-eclampsia. *BJOG: An International Journal of Obstetrics and Gynaecology.* 120(6): 774-775. 2013. <https://dx.doi.org/10.1111/1471-0528.12169> **KQ1E1a, KQ2E1a, KQ3E1a**
141. Tong, S, Mol, BW, et al. Preventing preeclampsia with aspirin: does dose or timing matter?. *Am J Obstet Gynecol.* 216(2): 95-97. 2017. <https://dx.doi.org/10.1016/j.ajog.2016.12.003> **KQ1E1a, KQ2E1a, KQ3E1a**
142. Vainio, M, Kujansuu, E, et al. Low dose acetylsalicylic acid in prevention of pregnancy-induced hypertension and intrauterine growth retardation in women with bilateral uterine artery notches. *BJOG.* 109(2): 161-167. 2002. **KQ1E5, KQ2E5, KQ3E5**
143. van Vliet, EO, Askie, LA, et al. Antiplatelet Agents and the Prevention of Spontaneous Preterm Birth: A Systematic Review and Meta-analysis. *Obstet Gynecol.* 129(2): 327-336. 2017. <https://dx.doi.org/10.1097/aog.0000000000001848> **KQ1E5, KQ2E5, KQ3E5**
144. Villa, PM, Kajantie, E, et al. Aspirin in the prevention of pre-eclampsia in high-risk women. *BJOG: An International Journal of Obstetrics and Gynaecology.* 120(6): 773. 2013. <https://dx.doi.org/10.1111/1471-0528.12135> **KQ1E1a, KQ2E1a, KQ3E1a**
145. Voutetakis, A, Pervanidou, P, et al. Aspirin for the Prevention of Preeclampsia and Potential Consequences for Fetal Brain

Appendix D. Excluded Studies

- Development. *JAMA Pediatr.* 2019. PMID: 31135877. <https://dx.doi.org/10.1001/jamapediatrics.2019.1260> **KQ1E1a, KQ2E1a, KQ3E1a**
146. Wheeler, S, Grotegut, C, et al. Aspirin for preeclampsia prevention, latency and chorioamnionitis in preterm premature rupture of membranes. *Am J Obstet Gynecol.* 213(6): 897. 2015. **KQ1E7b, KQ2E7b, KQ3E7b**
147. Wright, D, Nicolaides, KH. Aspirin delays the development of preeclampsia. *Am J Obstet Gynecol.* 2019. PMID: 30797761. <https://dx.doi.org/10.1016/j.ajog.2019.02.034> **KQ1E7a, KQ2E7a, KQ3E7a**
148. Wright, D, Poon, LC, et al. Aspirin for Evidence-Based Preeclampsia Prevention trial: influence of compliance on beneficial effect of aspirin in prevention of preterm preeclampsia. *Am J Obstet Gynecol.* 217(6): 685.e1-685.e5. 2017. PMID: 28888591. <https://dx.doi.org/10.1016/j.ajog.2017.08.110> **KQ1E7a, KQ2E7a, KQ3E7a**
149. Wright, D, Rolnik, DL, et al. Aspirin for Evidence-Based Preeclampsia Prevention trial: effect of aspirin on length of stay in the neonatal intensive care unit. *Am J Obstet Gynecol.* 218(6): 612.e1-612.e6. 2018. PMID: 29505771. <https://dx.doi.org/10.1016/j.ajog.2018.02.014> **KQ1E7a, KQ2E7a, KQ3E7a**
150. Zhao, m Y. Low-dose aspirin prescribed at bed time for the prevention of preeclampsia in high-risk pregnant women. *Shengzhi Yu Biyun.* 32(5): 355-359. 2012. **KQ1E9, KQ2E9, KQ3E9**

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m ²	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
ASPRE, 2017 ⁵⁸ Good	Aged 18 years or more, singleton pregnancy, live fetus at the time that scanning was performed at 11 to 13 weeks' gestation, high risk (>1 in 100) for preterm preeclampsia according to the screening algorithm	Unconscious or severely ill status, learning difficulties or serious mental illness, major fetal abnormality identified at the time that scanning was performed at 11 to 13 weeks of gestation, regular treatment with aspirin within 28 days before screening, bleeding disorder such as von Willebrand's disease, peptic ulceration, hypersensitivity to aspirin, long-term use of nonsteroidal anti-inflammatory medication, and participation in another drug trial within 28 days before screening	31.5*	White: 67.1 AA: 25.2 Asian: 6.4 Other: 1.3	IG: 26.7 CG: 26.5	IG: 68.5 CG: 66.1	IG: 7.1 CG: 7.2	Previous PreE IG: 10.9 CG: 10.2 Chronic HTN IG: 6.1 CG: 7.4 Chronic DM IG: 1.9 CG: 1.2
Ayala, 2012 ⁵⁶ Good	Higher risk for gestational HTN or preeclampsia, receiving medical care and follow-up at the Obstetric Physiopathology service (high-risk unit) of the hospital, ≤16 weeks' gestation, maternal age ≥18 years	Multiple pregnancy, chronic HTN or any other condition requiring the use of BP-lowering medication, cardiovascular disorders (unstable angina pectoris, heart failure, life-threatening arrhythmia, atrial fibrillation, kidney failure, and grade III–IV retinopathy), chronic liver disease, any disease requiring the use of anti-inflammatory medication, DM or any other endocrine disease such	30.7	NR	IG: 25.4 CG: 25.5	IG: 49.4 CG: 55.1	NR	NR

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
		as hyperthyroidism, history of drug/alcohol abuse, night/ shiftwork employment, AIDS, intolerance to ABPM, and inability to communicate and comply with all of the study requirements						
Benigni, 1989 ⁸⁷ Fair	At high risk of pregnancy-induced HTN, with HTN, or previous obstetrical history (fetal death due to placental insufficiency, severe IUGR, early onset PreE [<32 weeks])	Women with antiphospholipid antibodies (lupus-like, anticoagulant, anticardiolipin antibodies)	31.5	NR	NR	NR	IG: 6.3 CG: 0	Previous PreE IG: 50 CG: 35 Chronic HTN IG: 35.2 CG: 31.3
Caspi, 1994 ⁹³ Good	Uncomplicated twin pregnancies	Chronic renal, cardiovascular, pulmonary or hepatic disorders, past or present coagulopathy or peptic ulcer, gestational DM and known hypersensitivity to aspirin	28.3	NR	NR	IG: 41.6 CG: 30.4	NR	Chronic HTN IG: 1.0 CG: 3.0 Multifetal Gestation IG: 100 CG: 100
CLASP, 1994 ⁹⁴ Good	12 to 32 weeks' gestation, at sufficient risk of developing preeclampsia or IUGR for the use of low-dose aspirin to be considered; risk factors for prophylactic aspirin use defined as history of preeclampsia or IUGR in previous pregnancy, chronic	Increased risk of bleeding, asthma, allergy to aspirin, or a high likelihood of immediate delivery	28.5	NR	NR	IG: 27.9 CG: 27.9	IG: 20.7 CG: 20.1	Chronic HTN IG: 19.9 CG: 20.2 Chronic DM IG: 2.7 CG: 3.1

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
	HTN, renal disease, maternal age, family history, or multiple pregnancy; or, for therapeutic use, women with signs or symptoms of preeclampsia or IUGR in current pregnancy							
Davies, 1995 ⁹⁵ Fair	No previous pregnancy beyond 12 weeks' gestation, 12 to 19 weeks' gestation, hemoglobin concentrate > 13.2 g/dL, DBP <90 mmHg, no proteinuria at time of randomization	Multiple pregnancy, DM, recurrent spontaneous abortions, or any contraindication to aspirin therapy	25.0	White: 95.8	NR	IG: 100 CG: 100	IG: 10.3 CG: 13.3	Previous PreE IG: 0 CG: 0 Chronic HTN IG: 0 CG: 0 Chronic DM IG: 0 CG: 0 Multifetal Gestation IG: 0 CG: 0
Gallery, 1997 ⁹⁶ Fair	<16 weeks' gestation, at high risk of developing preeclampsia	Women with a history of aspirin allergy, aspirin-sensitive asthma, or preexisting bleeding diathesis	28.5 (22-38)	White: 95.5	NR	IG: 42.0 CG: 43.0	NR	Previous PreE IG: 17.0 CG: 22.0 Chronic HTN IG: 57.0 CG: 52.0
Grab, 2000 ¹⁰⁰ Fair	Singleton pregnancies, <20 weeks' gestation with early IUGR,	DM, preexisting proteinuric HTN, fetal malformations, or	NR	NR	NR	NR	NR	Previous PreE IG: 40.9

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
	impaired uteroplacental blood, chronic HTN, history of still birth, growth restriction, or preeclampsia	chromosome abnormalities						CG: 42.9 Chronic HTN IG: 50.0 CG: 38.1 Multifetal Gestation IG: 0 CG: 0
Hauth, 1993 ⁹⁰ Good	Nulliparous, age 28 years or younger, no later than 22 weeks of gestation	History of illness or conditions known to increase the incidence of PreE or PIH (i.e. renal disease, collagen vascular disease, DM, multifetal gestation, chronic HTN), failed "run-in" test for compliance	20.4	White: 28.5 AA: 71.5	NR	IG: 100 CG: 100	NR	Chronic HTN IG: 0 CG: 0 Chronic DM IG: 0 CG: 0
Hermida, 1997 ⁹⁷ Good	Absence of any condition requiring the use of antihypertensive medication, maternal age 18 to 40 years, <16 weeks' gestation.	Multiple pregnancy, chronic HTN, chronic liver disease, any disease requiring the use of anti-inflammatory medication, DM, or any other endocrine disease such as hyperthyroidism, intolerance to the use of ambulatory BP monitor, among others	30.2	White: 100	IG: 25.3 CG: 24.9	NR	NR	Multifetal Gestation IG: 0 CG: 0
McParland, 1990 ⁸⁹ Fair	Nulliparous, persistent abnormal Doppler flow-velocity waveforms	Known ASA allergy, maternal DM, bleeding disorders, peptic ulceration, systemic lupus erythematosus	26.1	White: 69.0 AA: 20.0 Asian: 7.0 Other: 4.0	IG: 24.1 CG: 24.6	IG: 100 CG: 100	IG: 16 CG: 23	NR

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
MFMU, 1998 ⁹⁸ Good	One of four high risk groups: DM, chronic HTN, multifetal gestation, or previous preeclampsia	Multifetal gestation in combination with DM, HTN, or proteinuria, history of PreE, current proteinuria	26.5	White: 32.7 AA: 56.3 Hispanic: 10.8 Other: 0.5	NR	NR	IG: 16.8 CG: 16.4	NR
Mone, 2018 ¹⁰⁴ Fair	Nulliparous, aged 18 years or older, 11 to 13 weeks' gestation, viable singleton pregnancy, did not meet criteria for taking aspirin based upon major preeclampsia risk factors (chronic kidney disease, autoimmune disease, DM, and chronic HTN)	Participants already taking part in a clinical trial, coexistence of a fetal congenital anomaly at recruitment, aspirin hypersensitivity	33.5 (19-44)	White: 96.8 AA: 0.8 Asian: 2.4	IG: 25.2 CG: 22.9	IG: 100 CG: 100	IG: 9.2 CG: 5.9	Chronic HTN IG: 0 CG: 0 Chronic DM IG: 0 CG: 0
Morris, 1996 ¹⁰⁶	Nulliparous with abnormal uterine artery waveforms at 18 weeks' gestation	Pregnancies dated at less than 17 weeks' or more than 19 weeks' of gestation	23.8	NR	NR	IG: 100 CG: 100	NR	NR
Rotchell, 1998 ⁹⁹ Good	12 to 32 weeks' gestation without contraindications	Increased risk of bleeding, known allergy to aspirin, high likelihood of immediate delivery or previous placental abruption	NR	NR	NR	IG: 44.0 CG: 44.0	NR	Chronic HTN IG: 0.44 CG: 0.44 Chronic DM IG: 0.71 CG: 0.44 Gestational DM IG: NR

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
								CG: NR Multifetal Gestation IG: 0.82 CG: 1.0
Scazzocchio, 2017 ¹⁰⁵ Good	Aged 18 years or older, singleton pregnancy, 11 to 14 weeks' gestation, crown-rump length of 45 to 84 mm, mean UtA pulsatility index > 95th percentile	Preexisting HTN, immune, renal, or cardiovascular disease, history of preeclampsia in a previous pregnancy, history of gastric ulcer, known allergy or hypersensitivity to aspirin, hemorrhagic disease, fetal malformation (including chromosomopathy), active treatment with heparin or aspirin before recruitment, multifetal gestation, maternal age <18 years	32.9	NR	IG: 23.5 CG: 23.5	IG: 62.5 CG: 64.0	IG: 11.3 CG: 12.0	Previous PreE IG: 0 CG: 0 Chronic HTN IG: 0 CG: 0

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
Schiff, 1989 ⁸⁸ Good	At least one of the following risk factors: nulliparity, twin gestation, history of PreE; had to screen positive during a roll-over test (which tested BP before & after rolling from the left side to back)	History of chronic HTN, long-term treatment with nonsteroidal anti-inflammatory drugs or the use of these drugs in the previous 6 weeks, PIH detected prior to screening, proteinuria detected before screening, history of thrombocytopenia, coagulation disorders, heart failure, chronic renal or pulmonary disease, hepatic or peptic ulcer disease, history of hypersensitivity to ASA	27.4	White: 100	NR	NR	NR	Previous PreE IG: 17.6 CG: 16.1 Chronic HTN IG: 0 CG: 0 Multifetal Gestation IG: 8.8 CG: 6.5
Sibai, 1993 ⁹¹ Good	Nulliparous, 13 to 25 weeks' gestation, BP <135/85 mmHg, no proteinuria	History of chronic HTN, renal disease, DM, or other medical illnesses	20.5	White: 17.9 AA: 49.8 Hispanic: 31.6	NR	IG: 100 CG: 100	IG: 11.5 CG: 10.7	Chronic HTN IG: 0 CG: 0 Chronic DM IG: 0 CG: 0 Multifetal Gestation IG: 1.3 CG: 1.2

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
Subtil, 2003 ¹⁰¹ Good	Nulliparous (no previous delivery at or after 22 weeks' gestation), 14 to 20 weeks' gestation, planned to continue prenatal care and give birth in the participating facility, provided written informed consent	History of HTN, potential indication (antiphospholipid antibodies, lupus) for or a contraindication (allergy, frequent hematomas or bleeding, history of hemorrhage during surgery, tooth extraction or other, recent gastric or duodenal ulcer, severe asthma) to aspirin or other anticoagulant treatment during this pregnancy	24.7	NR	IG: 22.5 CG: 22.3	IG: 100 CG: 100	IG: 25.0 CG: 24.9	Chronic HTN IG: 0 CG: 0 Multifetal Gestation IG: 1.2 CG: 1.6
Viinikka, 1993 ⁹² Fair	At risk of developing preeclampsia, defined as having arterial HTN (BP without treatment >140/90 mmHG already before pregnancy), or having severe preeclampsia in a previous pregnancy	None	33.0	NR	NR	IG: 25.2 CG: 23.8	NR	Previous PreE IG: 13.6 CG: 8.6 Chronic HTN IG: 86.4 CG: 91.4
Villa, 2012 ¹⁰³ Fair	Presenting with at least one of the following risk factors: <20 years of age, >40 years of age, BMI >30, chronic HTN (> or equal to 140/90 mmHg or medication for HTN before 20 weeks' gestation), Sjorgren's syndrome, lupus, or history of one of the following: gestational DM, PreE, SGA, fetus	Allergy to aspirin, tobacco smoking during this pregnancy, multiple pregnancy, history of one or more of the following: asthma, peptic ulcer, placental ablation, inflammatory bowel diseases, rheumatoid arthritis, hemophilia or thrombophilia	30.9 (20-40)	NR	IG: 27.9 CG: 29.7	IG: 26.2 CG: 15.0	IG: 0 CG: 0	Previous PreE IG: 32.8 CG: 28.3 Chronic HTN IG: 13.1 CG: 20.0 Gestational DM IG: 6.6 CG: 16.7

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
	mortus (fetal death after 22 weeks' gestation or >500 g weight in a previous pregnancy), and transvaginal color doppler ultrasound at all patients from 12 +0 weeks to 13 +6 weeks' gestation, second degree diastolic notch present eligible for randomization							
Wallenburg, 1986 ⁸⁶ Good	Healthy, nulliparous, uncomplicated pregnancy of 26 weeks' duration, maximum diastolic blood pressure of 80mmHg, positive angiotensin II sensitivity test	History of HTN or cardiovascular renal disease	24 (17-38)	NR	NR	IG: 100 CG: 100	IG: 21.7 CG: 17.4	Previous PreE IG: 0 CG: 0 Chronic HTN IG: 0 CG: 0 Gestational HTN IG: 0 CG: 0
Yu, 2003 ¹⁰² Good	Mean PI above 1.6, early diastolic notching of uterine arteries identified by transvaginal color Doppler ultrasound	Pre-existing HTN, renal, or cardiovascular disease, DM, bleeding disorders, systemic lupus erythematosus, peptic ulceration, hypersensitivity to aspirin, and the finding at the 23-week scan of a fetal abnormality or fetal growth restriction	29* (23-33)	White: 62.3 AA: 27.6 Other: 10.1	IG: 25.0 CG: 25.6	IG: 26.8 CG: 23.4	IG: 9.4 CG: 9.7	Previous PreE IG: 11.2 CG: 8.6 Chronic HTN IG: 0 CG: 0 Chronic DM

Appendix E Table 1. Study Population Baseline Characteristics of Included Studies

Author, year Quality	Included populations	Excluded populations	Mean age (range)	Race/ethnicity, %	Mean BMI, kg/m2	Nulliparous %	Smoking, %	Other health behaviors and conditions, %
								IG: 0 CG: 0

Abbreviations: AA = African American; ABPM = Ambulatory blood pressure monitoring; AIDS = Acquired immunodeficiency syndrome; ASA = Acetylsalicylic acid; CG = Control group; DBP = Diastolic blood pressure; DM = Diabetes mellitus; Hisp = Hispanic; HTN = Hypertension; IG = Intervention group; IUGR = Intrauterine growth restriction; NR = Not reported; PreE = Preeclampsia; PI = pulsatility index; PIH = Pregnancy-induced hypertension; SGA = Small for gestational age

*Median

Appendix E Table 2. Maternal Health Outcomes, Increased Risk Population

Author, year	Outcome	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95%CI)*
CLASP, 1994 ⁹⁴	Eclampsia	NR	7/4,659 [†] (0.2)	7/4,650 (0.2)	-
Morris, 1996 ¹⁰⁶	Eclampsia	Severe HTN developed postnatally associated with abnormal renal function and hyperuricemia, culminating in a generalized convulsion 6 days after delivery	1/52 (1.9)	0/50 (0.0)	-
Wallenburg, 1986 ⁸⁶	Eclampsia	NR	0/21 (0)	1/23 (4.3)	-
Villa, 2012 ¹⁰³	HELLP Syndrome	Hemolysis, elevated liver enzymes, and low platelets	0/61 (0)	1/60 (0.02)	-
Scazzacchio, 2017 ¹⁰⁵	Incomplete HELLP syndrome [‡]	NR	NR	NR	-

Abbreviations: CG = Control group; CI = Confidence interval; CLASP = Collaborative Low-dose Aspirin Study in Pregnancy; HELLP = Hemolysis, elevated liver enzymes, low platelet count; IG = Intervention group, NR = Not reported

*Dash symbol (“-”) indicates insufficient data to calculate effect

[†] All pregnancies with data

[‡] One participant in each study group developed severe preeclampsia; one with severe hypertension and the other with incomplete HELLP syndrome; authors do not specify to which study group these participants belong

Appendix E Table 3. Effect of Aspirin on Perinatal Mortality, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect
ASPRE, 2017 ⁵⁸	Perinatal mortality without preeclampsia	8/798 (1.0)	14/822 (1.7)	OR*: 0.59 (99% CI: 0.19 to 1.85)
Ayala, 2012 ⁵⁶	Perinatal mortality	2/176 (1.1)	5/174 (2.9)	RR [†] : 0.40 (95% CI: 0.08 to 2.01)
Benigni, 1989 ⁸⁷	Perinatal mortality	0/17 (0.0)	1/16 (6.3)	RR [†] : 0.31 (95% CI: 0.01 to 7.21)
Caspi, 1994 ⁹³	Perinatal mortality	2/48 (4.2)	2/46 (4.3)	RR [†] : 0.96 (95% CI: 0.14 to 6.52)
CLASP, 1994 ⁹⁴	Stillbirths included all intrauterine deaths at or after 24 weeks, and neonatal deaths included all deaths after birth up to the age of 28 days.	77/4123 (1.9)	97/4134 (2.3)	RR [†] : 0.80 (95% CI: 0.59 to 1.07)
Davies, 1995 ⁹⁵	Perinatal mortality	0/58 (0.0)	0/60 (0.0)	RR ^{†‡} : 1.03 (95% CI: 0.02 to 51.26)
Gallery, 1997 ⁹⁶	Perinatal mortality	4/58 (6.9)	2/50 (4.0)	RR [†] : 1.72 (95% CI: 0.33 to 9.02)
Hermida, 1997 ⁹⁷	Perinatal mortality	0/50 (0.0)	0/50 (0.0)	RR ^{†‡} : 1.00 (95% CI: 0.02 to 49.44)
McParland, 1990 ⁸⁹	Perinatal mortality	1/48 (2.1)	3/52 (5.8)	RR [†] : 0.36 (95% CI: 0.04 to 3.35)
MFMU, 1998 ⁹⁸	Perinatal mortality	43/1254 (3.4)	56/1249 (4.5)	RR: 0.80 (95% CI: 0.50 to 1.10)
Morris, 1996 ¹⁰⁶	Perinatal mortality	0/52 (0.0)	0/50 (0.0)	NR
Schiff, 1989 ⁸⁸	Perinatal mortality	0/34 (0.0)	0/32 (0.0)	RR ^{†‡} : 0.94 (95% CI: 0.02 to 46.16)
Viinikka, 1993 ⁹²	Perinatal mortality	2/97 (2.1)	0/100 (0.0)	RR [†] : 5.15 (95% CI: 0.25 to 105.98)
Wallenburg, 1986 ⁸⁶	Perinatal mortality	1/21 (4.8)	1/23 (4.3)	RR [†] : 1.10 (95% CI: 0.07 to 16.43)
Yu, 2003 ¹⁰²	Perinatal mortality	7/276 (2.5)	4/278 (1.4)	OR [§] : 1.89 (95% CI: 0.52 to 6.83)

Abbreviations: CG = Control group; CI = Confidence interval; IG = Intervention group; OR = Odds ratio; RR = Risk ratio

*Adjustment for the effect of the estimated risk for preeclampsia at screening and the participating center

[†]Crude calculation

Appendix E Table 3. Effect of Aspirin on Perinatal Mortality, Increased Risk Population

‡Excluded from pooled analysis.

§Adjusted for Maternal age, parity, ethnicity, education, smoking status, partner, previous pre-eclampsia, family history of pre-eclampsia, body mass index, blood pressure at randomization, mean pulsatility index, and the presence of unilateral or bilateral notch

Appendix E Table 4. Effect of Aspirin on Preterm Birth, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
ASPRE, 2017 ⁵⁸	Delivery at <37 wks of gestation	53/798 (6.6)	84/822 (10.2)	RR*: 0.65 (95% CI: 0.47 to 0.90)
Ayala, 2012 ⁵⁶	Delivery at <37 wks of gestation	7/176 (4.0)	20/174 (11.5)	RR*: 0.35 (95% CI: 0.15 to 0.80)
Benigni, 1989 ⁸⁷	Delivery at <37 wks of gestation	2/17 (11.8)	5/16 (31.3)	RR*: 0.38 (95% CI: 0.08 to 1.67)
Caspi, 1994 ⁹³	Delivery at <37 wks of gestation	11/24 (45.8)	14/23 (60.9)	RR*: 0.75 (95% CI: 0.44 to 1.30)
CLASP, 1994 ⁹⁴	Delivery at <37 wks of gestation	686/3992 (17.2)	761/3982 (19.1)	RR*: 0.90 (95% CI: 0.82 to 0.99)
Davies, 1995 ⁹⁵	Delivery at <34 wks of gestation	1/58 (1.7)	1/60 (1.7)	RR*: 1.03 (95% CI: 0.07 to 16.15)
Gallery, 1997 ⁹⁶	Delivery at <37 wks of gestation	6/58 (10.3)	8/50 (16.0)	RR*: 0.65 (95% CI: 0.24 to 1.74)
Hermida, 1997 ⁹⁷	Delivery at <37 wks of gestation	1/50 (2.0)	5/50 (10.0)	RR*: 0.20 (95% CI: 0.02 to 1.65)
MFMU, 1998 ⁹⁸	Delivery at <37 wks of gestation	502/1254 (40.0)	537/1249 (43.0)	RR: 0.90 (95% CI: 0.90 to 1.00)
Morris, 1996 ¹⁰⁶	Delivery at <37 wks of gestation	3/52 (5.8)	5/50 (10.0)	NR
Schiff, 1989 ⁸⁸	Delivery at <37 wks of gestation	2/34 (5.9)	6/32 (18.8)	RR*: 0.31 (95% CI: 0.07 to 1.44)
Wallenburg, 1986 ⁸⁶	Delivery at <37 wks of gestation	0/21 (0.0)	4/23 (17.4)	RR*: 0.12 (95% CI: 0.01 to 2.12)
Yu, 2003 ¹⁰²	Delivery at <37 wks of gestation	67/276 (24.3)	75/278 (27.0)	OR†: 0.86 (95% CI: 0.57 to 1.29)
ASPRE, 2017 ⁵⁸	Delivery at <37 wks of gestation with preeclampsia	13/798 (1.6)	35/822 (4.3)	OR‡: 0.38 (95% CI: 0.20 to 0.74)
ASPRE, 2017 ⁵⁸	Delivery at <34 wks of gestation with preeclampsia	3/798 (0.4)	15/822 (1.8)	OR‡: 0.18 (99% CI: 0.03 to 1.03)

Abbreviations: CG = Control group; CI = Confidence interval; IG = Intervention group; OR = Odds ratio; RR = Risk ratio; wks = Weeks

*Crude calculation

†Maternal age, parity, ethnicity, education, smoking status, partner, previous pre-eclampsia, family history of pre-eclampsia, body mass index, blood pressure at randomization, mean pulsatility index, and the presence of unilateral or bilateral notch.

‡Adjustment for the effect of the estimated risk for preeclampsia at screening and the participating center

Appendix E Table 5. Effect of Aspirin on Intrauterine Growth Restriction, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
ASPRE, 2017 ⁵⁸	<10 th percentile	148/785 (18.9)	187/807 (23.2)	OR*: 0.77 (99% CI: 0.56 to 1.06)
Ayala, 2012 ⁵⁶	Monthly echography assessments in all participants	16/176 (9.1)	32/174 (18.4)	RR†: 0.49 (95% CI: 0.28 to 0.87)
Benigni, 1989 ⁸⁷	<10 th percentile	2/17 (11.8)	6/16 (37.5)	RR†: 0.31 (95% CI: 0.07 to 1.33)
Caspi, 1994 ⁹³	<10 th percentile	6/48 (12.5)	11/46 (23.9)	RR†: 0.52 (95% CI: 0.21 to 1.30)
CLASP, 1994 ⁹⁴	<3 rd percentile	244/4123 (5.9)	272/4134 (6.6)	RR†: 0.90 (95% CI: 0.76 to 1.06)
Davies, 1995 ⁹⁵	5 – 10 th percentile	3/58 (5.2)	3/60 (5.0)	RR†: 1.03 (95% CI: 0.22 to 4.92)
Hermida, 1997 ⁹⁰	NR	1/50 (2.0)	2/50 (4.0)	RR†: 0.50 (95% CI: 0.05 to 5.34)
McParland, 1990 ⁸⁹	<5 th percentile	7/48 (14.6)	7/52 (13.5)	RR†: 1.08 (95% CI: 0.41 to 2.86)
MFMU, 1998 ⁹⁸	<10 th percentile	129/1254 (10.3)	108/1249 (8.6)	RR: 1.20 (95% CI: 0.90 to 1.50)
Morris, 1996 ¹⁰⁶	<10 th percentile	14/52 (26.9)	11/50 (22.0)	NR
Scazzocchio, 2017 ¹⁰⁵	<10 th percentile	7/80 (8.8)	13/75 (17.3)	RR†: 0.50 (95% CI: 0.21 to 1.20)
Schiff, 1989 ⁸⁸	<10 th percentile	2/34 (5.9)	6/31 (19.4)	RR†: 0.30 (95% CI: 0.07 to 1.40)
Viinikka, 1993 ⁹²	Birth weight ≤2 SD	4/97 (4.1)	9/100 (9.0)	RR†: 0.46 (95% CI: 0.15 to 1.44)
Villa, 2012 ¹⁰³	Birth weight <-2 SD	2/61 (3.3)	6/60 (10.0)	RR: 0.50 (95% CI: 0.10 to 1.90)
Wallenburg, 1986 ⁸⁶	<10 th percentile	4/21 (19.0)	6/23 (26.1)	RR†: 0.73 (95% CI: 0.24 to 2.23)
Yu, 2003 ¹⁰²	<5 th percentile	61/276 (22.1)	68/278 (24.5)	OR‡: 0.83 (95% CI: 0.54 to 1.28)

Abbreviations: CG = Control group; CI = Confidence interval; IG = Intervention group; NR = Not reported; OR = Odds ratio; RR = Risk ratio; SD = Standard deviation

Appendix E Table 5. Effect of Aspirin on Intrauterine Growth Restriction, Increased Risk Population

*Adjusted for the effect of the estimated risk for preeclampsia at screening and the participating center

†Crude calculation

‡Adjusted for maternal age, parity, ethnicity, education, smoking status, partner, previous pre-eclampsia, family history of pre-eclampsia, body mass index, blood pressure at randomization, mean pulsatility index, and the presence of unilateral or bilateral notch

Appendix E Table 6. Effect of Aspirin on Birth Weight, Increased Risk Population

Author, year	Outcome definition	IG mean (SD)	CG mean (SD)	Effect (95% CI)*
Ayala, 2012 ⁵⁶	Birth weight, g	Mean: 3286 (519)	Mean: 3162 (580)	Calc MeanDiff: 124.00 (95% CI: 8.72 to 239.28)
Benigni, 1989 ⁸⁷	Birth weight, g	Mean: 2922 (599)	Mean: 2264 (1072)	Calc MeanDiff: 658.00 (95% CI: 70.24 to 1245.76)
Caspi, 1994 ⁹³	Birth weight, g	Mean: 2502 (561)	Mean: 2108 (542)	Calc MeanDiff: 394.00 (95% CI: 170.86 to 617.14)
CLASP, 1994 ⁹⁴	Birth weight, g	Mean: 3024 (788)	Mean: 2991 (810)	MeanDiff: 32.00 (95% CI: 0.00 to 64.00)
Gallery, 1997 ⁹⁶	Birth weight, g ⁺	Mean: 3291 (514.3928)	Mean: 3029 (714.1779)	Calc MeanDiff: 262.00 (95% CI: 24.09 to 499.91)
Grab, 2000 ¹⁰⁰	Birth weight, g	Median: 3150 (Range: 560 to 3770)	Median: 2900 (Range: 800 to 4070)	-
Hermida, 1997 ⁹⁷	Birth weight, g	Mean: 3265 (452.5483)	Mean: 3155 (678.8225)	Calc MeanDiff: 110.00 (95% CI: -116.14 to 336.14)
McParland, 1990 ⁸⁹	Birth weight, g	Mean: 3068 (555)	Mean: 2954 (852)	Calc MeanDiff: 114.00 (95% CI: -170.39 to 398.39)
Morris, 1996 ¹⁰⁶	Birth weight, g	Mean: 3145 (633)	Mean: 3049 (576)	NR
Scazzocchio, 2017 ¹⁰⁵	Birth weight, g	Mean: 3112 (526)	Mean: 3117 (529)	Calc MeanDiff: -5.00 (95% CI: -171.16 to 161.16)
Schiff, 1989 ⁸⁸	Adjusted birth weight percentile [†]	Mean: 43.9 (13.7)	Mean: 38.6 (12.2)	Calc MeanDiff: 5.30 (95% CI: -1.03 to 11.63)
Schiff, 1989 ⁸⁸	Birth weight, g	Mean: 3037 (NR)	Mean: 2706 (NR)	-
Viinikka, 1993 ⁹²	Birth weight, g	Mean: 3348 (707)	Mean: 3170 (665)	Calc MeanDiff: 178.00 (95% CI: -13.61 to 369.61)
Villa, 2012 ¹⁰³	Birth weight, g	Mean: 3413 (630)	Mean: 3321 (871)	Calc MeanDiff: 92.00 (95% CI: -178.52 to 362.52)
Wallenburg, 1986 ⁸⁶	Birth weight, g	Median: 3190 (Range: 2380 to 4320)	Median: 3040 (Range: 530 to 4035)	-

Abbreviations: Calc MeanDiff = Calculated Mean Difference; CG = Control group; CI = Confidence interval; IG = Intervention group; SD = Standard deviation

*Dash symbol (“-”) indicates insufficient data to calculate effect

†Adjusted for gestational age at delivery

Appendix E Table 7. Effect of Aspirin on Gestational Age, Increased Risk Population

Author, year	Outcome description	IG mean (SD)	CG mean (SD)	Effect (95% CI)*
Ayala, 2012 ⁵⁶	Monthly echography assessments in all participants	Mean: 39.5 (1.6)	Mean: 39.2 (1.9)	Calc MeanDiff: 0.30 (95% CI: -0.07 to 0.67)
Benigni, 1989 ⁸⁷	Duration of pregnancy (weeks)	Mean: 39 (2)	Mean: 35 (4)	Calc MeanDiff: 4.00 (95% CI: 1.86 to 6.14)
Caspi, 1994 ⁹³	Gestational age at birth (wk), mean; gestational age was determined by last menstrual period, ovulation or ovum pickup in in vitro fertilization (IVF) cases, and confirmed by first or early second trimester ultrasound scan	Mean: 36 (31)	Mean: 35 (2.3)	Calc MeanDiff: 1.00 (95% CI: -11.71 to 13.71)
CLASP, 1994 ⁹⁴	Average duration of pregnancy (weeks)	Mean: 38.15 (3.82)	Mean: 37.99 (3.93)	Calc MeanDiff: 0.16 (95% CI: 0.00 to 0.32)
Davies, 1995 ⁹⁵	Mean age of birth (weeks)	Mean: 39.9 (NR)	Mean: 39.6 (NR)	-
Gallery, 1997 ⁹⁶	Mean gestational age at delivery (calculated without 4 intrauterine deaths, as the extent of postmortem weight alteration could not be assessed)	Mean: 38.6 (1.469694)	Mean: 37.5 (2.828427)	Calc MeanDiff: 1.10 (95% CI: 0.24 to 1.96)
Grab, 2000 ¹⁰⁰	Gestational age at delivery (wks)	Median: 39.7 (Range: 26.6 to 42.1)	Median: 38.6 (Range: 28.3 to 41.4)	-
Hermida, 1997 ⁹⁷	Gestational age at delivery (wks)	Mean: 39.6 (1.414214)	Mean: 39.4 (2.12132)	Calc MeanDiff: 0.20 (95% CI: -0.51 to 0.91)
McParland, 1990 ⁸⁹	Gestation at delivery (wks)	Mean: 39.5 (2.1)	Mean: 38.7 (3.9)	Calc MeanDiff: 0.80 (95% CI: -0.44 to 2.04)
Schiff, 1989 ⁸⁸	Gestational age at birth (wks)	Mean: 38.9 (1.14)	Mean: 37.3 (.86)	Calc MeanDiff: 1.60 (95% CI: 1.11 to 2.09)
Viinikka, 1993 ⁹²	Age of gestation at delivery (weeks)	Mean: 38.6 (2.1)	Mean: 38.2 (2)	Calc MeanDiff: 0.40 (95% CI: -0.17 to 0.97)
Wallenburg, 1986 ⁸⁶	Week of delivery	Median: 40 (Range: 37 to 42)	Median: 39 (Range: 30 to 43)	-

Abbreviations: Calc MeanDiff = Calculated Mean Difference; CG = Control group; CI = Confidence interval; IG = Intervention group; SD = Standard deviation

*Dash symbol (“-”) indicates insufficient data to calculate effect

Appendix E Table 8. Effect of Aspirin on Preeclampsia, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
ASPRE, 2017 ⁵⁸	SBP \geq 140 mmHg and/or DBP \geq 90 mmHg on at least two occasions four hours apart developing after 20 weeks of gestation in previously normotensive women. HTN should be accompanied by proteinuria \geq 300 mg in 24 hours or two readings of at least ++ on dipstick analysis of midstream or catheter urine specimens if no 24-hour collection is available. In preeclampsia superimposed on chronic HTN significant proteinuria (as defined above) should develop after 20 weeks' gestation in women with known chronic HTN (history of HTN before conception or the presence of HTN at the booking visit before 20 weeks' gestation in the absence of trophoblastic disease)	66/798 (8.3)	94/822 (11.4)	RR*: 0.72 (95% CI: 0.54 to 0.98)
Ayala, 2012 ⁵⁶ (657)	Hyperbaric index consistently above the threshold for diagnosis of HTN in pregnancy after the 20th wk of gestation and proteinuria, \geq 300 mg/24-h urine, diagnosed after the 20th wk of gestation in a previously normotensive woman	11/176 (6.3)	22/174 (12.6)	RR*: 0.49 (95% CI: 0.25 to 0.99)
Caspi, 1994 ⁹³ (114)	Sustained elevation of more than 15 mmHg of DBP or 30 mmHg of SBP (on at least two occasions six or more hours apart) compared to levels in the second trimester before treatment, presence of proteinuria of more than 300 mg per day (in the absence of urinary tract infection associated with HTN)	0/24 (0.0)	2/23 (8.7)	RR*: 0.19 (95% CI: 0.01 to 3.80)
CLASP, 1994 ⁹⁴	Development of HTN and proteinuria after randomization. For those with baseline DBP below 90 mmHg, HTN defined as a rise of at least 25 mmHg, to 90 mmHg or higher. For those with an initial DBP of 90 mmHg or above, an increment of at least 15 mmHg was required. Proteinuria defined as the appearance after randomization of at least 1+ on protein stick-testing during pregnancy, without evidence of urinary tract infection.	267/3992 (6.7)	302/3982 (7.6)	RR*: 0.88 (95% CI: 0.75 to 1.03)
Davies, 1995 ⁹⁵	Two DBP readings of $>$ 90 mmHg at least 6 hours apart or a single reading of $>$ 110 mmHg, proteinuria defined as ++ on reagent strip testing or greater than 300 mg/L in a 24-h urine collection.	5/58 (8.6)	7/60 (11.7)	OR: 0.70 (95% CI: 0.20 to 2.40)
Grab, 2000 ¹⁰⁰	Proteinuric HTN	3/22 (13.6)	2/21 (9.5)	RR*: 1.43 (95% CI: 0.27 to 7.73)

Appendix E Table 8. Effect of Aspirin on Preeclampsia, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Hermida, 1997 ⁹⁷	Gestational HTN and proteinuria, above 300 mg/24 h, with or without edema	3/50 (6.0)	7/50 (14.0)	RR*: 0.43 (95% CI: 0.12 to 1.56)
McParland, 1990 ⁸⁹	Proteinuric HTN, specific SBP/DBP, proteinuria not defined	1/48 (2.1)	10/52 (19.2)	RR*: 0.11 (95% CI: 0.01 to 0.81)
MFMU, 1998 ⁸⁸	Women who did not have HTN or proteinuria at BL who developed HTN plus one of the following: proteinuria, thrombocytopenia, or pulmonary edema. HTN defined as either SBP \geq 140 mmHg or DBP \geq 90 mmHg on two occasions at least 4 hours apart. Proteinuria defined as excretion of 300 mg of protein in a 24-hour urine collection, or two dipstick-test results of \geq 2+ (\geq 100 mg per deciliter), the values recorded at least four hours apart, with no evidence of urinary tract infection. Thrombocytopenia defined as a platelet count of less than 100,000 per cubic millimeter.	226/1254 (18.0)	250/1249 (20.0)	RR: 0.90 (95% CI: 0.80 to 1.10)
Morris, 1996 ¹⁰⁶	PIH plus proteinuria (1+ or more on dipstick testing on at least two occasions 6 hours apart) or hyperuricemia	4/52 (7.7)	7/50 (14.0)	NR
Scazzocchio, 2017 ¹⁰⁵	SBP \geq 140 mmHg or DBP \geq 90 mmHg on two readings at least four hours apart in previously normotensive women after 20 weeks gestation, and proteinuria >300 mg/24 hr	4/80 (5.0)	3/75 (4.0)	RR*: 1.25 (95% CI: 0.29 to 5.40)
Scazzocchio, 2017 ¹⁰⁵	Severe preeclampsia, defined as: BP \geq 160/110 mmHg on two or more occasions, proteinuria \geq 5g/24 hr, or the presence of maternal complications including: eclampsia, HELLP syndrome (lactate dehydrogenase > 600 IU/L, aspartate transaminase > 62 IU/L, platelet count <100 x 10 ⁹ /L), acute renal failure (creatinine > 1.2 mg/dL), subcapsular hepatic hematoma, pulmonary edema, placental abruption, or the presence of disseminated intravascular disease	1/80 (1.3)	1/75 (1.3)	RR*: 0.94 (95% CI: 0.06 to 14.72)
Schiff, 1989 ⁸⁸	SBP > 140 mmHg, DBP > 90 mmHg, or both when measured on at least two occasions within 24 hours of each other; accompanied by proteinuria (>1 g in 24 hours)	1/34 (2.9)	7/31 (22.6)	RR*: 0.13 (95% CI: 0.02 to 1.00)
Viinikka, 1993 ⁹²	Exacerbation of preexisting HTN (>160/120 mmHg) or the emergence of HTN (>160/110 mmHg) in women normotensive before pregnancy, and proteinuria > 300 mg/24 hours.	9/97 (9.3)	11/100 (11.0)	RR*: 0.84 (95% CI: 0.37 to 1.95)

Appendix E Table 8. Effect of Aspirin on Preeclampsia, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Villa, 2012 ¹⁰³	Blood pressure > or equal to 140 and/or 90 mmHg in two consecutive measurements and proteinuria > or equal to 0.3 g/24 hours	8/61 (13.1)	11/60 (18.3)	RR: 0.80 (95% CI: 0.40 to 1.80)
Villa, 2012 ¹⁰³	Early preeclampsia; preeclampsia diagnosed before 34 + 0 weeks of gestation	1/61 (1.6)	4/60 (6.7)	RR: 0.50 (95% CI: 0.10 to 2.60)
Villa, 2012 ¹⁰³	Preterm preeclampsia; preeclampsia diagnosed before 37 + 0 weeks of gestation	3/61 (4.9)	5/60 (8.3)	RR: 0.80 (95% CI: 0.20 to 2.80)
Villa, 2012 ¹⁰³	SBP ≥160 and/or DBP ≥110 and/or proteinuria ≥5 g/24 hours)	3/61 (4.9)	8/60 (13.3)	RR: 0.50 (95% CI: 0.20 to 1.60)
Wallenburg, 1986 ⁸⁶ (95)	DBP >95 mmHg on at least two occasions six or more hours apart and concomitant proteinuria (0.5 g/L) in the absence of a urinary tract infection	0/21 (0.0)	7/23 (30.4)	RR*: 0.07 (95% CI: 0.00 to 1.20)
Yu, 2003 ¹⁰²	Two recordings of DBP of 90 mmHg or higher at least four hours apart or one recording of DBP of at least 120 mmHg, in a previously normotensive woman, and a urine protein excretion of at least 300 mg in 24h or two readings of 2+ or higher on dipstick analysis of midstream or catheter urine specimens if no 24h collection is available.	49/276 (17.8)	52/278 (18.7)	OR†: 0.88 (95% CI: 0.56 to 1.40)

Abbreviations: BL = Baseline; CG = Control group; CI = Confidence interval; DBP = Diastolic blood pressure; HELLP = Hemolysis, Elevated Liver enzyme levels, and Low Platelet levels; hr = Hour; IG = Intervention group; SBP = Systolic blood pressure; OR = Odds ratio; RR = Risk ratio

*Crude calculation

†Adjusted for maternal age, parity, ethnicity, education, smoking status, partner, previous pre-eclampsia, family history of pre-eclampsia, body mass index, blood pressure at randomization, mean pulsatility index, and the presence of unilateral or bilateral notch.

Appendix E Table 9. Effect of Aspirin on Gestational Hypertension, Increased Risk Population

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
ASPRE, 2017 ⁵⁸	SBP should be 140 mmHg or more and/or the DBP should be 90 mmHg or more on at least two occasions four hours apart developing after 20 weeks of gestation in previously normotensive women.	80/798 (10.0)	69/822 (8.4)	RR*: 1.19 (95% CI: 0.88 to 1.62)
Ayala, 2012 ⁵⁶	Hyperbaric index (HBI): total area of BP excess summed over the 24-h period above the upper limit of the time-varying tolerance interval calculated as a function of gestational age; consistently above the threshold for diagnosis of HTN in pregnancy after the 20 th wk of gestation	26/176 (14.8)	49/174 (28.2)	RR*: 0.52 (95% CI: 0.34 to 0.80)
Benigni, 1989 ⁸⁷	Pregnancy-induced HTN, blood pressure levels higher than 140/90 mmHg in women whose blood pressure had previously been normal.	0/17 (0.0)	3/16 (18.8)	RR*: 0.13 (95% CI: 0.01 to 2.42)
Caspi, 1994 ⁹³	Sustained elevation of more than 15 mmHg of DBP or 30 mmHg of SBP (on at least two occasions 6 or more hours apart) compared to levels in the second trimester before treatment.	1/24 (4.2)	6/23 (26.1)	RR*: 0.16 (95% CI: 0.02 to 1.23)
Davies, 1995 ⁹⁵	Two DBP readings of > 90 mmHg at least 6 hours apart or a single reading of > 110 mmHg	2/58 (3.4)	4/60 (6.7)	RR*: 0.52 (95% CI: 0.10 to 2.72)
Hermida, 1997 ⁹⁷	Conventional BP values above 140/90 mmHg for systolic/diastolic BP without clinical record of HTN previous to pregnancy	5/50 (10.0)	9/50 (18.0)	RR*: 0.56 (95% CI: 0.20 to 1.54)
McParland, 1990 ⁸⁹	Pregnancy-induced HTN; specific SBP/DPB not defined	6/48 (12.5)	13/52 (25.0)	RR*: 0.50 (95% CI: 0.21 to 1.21)
Morris, 1996 ¹⁰⁶	Antenatal blood pressure > 140/90, with a rise in DBP of at least 15 mmHg on at least two occasions 6 hours apart	8/52 (15.4)	7/50 (14.0)	NR
Villa, 2012 ¹⁰³	New onset HTN after 20 weeks gestation	10/61 (16.4)	6/60 (10.0)	RR: 1.40 (95% CI: 0.60 to 3.50)

Abbreviations: CI = Confidence interval; CG = Control group; DBP = Diastolic blood pressure; HBI = Hyperbaric index; HTN = Hypertension; IG = Intervention group; RR = Risk ratio; SBP = Systolic blood pressure; wk = Week

*Crude calculation

Appendix E Table 10. Harms of Maternal Bleeding, Average and Increased Risk Populations

Author, year	Outcome	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Ayala, 2012 ⁵⁶	Antepartum hemorrhage	NR	6/176 (3.4)	9/174 (5.2)	RR*: 0.66 (95% CI: 0.24 to 1.81)
Sibai, 1993 ⁹¹	Bleeding During Delivery	Excessive blood loss (>500 mL for vaginal delivery and >1000 mL for a cesarean delivery)	102/1485 (6.9)	109/1500 (7.3)	RR*: 0.95 (95% CI: 0.73 to 1.23)
CLASP, 1994 ⁹⁴	Blood Transfusion Required	NR	188/4659 (4.0)	147/4650 (3.2)	
Mone, 2018 ¹⁰⁴	Blood Transfusion Required	NR	3/192 (1.6)	4/354 (1.1)	OR: 0.50 (95% CI: 0.10 to 2.70)
Morris, 1996 ¹⁰⁶	Antepartum hemorrhage	NR	2/52 (3.8)	1/50 (2.0)	NR
Rotchell, 1998 ⁹⁹	Blood Transfusion Required	NR	19/1819 (1.0)	18/1822 (1.0)	
Sibai, 1993 ⁹¹	Blood Transfusion Required	NR	7/1485 (0.5)	10/1500 (0.7)	RR*: 0.71 (95% CI: 0.27 to 1.85)
Sibai, 2003 ¹⁰¹	Blood Transfusion Required	NR	3/1634 (0.2)	9/1640 (0.5)	
Yu, 2003 ¹⁰²	Blood Transfusion Required	NR	6/276 (2.2)	7/278 (2.5)	
Ayala, 2012 ⁵⁶	Postpartum hemorrhage	NR	3/176 (1.7)	6/174 (3.4)	RR*: 0.49 (95% CI: 0.13 to 1.95)
CLASP, 1994 ⁹⁴	Postpartum hemorrhage	Placental bleed \geq 500 mL	1200/4659 (25.8)	1182/4650 (25.4)	RR*: 1.01 (95% CI: 0.95 to 1.09)
Hermida, 1997 ⁹⁷	Postpartum hemorrhage	Antepartum hemorrhage, no complication with maternal bleeding	0/50 (0.0)	0/50 (0.0)	RR*†: 1.00 (95% CI: 0.02 to 49.44)
MFMU, 1998 ⁹⁸	Postpartum hemorrhage	NR	73/1254 (5.8)	77/1249 (6.2)	RR: 0.90 (95% CI: 0.70 to 1.30)
Mone, 2018 ¹⁰⁴	Postpartum hemorrhage	>500 mL blood loss	25/179 (14.0)	9/183 (4.9)	RR*: 2.84 (95% CI: 1.36 to 5.91)
Mone, 2018 ¹⁰⁴	Postpartum hemorrhage	>1000 mL blood loss	7/179 (3.9)	5/183 (2.7)	RR*: 1.43 (95% CI: 0.46 to 4.43)
Rotchell, 1998 ⁹⁹	Postpartum hemorrhage	\geq 500 mL blood loss	178/1819 (9.8)	175/1822 (9.6)	RR*: 1.02 (95% CI: 0.84 to 1.24)
Scazzocchio, 2017 ¹⁰⁵	Postpartum hemorrhage	NR	2/80 (2.5)	5/75 (6.7)	RR*: 0.38 (95% CI: 0.08 to 1.87)

Appendix E Table 10. Harms of Maternal Bleeding, Average and Increased Risk Populations

Author, year	Outcome	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Sibai, 1993 ⁹¹	Postpartum hemorrhage	NR	45/1485 (3.0)	37/1500 (2.5)	RR*: 1.23 (95% CI: 0.80 to 1.89)
Subtil, 2003 ¹⁰¹	Postpartum hemorrhage	≥500 mL blood loss	63/1634 (3.9)	49/1640 (3.0)	RR*: 1.29 (95% CI: 0.89 to 1.86)
Wallenburg, 1986 ⁸⁶	Postpartum hemorrhage	Postpartum hemorrhage	0/21 (0.0)	0/23 (0.0)	RR*†: 1.09 (95% CI: 0.02 to 52.67)
Yu, 2003 ¹⁰²	Postpartum hemorrhage	>500 mL blood loss	73/276 (26.4)	71/278 (25.5)	OR‡: 1.09 (95% CI: 0.73 to 1.62)

Abbreviations: CG = Control group; CI = Confidence interval; IG = Intervention group; NR = Not reported; OR = Odds ratio; RR = Risk ratio

*Crude calculation

†Excluded in the pooled metaanalysis.

‡Adjusted for maternal age, parity, ethnicity, education, smoking status, partner, previous pre-eclampsia, family history of pre-eclampsia, body mass index, blood pressure at randomization, mean pulsatility index, and the presence of unilateral or bilateral notch.

Appendix E Table 11. Harms of Placental Abruption, Average and Increased Risk Populations

Author, year	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
ASPRE, 2017 ⁵⁸	Placental abruption without preeclampsia	4/798 (0.5)	6/822 (0.7)	RR*: 0.69 (95% CI: 0.19 to 2.42)
Caspi, 1994 ⁹³	Placental abruption	0/24 (0.0)	0/23 (0.0)	RR*†: 0.96 (95% CI: 0.02 to 46.47)
CLASP, 1994 ⁹⁴	Placental abruption	86/4659 (1.8)	71/4650 (1.5)	RR*: 1.21 (95% CI: 0.89 to 1.65)
Davies, 1995 ⁹⁵	Placental abruption	2/58 (3.4)	1/60 (1.7)	OR: 1.24 (95% CI: 0.20 to 23.90)
Hauth, 1993 ⁹⁰	Placental abruption	1/302 (0.3)	0/302 (0.0)	RR*: 3.00 (95% CI: 0.12 to 73.35)
Hermida, 1997 ⁹⁷	Placental abruption	0/50 (0.0)	0/50 (0.0)	RR*†: 1.00 (95% CI: 0.02 to 49.44)
MFMU, 1998 ⁹⁸	Diagnosed according to clinical criteria: vaginal bleeding, uterine tenderness, as well as examination of the placenta.	17/1254 (1.4)	25/1249 (2.0)	RR: 0.70 (95% CI: 0.40 to 1.30)
Mone, 2018 ¹⁰⁴	Placental abruption	1/179 (0.6)	0/183 (0.0)	RR*: 3.07 (95% CI: 0.13 to 74.78)
Rotchell, 1998 ⁹⁹	Placental abruption	9/1819 (0.5)	14/1822 (0.8)	RR*: 0.64 (95% CI: 0.28 to 1.48)
Sibai, 1993 ⁹¹	Diagnosed according to clinical findings (vaginal bleeding, uterine tenderness) or placental examination	11/1485 (0.7)	2/1500 (0.1)	RR: 5.56 (95% CI: 1.24 to 25.06)
Subtil, 2003 ¹⁰¹	Placental abruption either clinical or detected in a pathology examination of the placenta	13/1634 (0.8)	9/1640 (0.5)	RR: 1.45 (95% CI: 0.62 to 3.38)
Viinikka, 1993 ⁹²	Placental abruption	0/97 (0.0)	0/100 (0.0)	RR*†: 1.03 (95% CI: 0.02 to 51.43)
Yu, 2003 ¹⁰²	Placental abruption	10/276 (3.6)	5/278 (1.8)	OR‡: 2.54 (95% CI: 0.78 to 8.21)

Abbreviations: CG = Control group; CI = Confidence interval; IG = Intervention group; OR = Odds ratio; RR = Risk ratio

*Crude calculation

†Exclude in the pooled metanalysis.

‡Adjusted for maternal age, parity, ethnicity, education, smoking status, partner, previous pre-eclampsia, family history of pre-eclampsia, body mass index, blood pressure at randomization, mean pulsatility index, and the presence of unilateral or bilateral notch.

Appendix E Table 12. Perinatal Harms, Average and Increased Risk Populations

Author, year	Outcome	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Sibai, 1993 ⁹¹	Any bleeding disorder	Any bleeding disorder (infant)	104/1480 (7.0)	98/1505 (6.5)	RR*: 1.08 (95% CI: 0.83 to 1.41)
Schiff, 1989 ⁸⁸	Cephalohematoma	Cephalohematoma	0/34 (0.0)	1/32 (3.1)	RR*: 0.31 (95% CI: 0.01 to 7.45)
Sibai, 1993 ⁹¹	Cephalohematoma	Cephalohematoma	68/1480 (4.6)	55/1505 (3.7)	RR*: 1.26 (95% CI: 0.89 to 1.78)
ASPRE, 2017 ⁵⁸	Congenital malformations	Fetal structural defects	8/798 (1.0)	16/822 (1.9)	RR*: 0.52 (95%CI -0.34 to 1.36)
Gallery, 1997 ⁹⁶	Congenital malformations	Congenital malformations	2/54 (3.7)	4/50 (8.0)	RR*: 0.46 (95% CI: 0.09 to 2.42)
Mone, 2018 ¹⁰⁴	Congenital malformations	Total congenital anomaly (includes cardiac, GI, neurological, renal)	3/179 (1.7)	4/183 (2.2)	RR*: 0.77 (95% CI: 0.17 to 3.38)
Viinikka, 1993 ⁹²	Congenital malformations	Fetal anencephaly	1/97 (1.0)	0/100 (0.0)	RR*: 3.09 (95% CI: 0.13 to 74.98)
ASPRE, 2017 ⁵⁸	Fetal intracranial bleeding	Intraventricular hemorrhage of grade ≥II	2/798 (0.3)	1/822 (0.1)	OR†: 2.23 (99% CI: 0.09 to 52.70)
Benigni, 1989 ⁸⁷	Fetal intracranial bleeding	Intracranial hemorrhage	0/17 (0.0)	0/16 (0.0)	RR*: 0.94 (95% CI: 0.02 to 44.98)
MFMU, 1998 ⁹⁸	Fetal intracranial bleeding	Defined according to ultrasound criteria	21/1254 (1.7)	14/1249 (1.1)	RR: 1.50 (95% CI: 0.80 to 2.80)
Sibai, 1993 ⁹¹	Fetal intracranial bleeding	Cerebral hemorrhage	10/1480 (0.7)	11/1505 (0.7)	RR*: 0.92 (95% CI: 0.39 to 2.17)
CLASP, 1994 ⁹⁴	Fetal intracranial bleeding	Intraventricular hemorrhage	33/4810 (0.7)	45/4821 (0.9)	RR*: 0.74 (95% CI: 0.47 to 1.15)
Rotchell, 1998 ⁹⁹	Fetal intracranial bleeding	Intraventricular hemorrhage	0/1834 (0.0)	1/1841 (0.1)	RR*: 0.33 (95% CI: 0.01 to 8.21)
Schiff, 1989 ⁸⁸	Fetal intracranial bleeding	Intraventricular hemorrhage	0/34 (0.0)	0/32 (0.0)	RR*: 0.94 (95% CI: 0.02 to 46.16)
Subtil, 2003 ¹⁰¹	Fetal intracranial bleeding	Intraventricular hemorrhage grade II, III, or IV	1/1645 (0.1)	6/1660 (0.4)	RR*: 0.17 (95% CI: 0.02 to 1.40)
ASPRE, 2017 ⁵⁸	Respiratory distress syndrome	Respiratory distress syndrome treated with surfactant and ventilation	11/798 (1.4)	22/822 (2.7)	OR†: 0.53 (99% CI: 0.20 to 1.40)

Appendix E Table 12. Perinatal Harms, Average and Increased Risk Populations

Study, Year Quality	Treatment group	N Analyzed	Child outcomes, n (%)
CLASP, 1994 ⁹⁴ Good 18-month followup	IG	2,146	<i>Gross motor failure:</i> 9 (0.4) <i>Fine motor failure:</i> 28 (1.3) <i>Height <3rd percentile:</i> 236 (11.0) <i>Weight <3rd percentile:</i> 112 (5.2)
	CG	2,219	<i>Gross motor failure:</i> 10 (0.5) <i>Fine motor failure:</i> 39 (1.8) <i>Height <3rd percentile:</i> 248 (11.2) <i>Weight <3rd percentile:</i> 129 (5.8)

Author, year	Outcome	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Gallery, 1997 ⁹⁶	Severe hyaline membrane disease	Neonatal death caused by severe hyaline membrane disease	0/58 (0.0)	1/50 (2.0)	RR*: 0.29 (95% CI: 0.01 to 6.92)
Gallery, 1997 ⁹⁶	Staphylococcus epidermis septicemia	One neonatal death from Staphylococcus epidermis septicemia	0/58 (0.0)	1/50 (2.0)	RR*: 0.29 (95% CI: 0.01 to 6.92)

Abbreviations: CG = Control group; CI = Confidence interval; GI = Gastrointestinal; IG = Intervention group; OR = Odds ratio; RR = Risk ratio

*Crude calculation

†Adjustment for the effect of the estimated risk for preeclampsia at screening and the participating center

Appendix E Table 14. Other Maternal Outcomes, Average and Increased Risk Populations

Author, year	Outcome	Outcome definition	IG n/N (%)	CG n/N (%)	Effect (95% CI)
Mone, 2018 ¹⁰⁴	Any Adverse Event	NR	123/192 (64.1)	143/354 (40.4)	OR: 2.60 (95% CI: 1.80 to 3.80)
Mone, 2018 ¹⁰⁴	Serious Adverse Event	NR	32/179 (17.9)	28/183 (15.3)	RR*: 1.17 (95% CI: 0.74 to 1.86)
Morris, 1996 ¹⁰⁶	Any Adverse Event	NR	29/52 (55.7)	28/50 (56.0)	NR
ASPRE, 2017 ⁵⁸	Maternal death	Maternal death due to pulmonary embolism	0/798 (0.0)	1/822 (0.1)	RR*: 0.34 (95% CI: 0.01 to 8.42)
Hauth, 1993 ⁹⁰	Preterm Rupture of Membranes	Preterm premature rupture of membranes	9/302 (3.0)	8/302 (2.6)	RR*: 1.13 (95% CI: 0.44 to 2.88)
Sibai, 1993 ⁹¹	Preterm Rupture of Membranes	Premature rupture of membranes with preterm delivery	49/1485 (3.3)	54/1500 (3.6)	RR*: 0.92 (95% CI: 0.63 to 1.34)

Abbreviations: CG = Control group; CI = Confidence interval; IG = Intervention group; NR = Not reported; OR = Odds ratio; RR = Risk ratio

*Crude calculation

Appendix F Table 1. Within Trial Subgroup Differences in Aspirin Effectiveness Based on Reported Tests for Interaction

Study	Outcome	Subgroup Comparison	N	IG n/N (%)	CG n/N (%)	RR or OR (95%CI)	P value for interaction test*
CLASP, 1994 ⁹⁴	Preeclampsia	Aspirin \leq 20 weeks' gestation	5482	176/2733 (6.4)	222/2749 (8.1)	0.79 (0.66, 0.96)	NS (0.06) [†]
		Aspirin >20 week's gestation	2492	91/1259 (7.2)	80/1233 (6.5)	1.11 (0.83, 1.45)	
		Nulliparous	1837	57/922 (6.2)	62/975 (6.8)	0.91 (0.64, 1.29)	NS
		Multiparous	6137	210/3070 (6.8)	240/3067 (7.8)	0.87 (0.73, 1.04)	
	Preterm birth (<37 weeks gestation)	Aspirin \leq 20 weeks' gestation	5482	477/2733 (17.5)	497/2749 (18.1)	0.97 (0.86, 1.08)	NS (0.03) [†]
		Aspirin >20 week's gestation	2492	209/1259 (16.6)	264/1233 (21.4)	0.78 (0.66, 0.91)	
		Nulliparous	1837	144/922 (15.6)	184/915 (20.1)	0.77 (0.64, 0.95)	NS
		Multiparous	6137	542/3070 (17.7)	577/3067 (18.8)	0.94 (0.84, 1.04)	
	Perinatal mortality	Entered study \leq 20 weeks' gestation	5635	55/2802 (2.0)	62/2833 (2.2)	0.89 (0.63, 1.28)	NS
		Entered study >20 week's gestation	2622	22/1321 (1.7)	35/1301 (2.7)	0.62 (0.37, 1.05)	
		Nulliparous	1961	11/981 (1.1)	22/980 (2.2)	0.50 (0.24, 1.02)	NS
		Multiparous	6296	66/3142 (2.1)	75/3154 (2.4)	0.88 (0.64, 1.23)	
	IUGR	Aspirin \leq 20 weeks' gestation	5635	151/2802 (5.4)	185/2833 (6.5)	0.82 (0.67, 1.02)	NS
		Aspirin >20 week's gestation	2622	93/1321 (7.0)	87/1301 (6.7)	1.05 (0.79, 1.40)	
		Nulliparous	1961	53/981 (5.4)	63/980 (6.4)	0.84 (0.59, 1.20)	NS
		Multiparous	6296	191/3142 (6.1)	209/3154 (6.6)	0.92 (0.76, 1.11)	
MFMU, 1998 ^{98‡}	Preeclampsia [§]	Pregestational diabetes risk group	462	(18)	(22)	0.9 (0.6, 1.2)	NS [†]
		Hypertension risk group	763	(26)	(25)	1.1 (0.8, 1.4)	
		Multifetal gestation risk group	678	(12)	(16)	0.7 (0.5, 1.1)	
		Previous preeclampsia risk group	600	(17)	(19)	0.9 (0.6, 1.2)	
		Race white	814	(18)	(22)	0.8 (0.6, 1.1)	NS [†]
		Race nonwhite	1689	(18)	(20)	0.9 (0.8, 1.2)	
		Nulliparous	668	(25)	(27)	0.9 (0.7, 1.2)	NS [†]
		Multiparous	1835	(16)	(18)	0.9 (0.7, 1.1)	

Study	Outcome	Subgroup Comparison	N	IG n/N (%)	CG n/N (%)	RR or OR (95%CI)	P value for interaction test*
		Systolic BP <120 mmHg at entry ^f	1151	(11)	(14)	0.8 (0.6, 1.1)	NS ⁱ
		Systolic BP 120–134 mmHg at entry ^l	435	(22)	(26)	0.8 (0.6, 1.2)	
		Aspirin <16 weeks' gestation	461	49/225 (21.8)	55/236 (23.3)	0.93 (0.67, 1.31)	0.87
		Aspirin ≥16 weeks' gestation	2042	182/1029 (17.7)	199/1013 (19.6)	0.90 (0.75, 1.08)	
		BMI <30	1512	131/756 (17.3)	144/756 (19.1)	0.91 (0.7, 1.13)	0.85
		BMI ≥30	967	97/487 (19.9)	108/480 (22.5)	0.89 (0.7, 1.13)	
		Nonsmokers	2079	203/1041 (19.5)	218/1038 (21.0)	0.93 (0.78, 1.10)	0.58
		Smokers	421	29/214 (13.6)	35/207 (16.9)	0.80 (0.51, 1.26)	
	Preterm birth (<37 weeks gestation)	Nonsmokers	2079	465/1038 (44.8)	512/1041 (49.2)	0.91 (0.84, 0.99)	0.01
		Smokers	421	100/214 (46.5)	113/207 (54.5)	1.17 (0.99, 1.39)	
	SGA	Nonsmokers	2079	74/1038 (7.1)	74/1041 (7.1)	1.0 (0.7, 1.4)	0.36
		Smokers	421	175/1252 (14.0)	132/1248 (10.6)	1.3 (0.8, 2.2)	
	ASPRE, 2017 ^{58#}	Preterm birth with preeclampsia	Maternal age <30 years	649	7/321 (2.2)	14/331 (4.2)	0.51 (0.20, 1.32)
Maternal age ≥30 years			971	6/477 (1.3)	21/491 (4.3)	0.29 (0.11, 0.75)	
BMI <25 kg/m ²			649	5/311 (1.0)	14/338 (4.1)	0.34 (0.12, 0.99)	0.60
BMI ≥25 kg/m ²			971	8/487 (1.6)	21/484 (4.3)	0.41 (0.18, 0.96)	
Racial origin Afro Caribbean			409	5/208 (2.4)	11/201 (5.5)	0.45 (0.15, 1.39)	0.91
Racial origin Caucasian			1087	7/528 (1.3)	22/559 (3.9)	0.34 (0.14, 0.82)	
Racial origin Other			124	1/62 (0.18)	2/62 (3.2)	0.41 (0.04, 4.85)	
Method conception natural			1526	12/747 (1.6)	32/779 (4.1)	0.40 (0.20, 0.79)	0.13
Method conception assisted			94	1/51 (2.0)	3/43 (7.0)	0.19 (0.02, 2.05)	
Nonsmokers			1504	12/741 (1.6)	32/763 (4.2)	0.39 (0.20, 0.78)	0.79
Smokers			116	1/57 (1.8)	3/59 (5.1)	0.28 (0.03, 3.04)	
Family history preeclampsia			166	2/76 (2.6)	5/90 (5.6)	0.37 (0.18, 2.75)	0.77

Study	Outcome	Subgroup Comparison	N	IG n/N (%)	CG n/N (%)	RR or OR (95%CI)	P value for interaction test*
		No family history of preeclampsia	1454	11/722 (1.5)	30/732 (4.1)	0.49 (0.09, 2.73)	
		Nulliparous	1090	7/547 (1.3)	24/543 (4.4)	0.27 (0.11, 0.64)	0.39
		Multiparous without preeclampsia	359	4/164 (2.4)	7/195 (3.6)	0.79 (0.22, 2.90)	
		Multiparous with preeclampsia	171	2/87 (2.3)	4/84 (4.8)	0.50 (0.08, 2.93)	
		History of chronic hypertension	110	5/49 (10.2)	5/61 (8.2)	1.30 (0.33, 5.12)	NS (0.06) [#]
		No history of chronic hypertension	1510	8/749 (1.1)	30/761 (3.9)	0.27 (0.12, 0.60)	

Abbreviations: BMI = Body mass index; CG = Control group; CI = Confidence interval; IG = Intervention group; IUGR = Intrauterine growth restriction; NS = Not significant; OR = Odds ratio; RR = Risk ratio; SGA = Small for gestational age

[†]The CLASP and MFMU trials reported relative risks and the ASPRE trial reported odds ratios.

[‡]Statistical significance in this trial was set at <.01 for subgroup comparisons.

[§]Comparison numbers calculated from main report, only percentages given


[§]Additional stratified comparisons reported in text, none statistically significant. We report in this table the comparisons discussed in our report.

[¶]Interaction test results not reported, but the authors reported in the text that there were no statistically significant subgroup differences. The study was underpowered, however, for comparing the small magnitude differences between groups.

[¶]Excludes participants with hypertension at baseline


[#]Study reported that p-values for effect estimates should be compared to 0.00625 to determine significance at the 5% level to adjust for multiple comparisons.

Appendix F Figure 1. Pooled Estimate of Mean Birthweight, Grams



Abbreviations: CI = Confidence interval; mg = Milligrams; PreE = Preeclampsia; n = Sample size; NR = Not reported; SD = Standard deviation; WMD = Weighted mean difference


Appendix F Figure 2. Gestational Hypertension, Sorted by Study Size



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; mg = Milligrams; PreE = Preeclampsia; NR = Not reported; RR = Risk ratio


Appendix F Figure 3. Subgroup Analyses of Aspirin Effectiveness Comparisons for Dosage, Timing, Study Design, and Population Characteristics on Small for Gestational Age or Intrauterine Growth Restriction



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; K = Number of studies; mg = milligrams; N = Number of observations; RR= Relative Risk


Appendix F Figure 4. Subgroup Analyses of Aspirin Effectiveness Comparisons for Dosage, Timing, Study Design, and Population Characteristics on Preterm Delivery



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; K = Number of studies; mg = milligrams; N = Number of observations; RR= Relative Risk

Appendix F Figure 5. Subgroup Analyses of Aspirin Effectiveness Comparisons for Dosage, Timing, Study Design, and Population Characteristics on Preeclampsia



Random-effects REML model with Knapp-Hartung confidence intervals

Abbreviations: CI = Confidence interval; K = Number of studies; mg = milligrams; N = Number of observations; RR= Relative Risk

Appendix G. Ongoing Studies

Trial identifier	Study name	Location	Estimated N and Age range	Interventions	Outcome Measures	Status
NCT03735433	The Effect of Two Aspirin Dosing Strategies for Obese Women at High Risk for Preeclampsia	USA	400 18-45 years	ASA, 81 mg vs. ASA, 162 mg	Incidence of Diagnosis of preeclampsia Incidence of aspirin resistance based on incomplete inhibition of TBx2	Recruiting Start date: Jan 2019 Est completion date: Jan 2020
NCT03961360	Effectiveness of Higher Aspirin Dosing for Prevention of Preeclampsia in High Risk Obese Gravida (ASPREO)	USA	220 18-45 years	ASA, 181 mg vs. ASA, 162 mg	Preeclampsia diagnosis Maternal Outcomes-Incidence of preterm preeclampsia Maternal Outcomes-Gestational Hypertension Maternal Outcomes-Placenta Abruptio Maternal Outcomes-Eclampsia Maternal Outcomes-HELLP syndrome Maternal Outcomes-Postpartum Hemorrhage Maternal Outcomes-Other maternal bleeding Maternal Outcomes-Need for blood transfusion Neonatal Outcomes- Gestational age at delivery Neonatal Outcomes-Delivery at < 37 weeks Neonatal Outcomes-Apgar score at 5 min ≤ 5 Neonatal Outcomes-Small for gestational age Neonatal Outcomes-Neonatal Intensive Care Unit (NICU) length of stay Neonatal Outcomes-Intraventricular Hemorrhage Grade III-IV Neonatal Outcomes-Bronchopulmonary Dysplasia Neonatal Outcomes-Necrotizing Enterocolitis	Recruiting Start date: May 2019 Est completion date: May 2020
NCT03893630	Role of ASpirin in Placental and Maternal Endothelial Cell Regulation IN Pre-eclampsia (ASPERIN)	USA	150 18-45 years	ASA, 81 mg vs. ASA, 162 mg vs. Standard of care	Change in Pulsatility Index (PI) Onset of Pre-eclampsia Severity of Pre-eclampsia Composite Neonatal outcomes including frequency of Intraventricular hemorrhage (IVH), Bronchopulmonary dysplasia (BPD),	Recruiting Start date: April 2019 Est

Appendix G. Ongoing Studies

Trial identifier	Study name	Location	Estimated N and Age range	Interventions	Outcome Measures	Status
					Respiratory distress syndrome (RDS), Necrotizing enterocolitis(NEC) Change in s-ICAM levels over time Change in PIGF levels over time Change in CRP levels over time Change in IL-6 over time Change in TNF over time	completion date: June 2020
NCT03574909	IRELANd: Investigating the Role of Early Low-dose Aspirin in Diabetes (IRELANd)	Ireland	300 18 years or older	ASA, 150 mg vs placebo	Preeclampsia Birth weight Preterm birth Perinatal mortality Parameters of neonatal morbidity: Gestational age at delivery Parameters of neonatal morbidity: Birth weight Parameters of neonatal morbidity: NICU admission Respiratory morbidity: Duration of invasive ventilation Duration of O2 Duration of hospital stay Use of nitric oxide Number and type of inotropes Duration of inotrope use Apgar score Umbilical artery acidosis at birth Intracranial hemorrhage Culture-proven sepsis Necrotizing enterocolitis Hypoxic ischemic encephalopathy Shoulder dystocia Composite measure of Maternal outcomes not directly related to primary outcome Hemorrhage Sepsis	Not yet recruiting Start date: Sept 2018 Est completion date: Sept 2021
NCT04158830	Aspirin (ASA) Therapy and Preeclampsia Prevention	USA	900 14-50 years	ASA, 81 mg vs. 2 x ASA, 81 mg	Preeclampsia	Not yet recruiting Start date: Dec 2019 Est completion date: Nov 2021

Appendix G. Ongoing Studies

Trial identifier	Study name	Location	Estimated N and Age range	Interventions	Outcome Measures	Status
NCT04070573	Low Doses of Aspirin in the Prevention of Preeclampsia (ASAPP)	USA	400 18-60 years	ASA, 81 mg vs. ASA, 162 mg	Incidence of preterm (<37 weeks) preeclampsia Incidence of preeclampsia with severe features Aspirin adherence Maternal and Fetal Outcomes Time-to-event for preeclampsia Aspirin compliance Impact of co-morbidities on incidence of preeclampsia	Recruiting Start date: Oct 2019 Est completion date: Aug 2024