

# Effectiveness of Primary Care–Relevant Treatments for Obesity in Adults: A Systematic Evidence Review for the U.S. Preventive Services Task Force

Erin S. LeBlanc, MD, MPH; Elizabeth O'Connor, PhD; Evelyn P. Whitlock, MD, MPH; Carrie D. Patnode, PhD, MPH; and Tanya Kapka, MD, MPH

**Background:** Overweight and obesity in adults are common and adversely affect health.

**Purpose:** To summarize effectiveness and harms of primary care–relevant weight-loss interventions for overweight and obese adults.

**Data Sources:** MEDLINE, Cochrane Central Register of Controlled Trials, and PsycINFO from January 2005 to September 2010; systematic reviews for identifying trials before 2005.

**Study Selection:** Two investigators appraised 6498 abstracts and 648 articles. Clinical trials were included if control groups received minimal interventions. Articles were rated as good, fair, or poor by using design-specific criteria.

**Data Extraction:** One investigator abstracted study characteristics and findings for good- and fair-quality studies; a second checked them.

**Data Synthesis:** Behaviorally based treatment resulted in 3-kg (6.6-lb) greater weight loss in intervention than control participants after 12 to 18 months, with more treatment sessions associated with greater loss. Limited data suggest weight-loss maintenance for 1 year or more. Orlistat plus behavioral intervention resulted in 3-kg (6.6-lb) more weight loss than did placebo after 12 months. Met-

formin resulted in less weight loss. Data on effects of weight-loss treatment on long-term health outcomes (for example, death and cardiovascular disease) were insufficient. Weight-loss treatment reduced diabetes incidence in participants with prediabetes. Effects on intermediate outcomes (for example, lipids and blood pressure) were mixed and small. Data on serious medication harms were insufficient. Medications commonly caused withdrawals due to gastrointestinal symptoms.

**Limitations:** Few studies reported health outcomes. Behaviorally based treatments were heterogeneous and specific elements were not well-described. Many studies could not be pooled because of insufficient reporting of variance data. Medication trials had high attrition, lacked postdiscontinuation data, and were inadequately powered for rare adverse effects.

**Conclusion:** Behaviorally based treatments are safe and effective for weight loss and maintenance.

**Primary Funding Source:** Agency for Healthcare Research and Quality.

*Ann Intern Med.* 2011;155:434-447.  
For author affiliations, see end of text.

www.annals.org

The prevalence of adult obesity—defined as a body mass index (BMI) greater than 30 kg/m<sup>2</sup>—is high in the United States, exceeding 30% in most age- and sex-specific groups. In 2007–2008, 32% of men and 36% of women were obese. In addition, 40% of men and 28% of women met overweight criteria (BMI >25 kg/m<sup>2</sup>) (1). The prevalence of obesity and of overweight have increased by 134% and 48%, respectively, since 1976–1980 (2).

Obesity is associated with increased mortality (particularly in adults <65 years) (3–5), coronary heart disease (6), type 2 diabetes (7), some types of cancer (8), and many other deleterious effects (9). Whether being overweight is

associated with an increased mortality risk is less clear, possibly because the association varies by sex, ethnicity, and age and depends on the obesity measure used (for example, BMI vs. waist circumference) (10–12). Maternal obesity is associated with pregnancy complications and adverse fetal and neonatal health outcomes (13).

In 2003, the U.S. Preventive Services Task Force (USPSTF) recommended that clinicians screen all adults for obesity and offer intensive counseling and behavioral interventions to promote sustained weight loss for obese adults (B recommendation: high certainty that net benefit was moderate or moderate certainty that net benefit was moderate to substantial). The USPSTF, however, concluded that evidence was insufficient to recommend for or against moderate- or low-intensity counseling together with behavioral interventions to promote sustained weight loss in obese adults (I recommendation: insufficient evidence to assess benefit and harm balance). The USPSTF concluded that evidence was insufficient to recommend for or against counseling of any intensity and/or behavioral interventions to promote sustained weight loss in overweight adults (I recommendation).

We undertook this systematic review to help update these recommendations. To conduct it, we developed an

See also:

**Print**

Editors' Notes . . . . . 435

**Web-Only**

- Appendix Table
- Appendix Figures
- CME quiz
- Conversion of graphics into slides

analytic framework with 4 key questions (**Appendix Figure 1**, available at [www.annals.org](http://www.annals.org)). The first was whether primary care screening programs to identify obesity or overweight in adults improved health or physiologic outcomes or resulted in weight loss. The other questions asked whether primary care feasible or referable weight-loss interventions (behaviorally based, with or without pharmacologic adjuncts) improved health outcomes, improved physiologic outcomes, resulted in short-term (12 to 18 months) or long-term (>18 months) weight loss, or caused harm.

## METHODS

The full report (9) describes our methods in detail.

### Data Sources and Searches

We relied on existing reviews to cover part of the search window from the previous USPSTF review, following previously detailed guidance (14). We identified a 2006 National Institute for Clinical Excellence systematic review on behavioral weight-loss interventions and orlistat (15) and a 2008 review of metformin trials (16). Their inclusion and exclusion criteria were congruent with ours, and investigators for both searched multiple databases and examined reference lists of pertinent reports. The reviews' search and selection strategies were judged acceptable to substitute for ours through 2005. We bridge-searched MEDLINE, PsycINFO, and the Cochrane Central Register of Controlled Trials from 2005 through 9 September 2010. We supplemented our search with relevant existing systematic reviews identified through databases (Cochrane Database of Systematic Reviews, the Database of Abstracts of Reviews of Effects, and MEDLINE) and Web sites (Institute of Medicine, National Institutes of Health, and National Institute for Health and Clinical Excellence). We supplemented our searches with experts' suggestions and reference lists from relevant publications, including evaluating all studies from the previous USPSTF review (17).

### Study Selection

Two investigators independently reviewed 6498 abstracts and 648 articles against prespecified inclusion and exclusion criteria (**Appendix Figure 2**, available at [www.annals.org](http://www.annals.org)). For key questions 1 to 3, we included randomized or controlled clinical trials with interventions focused on weight loss in adults (age  $\geq 18$  years) conducted in settings relevant to primary care (studies conducted in primary care or those that could in theory be implemented in a health care system, to which primary care clinicians could refer patients). We defined criteria for acceptable control groups a priori so that they would represent usual care and not overlap with low-intensity intervention groups. Acceptable control groups could not receive a personalized intervention, at-home workbook materials, or advice more frequently than annually; they also could not participate in frequent weigh-ins (<3 months). Healthy lifestyle messages were considered equivalent to weight-loss

### Context

Experts recommend that primary care clinicians offer obese adults interventions that promote weight loss.

### Contribution

In this systematic review of 58 trials, overweight adults in behavioral treatment trials that provided 12 to 26 intervention sessions during the first year lost 9 to 15 lb, whereas control groups lost little or no weight. Adults who received orlistat plus intensive behavioral interventions lost 11 to 22 lb, and those receiving placebo lost 7 to 13 lb.

### Caution

Behavioral treatment trials studied heterogeneous interventions, and orlistat trials had high rates of attrition.

### Implication

Behavior-oriented interventions can help overweight adults achieve meaningful weight loss.

—The Editors

messages. For harms (key question 4), we included additional study designs (large cohort studies or case-control studies; large event monitoring; systematic evidence reviews of randomized, controlled trials [RCTs] or controlled clinical trials) and did not require 12 months of follow-up.

### Data Extraction and Quality Assessment

Two independent investigators appraised all included articles as good, fair, or poor quality according to design-specific criteria and USPSTF methods (18). A third investigator resolved discrepancies. We assessed validity of randomization and measurement procedures, attrition, baseline characteristics, intervention fidelity, and statistical methods. Good-quality trials blinded researchers to participant randomization if they performed tasks related to assessment, had follow-up data on 90% or more of participants with fewer than a 10-percentage point difference between groups, and described anthropomorphic measurements in detail. Trials were rated poor quality and excluded if attrition was greater than 40%, was missing, or differed by more than 20% between groups (except for harms data); key baseline characteristics differed substantially between groups and were not controlled for in analyses; or outcomes were measured unequally between groups. Additional issues caused trials to be downgraded but not excluded; these included inconsistently applied interventions, selective reporting, and unclear or suboptimal blinding or randomization procedures (9). A table of excluded studies is available in our full report (9).

For included studies, one investigator abstracted data on study design, setting, population characteristics, baseline health and weight, intervention characteristics, prespecified outcomes, funding source, and adverse events

into standardized evidence tables (9). A second investigator reviewed abstraction for accuracy.

### Data Synthesis and Analysis

We conducted separate random-effects meta-analyses to estimate the effect size of behavioral and pharmacologic interventions on weight loss (expressed in kg) and intermediate health outcomes (adiposity, systolic and diastolic blood pressure, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglycerides, and glucose). Within each intervention type, trials were grouped according to the study population's risk status—cardiovascular risk (diabetes, dyslipidemia, hypertension), subclinical risk (prediabetes [19], borderline high lipids [20], prehypertension [21], abdominal obesity as defined by study researchers [22, 23]), and unselected/low risk—and then ordered by the behavioral intervention's intensity (number of sessions for behavioral trials and brief or intensive behavioral component accompanying medication trials).

We assessed the presence of statistical heterogeneity among studies by using standard chi-square tests and estimated heterogeneity magnitude by using the  $I^2$  statistic (24). Tests of publication bias included funnel plots and the Egger linear regression method (25) when there were 10 or more studies (26).

We explored heterogeneity of the effect size for weight loss with a series of meta-regressions (9). Factors we included were population risk status, recruitment strategy, retention, study focus (weight maintenance vs. loss), whether the trial was conducted in primary care, setting (United States or not), quality, and selected patient characteristics. For behavioral trials, we also examined the number of sessions during the first year and presence of several key intervention components (9). For medications, we also examined the percentage retained after run-in, medication type, and intensity of accompanying behavioral intervention.

All analyses were performed by using Stata 10.0 (Stata-Corp LP, College Station, Texas).

### Role of the Funding Source

We worked with 4 USPSTF liaisons at key points throughout the process to develop and refine the analytic framework, address methodological issues, and define scope. This research was funded by the Agency for Healthcare Research and Quality under a contract to support the work of the USPSTF. Agency for Healthcare Research and Quality staff provided project oversight, reviewed the draft report, and assisted in external draft report review.

## RESULTS

### Key Question 1: Screening for Obesity/Overweight

We identified no trials comparing screening with not screening for adult obesity.

### Key Questions 2 and 3: Benefits of Weight-Loss Interventions

We identified 58 trials of benefits of weight-loss interventions. Thirty-eight trials (13 495 participants) involved behaviorally based interventions (23, 27–64), 18 (11 256 participants) involved orlistat plus behavioral interventions (65–82), and 3 (2652 participants) involved metformin plus behavioral interventions (22, 27, 83). About one third of weight-loss trials could not be included in a weight-loss meta-analysis because of missing information, usually a measure of variability around the mean.


Behavioral trial participants had mean BMIs ranging from 25 to 39 kg/m<sup>2</sup>, with an average baseline BMI across all trials (weighted for sample size) of 31.9 kg/m<sup>2</sup>. Participants had mean ages of 34 to 70 years. Overall, 60% of participants were female; less than 40% of patients were nonwhite. Orlistat trial participants were 66% female, and less than 12% were nonwhite. Only 1 metformin trial reported ethnicity; 45.3% of patients were nonwhite (84). Fifty-five percent of behavioral trials and 57% of orlistat trials examined participants with clinical or subclinical cardiovascular risk factors. Metformin trials examined participants with diabetes risk factors (prediabetes or elevated waist-to-hip ratio).

### Weight Loss

Behavioral treatment trials were fairly high quality, with 24% being rated “good” (Appendix Table, available at [www.annals.org](http://www.annals.org)) (27, 28, 38, 53–56, 58, 61). Among those rated “fair,” allocation concealment and blinding of outcomes assessment were frequently unclear or not reported (59% and 83% of trials, respectively). Approximately 30% of fair-quality trials had follow-up of 90% or more at 12 months (Appendix Table) (23, 29, 32, 40, 42, 48, 57, 60). Just over one half of all trials limited analyses to completers (23, 27–30, 34, 39, 40, 43, 46, 47, 49–51, 53–55, 57, 58, 60, 62, 63), although 3 trials had low ( $\leq 3\%$ ) attrition (32, 58, 61). When data substitution was used, studies used baseline-observation-carried-forward (33, 37, 38, 42, 52), multiple imputation (35, 56), last-observation-carried-forward (37, 41, 44, 59), imputation of missing data through random-effects regression (35, 45, 64), multiple imputation method (35, 56), or unspecified methods (31, 48, 61). We did not find an association between effect size and study quality, attrition, or presence of data imputation through meta-regression.

Most trials showed that behavioral interventions had a statistically significant effect on weight loss at 12 to 18 months (23, 27–64). Controls generally lost little or no weight, whereas intervention groups lost 1.5 to 5 kg (3.3 to 11 lb), an average of 4% of baseline weight. In 21 trials that could be combined by meta-analysis, patients receiving behavioral interventions lost 3.0 kg (6.6 lb) more (95% CI,  $-4.0$  to  $-2.0$  kg) than controls after 12 to 18 months (Figure 1). Statistical heterogeneity was high ( $I^2 = 95\%$ )

Figure 1. Difference between intervention and control groups in weight change at 12 to 18 months.



Weights are from random-effects analysis. ADAPT = Activity, Diet and Blood Pressure Trial; CV = cardiovascular; DISH = Dietary Intervention to Study Hypertension; DPP = Diabetes Prevention Program; FDPS = Finnish Diabetes Prevention Study; HEED = Help Educate to Eliminate Diabetes; HI = intensive intervention; IV = intervention; LO = brief intervention; NR = not reported; ORBIT = Obesity Reduction Black Intervention Trial; PATH = Physical Activity for Total Health; PREDIAS = Prevention of Diabetes Self-Management Program; Subclinical = trials limited to those with elevated risk but without known disease (prehypertension; impaired glucose tolerance or elevated fasting glucose; borderline high total cholesterol, low-density lipoprotein, or triglyceride levels; low high-density lipoprotein levels; abdominal obesity); TOHP = Trials of Hypertension Prevention; WHLP = Women's Healthy Lifestyle Project; WMD = weighted mean difference.

because the amount of weight change varied greatly. Behavioral interventions lasting longer (24 to 54 months) continued to show greater weight loss (2 to 4 kg [4.4 to 8.8 lb]) compared with controls (28, 46, 53, 55, 58, 61). Weight loss could be maintained for an additional year or more after completion of an active weight-loss phase, particularly with additional support (27, 34, 39, 41, 52, 60).

Interventions with more sessions showed more weight loss—patients receiving 12 to 26 intervention sessions generally lost 4 to 7 kg (8.8 to 15.4 lb) (6% of baseline weight) compared with 1.5 to 4 kg (3.3 to 8.8 lb) (2.8% of baseline weight) in intervention groups with fewer than 12 sessions in the first year. After adjustment for number of sessions in the first year, none of the following demonstrated a relationship with effect size: physical activity sessions, group sessions, individual sessions, technology-based intervention, specific weight-loss goals, spouse or family involvement, addressing barriers to weight loss, motivational assessment (for example, pros and cons of weight loss), self-monitoring, incentives, or support after active intervention phase. However, our confidence in these null results is limited because some behavioral trials did not detail their interventions. These studies may have provided one or more components but not reported them. In addition, more intensive interventions tended to involve more components; disentangling the effect of intensity from specific components was not possible.

The orlistat data were limited in that there was only 1 good-quality trial (69) (**Appendix Table**). Randomization procedures (including allocation concealment) and medication adherence rates were rarely reported. Only 1 study specifically stated that funding was not from a pharmaceutical company (69). Only 5 studies had greater than 80% follow-up at 12 to 18 months (range, 61% to 96%) (69, 72, 74, 79, 82). Follow-up in control groups was often greater than 10% lower than in orlistat groups (72, 74, 79, 82). Over 70% of orlistat trials (65–67, 70–73, 77–82) and 33% of metformin trials (22) used last-observation-carried-forward for data substitution. The remaining analyzed only those with complete data (27, 68, 75, 83) or did not describe data substitution methods (69, 74, 76). We did not note an association between effect size and attrition or presence of data imputation through meta-regression, although power and range of attrition would be somewhat limited.

Orlistat treatment with accompanying behavioral component resulted in weight loss of 5 to 10 kg (11 to 22 lb; 8% of baseline weight) compared with 3 to 6 kg (7 to 13 lb; 5% of baseline weight) with placebo and the same behavioral component (65–82). Almost all orlistat trials used intensive behavioral components. In the 12 trials that could be combined by meta-analysis, participants randomly assigned to orlistat lost 3.0 kg (6.6 lb) more (95% CI, –3.9 to –2.0 kg) than those receiving placebo after 12 months (**Figure 1**). With 1 exception (69), the studies were

not highly variable. Limited data showed no dose response (72, 79). Weight loss was maintained with up to 24 to 36 months of orlistat therapy (78, 79). No trials reported weight outcomes after orlistat therapy was stopped. Metformin plus a behavioral intervention was associated with a smaller degree of weight loss (2 to 4 kg [4.4 to 8.8 lb]) (22, 83, 85), although the best evidence was limited to patients with prediabetes (85).

The effect of weight-loss programs among participant subgroups was sparsely reported and often mixed. Behaviorally based interventions seemed on average to lead to less weight loss in black patients and women than nonblack patients and men; effects of baseline BMI and age were mixed (30, 38, 43, 54, 56, 57, 61, 63, 86–88). Medication trials did not examine subgroups, or their findings applied only to patients with prediabetes (27, 87).

### Health Outcomes

Included trials did not demonstrate an effect on mortality, cardiovascular disease, hospitalizations, or depression, although data were sparse for all outcomes (**Table 1**) (22, 27, 42, 49, 56, 58, 63, 66, 72, 79, 81, 82, 84, 89–91). The 2 good-quality trials reporting 1 or more of these health outcomes were not powered to detect group differences in any health outcomes other than depressive symptoms (27, 58, 89).

### Diabetes Incidence

All intervention types reduced diabetes incidence, particularly in patients with elevated risk (**Table 1**). Behaviorally based interventions (7 to 23 sessions in first year), which led to weight loss of 4 to 7 kg (8.8 to 15.4 lb), cut diabetes incidence by about 50% or more over 2 to 3 years (58, 84). Metformin and orlistat reduced diabetes incidence (22, 78, 82, 84). However, orlistat data may not be reliable and generalizable (78, 82); by year 4, 1 trial had 48% and 68% attrition in the orlistat and placebo groups, respectively (82), and the other administered orlistat after participants experienced at least 5% weight loss during an 8-week very-low-calorie diet (78).

### Glucose Tolerance

Behaviorally based interventions, orlistat, and metformin all led to declines in fasting glucose levels in prediabetic and diabetic patients at 12 to 18 months compared with controls (31, 42, 45, 46, 49, 58, 69, 71, 74, 77, 85). Mean decreases in glucose levels were 0.30 and 0.31 mmol/L (5.4 and 5.5 mg/dL) with behavioral interventions and metformin, respectively. Glucose reductions were greater with orlistat (0.672 mmol/L [12.1 mg/dL] greater than placebo), possibly because those studies were conducted in diabetic patients.

**Table 1. Summary of Evidence for Effect of Weight-Loss Interventions on Clinical Outcomes**

Outcome	Trials, <i>n</i>	Overall Quality	Summary of Findings
<b>Distal health outcomes</b>			
Behaviorally based interventions			
Death	2	Good	No differences in death rate, but small number of deaths limits conclusions
CVD	4	Fair to good	No differences in CVD events or CVD-related deaths in 3 large good-quality trials; additional fair-quality trial showed no difference in percentage taking cardiovascular medication at 1 y
Hospitalization	1	Fair to good	No differences in hospitalization, but low hospitalization rate limits conclusions
HRQL/depression	3	Fair	None of 3 trials found group differences in depression outcomes; small change in HRQL correlated with weight change in the 1 good-quality trial (DPP)
Pharmacotherapy			
Orlistat			
Death	4	Fair	Each study only had 1 death; in all studies, deaths were in the orlistat group but there was no clear relationship with treatment
HRQL/depression	2	Fair	No difference in depression scores; orlistat group had greater satisfaction with treatment, less overweight distress, and improvement on 1 of 8 subscales (vitality) of SF-36
Metformin			
Death	2	Fair to good	No difference between groups, but small number of deaths limits conclusions
Hospitalization	1	Good	No difference in hospitalization, but low hospitalization rate limits conclusions
CVD	2	Fair to good	No difference in CVD events
HRQL/depression	1	Good	No difference in depression
<b>Intermediate health outcome</b>			
Behaviorally based interventions			
DM	3	Good	In 2 large good-quality trials, approximately twice as many participants in the control group than in the lifestyle intervention group developed DM, but no DM reduction was seen in the small trial, which had very high baseline rates of elevated fasting glucose levels
Pharmacotherapy			
Orlistat			
DM	2	Fair	Both trials reported lower incidence of diabetes (by 9–10 percentage points) in orlistat group, but we had concerns about the generalizability and reliability of these findings
Metformin			
DM	2	Fair to good	Incidence of diabetes was reduced with metformin in the good-quality trial in participants with prediabetes after 3 y (21.7% vs. 28.9%); the smaller trial with unclear adjudication also found decreased risk for diabetes in participants randomly assigned to metformin

CVD = cardiovascular disease; DM = type 2 diabetes mellitus; DPP = Diabetes Prevention Program; HRQL = health-related quality of life; SF-36 = Short Form-36.

### Lipids

Pooled estimates for lipid changes with behavioral interventions were at high risk for reporting bias because lipid outcomes were rarely reported (Figure 2). Trials included in meta-analyses were more likely to show effects than those not included. Although some trials did find statistically significant results, effect sizes were consistently small (most had reductions in low-density lipoprotein cholesterol level  $\leq 0.26$  mmol/L [10 mg/dL]). We concluded that behavioral weight-loss interventions had low or very small effects on low-density and high-density lipoprotein cholesterol and triglyceride levels (23, 29–31, 38, 41, 42, 46, 48, 49, 53, 57, 58, 62–64). Orlistat reduced low-density lipoprotein cholesterol levels by a slightly greater amount (0.29 mmol/L [11 mg/dL] more than placebo), but high-density lipoprotein cholesterol levels were reduced and triglyceride levels did not change (65–82). Metformin did not improve lipid profiles (22, 83, 90).

### Blood Pressure

Absolute reductions of 2 to 5 mm Hg in systolic and diastolic blood pressure were reported in behaviorally based and orlistat (plus behavioral intervention) trials over 12 to 36 months. When examined with meta-analyses, this translated into approximately 2-mm Hg greater decreases than control conditions after 12 to 18 months with either treatment (Figure 3) (23, 29–31, 42, 49, 53–55, 57, 58, 60, 63, 67, 68, 72, 77, 79–81, 90, 91). Behavioral treatment reduced the risk for a hypertension diagnosis in participants with prehypertension (34% and 22% reduced risk at 12 and 18 months, respectively) (54, 55). Metformin did not have favorable effects on blood pressure (22, 90).

### Waist Circumference

Waist circumference decreased by 2.7 cm more (CI,  $-4.1$  to  $-1.4$  cm) in behavioral intervention groups

Figure 2. Difference between intervention and control groups in changes in LDL cholesterol levels.

Study, Year (Reference)	Risk Status	IV Sessions	WMD (95% CI)	Number of Patients, Mean (SD) Difference in LDL Cholesterol Level (in mg/dL)	
				Treatment Group	Control Group
<b>Behavioral</b>					
Christian et al, 2008 (31)	CV risk	4	-10.81 (-19.95 to -1.67)	141, -14.6 (38.5)	132, -3.81 (38.5)
ter Bogt et al, 2009 (57)	CV risk	5	-0.39 (-5.29 to 4.51)	201, 2.32 (25.5)	215, 2.7 (25.5)
Mensink et al, 2003 (46)	Subclinical	4	-5.79 (-13.43 to 1.84)	40, 0.386 (19.7)	48, 6.18 (16.2)
Parikh et al, 2010 (Project HEED) (49)	Subclinical	8	-5.00 (-19.89 to 9.89)	35, -1 (35)	37, 4 (29)
Simkin-Silverman et al, 2003 (WHLP) (53)	Unselected/low	20	-6.90 (-10.86 to -2.94)	236, -4.2 (21.9)	253, 2.7 (22.8)
Wood et al, 1988 (62)	Unselected/low	23	-3.86 (-14.68 to 6.96)	42, -12 (24.7)	42, -8.11 (25.9)
Wood et al, 1991 (63)	Unselected/low	25	-6.18 (-13.03 to 0.67)	81, -10.8 (24.3)	79, -4.63 (19.7)
Irwin et al, 2003 (PATH) (38)	Unselected/low	128	-0.30 (-11.41 to 10.81)	87, -5.7 (35.1)	86, -5.4 (39.3)
Subtotal ( $I^2 = 0.0\%$ ; $P = 0.458$ )			-4.94 (-7.32 to -2.56)	863	892
<b>Orlistat</b>					
Berne et al, 2004 (65)	CV risk	HI	-3.47 (-13.22 to 6.27)	111, -3.09 (37.1)	109, 0.386 (36.7)
Lindgärde, 2000 (76)	CV risk	HI	-6.95 (-15.16 to 1.26)	190, -9.65 (43.2)	186, -2.7 (37.8)
Derosa et al, 2003 (68)	CV risk	HI	-16.00 (-26.94 to -5.06)	25, -37 (19)	23, -2.1 (19.6)
Hanefeld and Sachse, 2002 (71)	CV risk	HI	-7.10 (-13.39 to -0.81)	189, -2 (26.7)	180, 5.1 (34.3)
Miles et al, 2002 (77)	CV risk	HI	-7.72 (-14.15 to -1.29)	250, -9.65 (35)	254, -1.93 (38.6)
Derosa et al, 2010 (69)	CV risk	HI	-25.00 (-28.12 to -21.88)	113, -27 (12.7)	121, -2 (11.5)
Swinburn et al, 2005 (81)	CV risk	HI	-8.88 (-14.10 to -3.66)	170, -4.63 (25.1)	169, 4.25 (23.9)
Hollander et al, 1998 (74)	CV risk	NR	-13.51 (-19.45 to -7.57)	162, -5.02 (24.7)	159, 8.49 (29.3)
Rössner et al, 2000 (79)	Unselected/low	HI	-10.42 (-16.27 to -4.58)	242, -12.7 (30.2)	237, -2.32 (34.9)
Hauptman et al, 2000 (72)	Unselected/low	HI	-14.29 (-21.17 to -7.40)	210, -4.63 (38.5)	212, 9.65 (33.5)
Sjöström et al, 1998 (80)	Unselected/low	NR	-8.49 (-11.54 to -5.44)	343, -3.47 (20.5)	340, 5.02 (20.1)
Finer et al, 2000 (70)	Unselected/low	NR	-12.36 (-18.32 to -6.39)	110, -4.25 (24.3)	108, 8.11 (20.5)
Subtotal ( $I^2 = 86.3\%$ ; $P = 0.000$ )			-11.37 (-15.75 to -7.00)	2115	2098
<b>Metformin</b>					
Fontbonne et al, 1996 (22)	Subclinical	LO	-4.63 (-10.65 to 1.38)	164, -0.772 (29)	160, 3.86 (26.3)
Gambineri et al, 2006 (83)	Subclinical	HI	-6.00 (-25.43 to 13.43)	20, -14 (33.8)	19, -8 (28)
Subtotal ( $I^2 = 0.0\%$ ; $P = 0.895$ )			-4.75 (-10.50 to 0.99)	184	179
Overall ( $I^2 = 83.2\%$ ; $P = 0.000$ )			-8.73 (-12.00 to -5.46)	3162	3169

Weights are from random-effects analysis. Pooled estimates for lipid changes with behavioral interventions were at high risk for reporting bias because lipid outcomes were rarely reported. To convert LDL cholesterol values to mmol/L, multiply by 0.0259. CV = cardiovascular; HEED = Help Educate to Eliminate Diabetes; HI = intensive intervention; IV = intervention; LDL = low-density lipoprotein; LO = brief intervention; NR = not reported; PATH = Physical Activity for Total Health; Subclinical = trials limited to those with elevated risk but without known disease (prehypertension; impaired glucose tolerance or elevated fasting glucose; borderline high total cholesterol, low-density lipoprotein, or triglyceride levels; low high-density lipoprotein levels; abdominal obesity); WHLP = Women's Healthy Lifestyle Project; WMD = weighted mean difference.

than in control groups. Statistical heterogeneity was high ( $I^2 = 93.8\%$ ), but most trials showed statistically significant group differences (23, 30, 31, 37, 38, 41, 42, 46, 48, 49, 57, 58, 60, 85). Orlistat and metformin reduced waist circumference by 2.3 cm (CI, -3.6 to -0.9 cm) and 1.5 cm (CI, -2.0 to -1.0 cm), respectively, compared with placebo (65, 66, 68, 69, 71, 74-76, 78, 79, 81-83, 85).

**Key Question 4: Harms of Weight-Loss Interventions**  
**Behavioral Studies**

Ten studies reported on possible harms of behavioral weight-loss interventions. Weight loss reduced total (61) or

hip (53, 59) bone mineral density in 3 fair- to good-quality trials (53, 59, 61). Increased physical activity did not result in serious adverse effects or injuries over 1- to 2-year interventions (38, 46, 92, 93). One study reported no increased risk for eating disorder pathology in those participating in weight-loss interventions (94).

**Orlistat**


We included 18 RCTs from key questions 2 and 3 (65-82), 5 additional published RCTs not included in key questions 2 and 3 (95-99) (12 174 participants in all trials combined), and 1 United Kingdom event-

monitoring study (100) (16 021 persons) on the harms of orlistat (120 mg 3 times per day). Sixty-two percent of trials recruited participants with at least 1 clinical or subclinical cardiovascular risk factor (65, 66, 68, 69, 71, 74, 76–78, 81, 82, 95–98). Participants were 41 to 59 years of age; 66% of patients were female, and less than 15% were nonwhite. Although median trial duration was 52 weeks, 5 trials provided data beyond 52 weeks.

Participants randomly assigned to receive orlistat were more likely to experience adverse effects and withdraw

from trials because of adverse effects than those assigned to placebo (Table 2) (65–82, 95–99, 101). Withdrawals were primarily due to gastrointestinal symptoms. In included studies, serious adverse effects were not increased with orlistat compared with placebo (66, 68, 69, 75, 76, 78–82, 95, 96, 99). Orlistat was associated with a decrease in some fat-soluble vitamin levels compared with placebo (73, 74, 79, 82, 99). Data did not suggest that higher dosages were associated with elevated adverse effect rates, although results were mixed (72, 73, 79, 99).

Figure 3. Difference between intervention and control groups in changes in systolic blood pressure.



Weights are from random-effects analysis. ADAPT = Activity, Diet and Blood Pressure Trial; CV = cardiovascular; DPP = Diabetes Prevention Program; FDPS = Finnish Diabetes Prevention Study; HEED = Help Educate to Eliminate Diabetes; HI = intensive intervention; IV = intervention; LO = brief intervention; NR = not reported; ODES = Oslo Diet and Exercise Study; PATH = Physical Activity for Total Health; PREDIAS = Prevention of Diabetes Self-Management Program; Subclinical = trials limited to those with elevated risk but without known disease (prehypertension; impaired glucose tolerance or elevated fasting glucose; borderline high total cholesterol, low-density lipoprotein, or triglyceride levels; low high-density lipoprotein levels; abdominal obesity); TOHP = Trials of Hypertension Prevention; WHLP = Women's Healthy Lifestyle Project; WMD = weighted mean difference.

**Table 2. Summary of Medication Harms**

Adverse Events	Trials in Meta-analysis, Additional Trials, <i>n</i> , <i>n</i>	Meta-analysis Results: Relative Risk (95% CI)	Weighted Means, %	Results From Studies Not in Meta-analysis	Comments
<b>Orlistat</b>					
Withdrawals due to adverse effects	23, 0	1.67 (1.32–2.13)	IG: 8 CG: 4	–	GI symptoms were main reason for withdrawal
Any	8, 0	1.10 (1.03–1.17)	IG: 78 CG: 70	–	GI symptoms were main reason for AEs
Serious	11, 2	1.21 (0.88–1.68)	IG: 10 CG: 9	0 serious AEs in either treatment group in 2 trials	Examples of serious events included fecal incontinence, diverticulitis, abdominal pain
GI	18, 0	1.42 (1.33–1.52)	IG: 83 CG: 59	–	GI symptoms were of mild to moderate intensity and often resolved spontaneously
Hypoglycemia	0, 3	–	–	2 of 3 found increased incidence of hypoglycemia with orlistat	–
Bone mineral density	0, 1	–	–	In small subsample ( <i>n</i> = 30) of larger study, bone density did not differ between group	–
Vitamin deficiencies	0, 5	–	–	5 of 5 studies found lower vitamin E with orlistat; 4 of 4 studies found lower β-carotene levels with orlistat; 1 of 2 trials found lower vitamin A levels; 1 of 1 study found lower vitamin K levels; 5 of 5 studies found that orlistat recipients required more vitamin supplementation during the study	–
Liver injury	0, 1 (event monitoring cohort)	–	–	UK monitoring study reported elevated liver test results in 2 cases of 16 021 dispenses analyzed; no cases of serious hepatic adverse reactions	FDA recently added warning to label to orlistat about possible risk for severe liver disease
<b>Metformin</b>					
Withdrawals	2, 0	3.92 (1.23–12.57)	IG: 5 CG: 1	–	–
Any	2, 0	4.83 (0.84–27.63)	IG: 46 CG: 16	–	–
Serious	0	–	–	–	–
GI	1, 3	–	–	Increased risk of GI AEs in metformin group	Main GI symptoms included diarrhea, flatulence, nausea, vomiting
Hypoglycemia	0	–	–	–	–
Bone density	0	–	–	–	–

AE = adverse event; CG = control group; FDA = U.S. Food and Drug Administration; GI = gastrointestinal; IG = intervention group; UK = United Kingdom.

**Metformin**

We included 4 trials on harms of metformin (850 mg twice daily): 3 trials from key questions 2 and 3 (22, 83, 86) and 1 additionally published RCT (102) (2712 participants). Participants randomly assigned to receive metformin were more likely than placebo recipients to have and withdraw because of adverse events (Table 2) (22, 83, 102). Gastrointestinal symptoms were the main reason for excess adverse effects (22, 83, 86, 102). No studies reported serious adverse effects.

**DISCUSSION**

Because we found no trials directly examining the benefits and harms of obesity screening followed by expected appropriate treatment in adults, we focused on effectiveness and harms of primary care–relevant weight-loss inter-

ventions. Behavioral treatment trials were fairly recent and of high quality. However, orlistat trials were generally lower quality. Participants in behavioral treatment trials providing 12 to 26 intervention sessions during the first year lost 4 to 7 kg (8.8 to 15.4 lb) (average, 6% of baseline weight) at 12 to 18 months, compared with little to no weight loss in control groups. Participants receiving orlistat plus intensive behaviorally based intervention lost 5 to 10 kg (11 to 22 lb) (average, 8% of baseline weight), compared with 3 to 6 kg in the placebo groups. Metformin was associated with less weight loss (2 to 4 kg [4.4 to 8.8 lb]). Long-term weight loss (>18 months) was sparsely reported, but weight loss generally persisted with continued treatment. Five percent weight loss is considered clinically meaningful and is a primary weight-loss outcome according to the U.S. Food and Drug Administration (103).

Most (104–107), but not all (108), epidemiologic data suggest that intentional weight loss less than 9 kg is not associated with reduced mortality. Epidemiologic data, however, are mixed and confounded by several factors, particularly health status. Prospective cohort studies of patients undergoing bariatric surgery show substantial improvements in health. Weight loss with surgery, however, is generally 25 to 50 kg (55 to 110 lb) (109, 110).

Because health outcome data were insufficient, we examined the metabolic consequences of weight-loss interventions. Two fair- to good-quality trials showed that diabetes incidence was reduced by 30% to 50% with behavioral weight-loss interventions among overweight and obese patients with elevated plasma glucose levels. Behavioral weight-loss interventions had little to no effect on lipids. Improved cholesterol levels may require large amounts of weight loss (111). Lowering low-density lipoprotein cholesterol levels with orlistat may reflect reduced fat absorption (112). Although summary measures showed small blood pressure reductions, absolute reductions of 2 to 5 mm Hg were reported in some orlistat and behaviorally based trials over 12 to 36 months, consistent with findings of a previous meta-analysis (113). Reductions in diastolic blood pressure of 5 to 6 mm Hg over 5 to 10 years have been associated with small reductions in stroke and coronary heart disease events (114).

Higher treatment intensity was associated with greater weight loss, despite limitations in our measure of treatment intensity. Most higher-intensity interventions included self-monitoring, setting goals, addressing barriers to change, and strategizing about maintaining long-term changes. However, we found that no component was associated with degree of weight loss in meta-regression. Specific articulation of essential elements of effective interventions was not possible.

Methods used to measure obesity in clinical practice (for example, BMI and waist circumference) are low cost and cause no direct physical harms. Secondary harms could include labeling stigma, higher insurance premiums, or reinforcement of poor self-esteem. Misclassifying a person's risk status is possible if current BMI cut-offs are used during screening because BMI may predict future health risk differently among various ethnic and age groups (115, 116).

Weight loss may be associated with decreased bone density, but data are lacking on post-weight-loss bone density and subsequent fracture risk. The validity of measuring bone changes during weight loss is also unclear. Fat distribution changes may alter bone measurement despite stable bone (117–119). Limited data suggested no increased risk for serious injuries or eating disorders.

Orlistat and metformin caused mild to moderate gastrointestinal side effects resulting in discontinuation of therapy. Our inclusion and exclusion criteria were not designed to identify rare harms. We did not find studies

associating orlistat with liver, kidney, or pancreas damage, other than case reports. However, in May 2010, the U.S. Food and Drug Administration revised its label for orlistat, 120 mg (prescription) and 60 mg (over-the-counter), to include “new safety information about cases of severe liver injury that have been reported rarely with the use of this medication” (120). Orlistat has been recently associated with possible kidney and pancreas damage (121). Indeed, antiobesity medications have a long history of removal from the market or failing U.S. Food and Drug Administration approval (122–124).

Several factors limited this evidence review. Because of missing information, we could not include about one third of trials in meta-analyses. Intermediate physiologic outcomes were even less likely to be studied and available for meta-analyses. We excluded 143 studies because control groups had more than minimal interventions (considered comparative effectiveness), including Look AHEAD, a weight-loss intervention trial in diabetic patients (125). Its findings were similar to and slightly more positive than the findings of included trials. Comparative-effectiveness trials would shed more light on effective intervention components and are being reviewed elsewhere (126).

Few studies were conducted in primary care. Interventions were often intensive and possibly difficult to implement in primary care settings (although providers could refer to them). No trial had a mean BMI in the class III obese range ( $>40 \text{ kg/m}^2$ ), so generalizability to extreme obesity is unknown. Because most medication trials had run-in periods, participants were probably more motivated, adherent, and responsive than primary care populations. Our results, especially medication findings, are possibly biased by high attrition.

We did not systematically examine the best screening approach for obesity. A growing body of evidence suggests that waist-to-hip ratio or waist circumference may be better predictors of future health effects than is BMI (127–135), especially for some subgroups (116, 136). Systematically rereviewing the best screening tool for adult obesity should be of high priority. We did not systematically review cost-effectiveness data. The economic impact of weight-loss interventions is an important clinical consideration.

Metformin is the only off-label medication we included. Other medications used off-label for weight loss include zonisamide, an antiepileptic agent (137). We did not examine antiobesity drugs in development, including lorcaserin; a combination of phentermine plus topiramate; or a combination naltrexone plus bupropion. We did not include phentermine because it is approved only for short-term use.

In summary, we found no direct evidence on benefits and harms of primary care-based obesity screening but did find that behavioral weight-loss interventions with or without orlistat or metformin yielded clinically meaningful weight loss; however, health outcomes data were sparse.

Harms of behavioral weight-loss interventions were minimal; data were insufficient on serious harms from medications. Long-term weight and health outcomes data were lacking and should be a high priority for future study. Research should clarify which benefits are derived specifically from weight loss itself or from behavioral mediators, such as physical activity or dietary changes.

From Center for Health Research, Kaiser Permanente Northwest, Portland, Oregon.

**Disclaimer:** Agency for Healthcare Research and Quality staff provided oversight for the project and assisted in external review of the draft-evidence synthesis. USPSTF liaisons helped to resolve issues around the scope of the review but were not involved in the conduct of the review.

**Acknowledgment:** The authors thank Victor Stevens, PhD, MA, for providing his expertise; Daphne Plaut, MLS, for conducting the literature searches; Kevin Lutz, MFA, for his editorial support; Caitlyn Senger, MPH, Tracy Beil, MS, Catherine Livingston, MD, MPH, and Sarah Zuber, MSW, for their assistance in conducting the evidence review; and Leslie Perdue, MPH, for her assistance in conducting the evidence review and preparing the manuscript. They also thank the Agency for Healthcare Research and Quality and the USPSTF, as well as expert reviewers David Arterburn, MD, MPH; George Bray, MD; James Hill, PhD; Robert Jeffery, PhD; and Jarl Torgerson, MD, PhD, for their contribution to this evidence review.

**Grant Support:** By the Oregon Evidence-based Practice Center under contract to the Agency for Healthcare Research and Quality, Rockville, Maryland (contract HHS-290-2007-10057-I, task order no. 3).

**Potential Conflicts of Interest:** Dr. LeBlanc: *Grant (money to institution):* Agency for Healthcare Research and Quality. Dr. O'Connor: *Grant (money to institution):* Agency for Healthcare Research and Quality; *Support for travel to meetings for the study or other purposes (money to institution):* Agency for Healthcare Research and Quality; *Payment for writing or reviewing the manuscript (money to institution):* Agency for Healthcare Research and Quality; *Provision of writing assistance, medicines, equipment, or administrative support (money to institution):* Agency for Healthcare Research and Quality. Dr. Kapka: *Grant (money to institution):* Agency for Healthcare Research and Quality; *Support for travel to meetings for the study or other purposes (money to institution):* Agency for Healthcare Research and Quality; *Payment for writing or reviewing the manuscript (money to institution):* Agency for Healthcare Research and Quality; *Provision of writing assistance, medicines, equipment, or administrative support (money to institution):* Agency for Healthcare Research and Quality. Disclosures can also be viewed at [www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M11-0823](http://www.acponline.org/authors/icmje/ConflictOfInterestForms.do?msNum=M11-0823).

**Requests for Single Reprints:** Reprints are available from the Agency for Healthcare Research and Quality Web site ([www.preventiveservices.ahrq.gov](http://www.preventiveservices.ahrq.gov)).

Current author addresses and author contributions are available at [www.annals.org](http://www.annals.org).

## References

1. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA*. 2010;303:235-41. [PMID: 20071471]

2. Stein CJ, Colditz GA. The epidemic of obesity. *J Clin Endocrinol Metab*. 2004;89:2522-5. [PMID: 15181019]
3. Adams KF, Schatzkin A, Harris TB, Kipnis V, Mouw T, Ballard-Barbash R, et al. Overweight, obesity, and mortality in a large prospective cohort of persons 50 to 71 years old. *N Engl J Med*. 2006;355:763-78. [PMID: 16926275]
4. Flegal KM, Graubard BI, Williamson DF, Gail MH. Excess deaths associated with underweight, overweight, and obesity. *JAMA*. 2005;293:1861-7. [PMID: 15840860]
5. Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al; **Prospective Studies Collaboration**. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *Lancet*. 2009;373:1083-96. [PMID: 19299006]
6. Bogers RP, Bemelmans WJ, Hoogenveen RT, Boshuizen HC, Woodward M, Knekt P, et al; **for the BMI-CHD Collaboration Investigators**. Association of overweight with increased risk of coronary heart disease partly independent of blood pressure and cholesterol levels: a meta-analysis of 21 cohort studies including more than 300 000 persons. *Arch Intern Med*. 2007;167:1720-8. [PMID: 17846390]
7. Guh DP, Zhang W, Bansback N, Amarsi Z, Birmingham CL, Anis AH. The incidence of co-morbidities related to obesity and overweight: a systematic review and meta-analysis. *BMC Public Health*. 2009;9:88. [PMID: 19320986]
8. Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet*. 2008;371:569-78. [PMID: 18280327]
9. LeBlanc E, O'Connor E, Whitlock EP, Patnode CD, Kapka T. Screening and management of obesity and overweight in adults. Rockville, MD: Agency for Healthcare Research and Quality; 2011. Accessed at [www.uspreventiveservicestaskforce.org/uspstf/uspsobes.htm](http://www.uspreventiveservicestaskforce.org/uspstf/uspsobes.htm) on 14 August 2011.
10. Calle EE, Thun MJ, Petrelli JM, Rodriguez C, Heath CW Jr. Body-mass index and mortality in a prospective cohort of U.S. adults. *N Engl J Med*. 1999;341:1097-105. [PMID: 10511607]
11. Lenz M, Richter T, Mühlhauser I. The morbidity and mortality associated with overweight and obesity in adulthood: a systematic review. *Dtsch Arztebl Int*. 2009;106:641-8. [PMID: 19890430]
12. Pischon T, Boeing H, Hoffmann K, Bergmann M, Schulze MB, Overvad K, et al. General and abdominal adiposity and risk of death in Europe. *N Engl J Med*. 2008;359:2105-20. [PMID: 19005195]
13. Sarwer DB, Allison KC, Gibbons LM, Markowitz JT, Nelson DB. Pregnancy and obesity: a review and agenda for future research. *J Womens Health (Larchmt)*. 2006;15:720-33. [PMID: 16910904]
14. Cornell JE, Laine C. The science and art of deduction: complex systematic overviews [Editorial]. *Ann Intern Med*. 2008;148:786-8. [PMID: 18490692]
15. **National Institute for Health and Clinical Excellence**. Obesity: guidance on the prevention, identification, assessment and management of overweight and obesity in adults and children. Clinical Guideline 43, 1-84. 2006. Accessed at [www.nice.org.uk/nicemedia/live/11000/30365/30365.pdf](http://www.nice.org.uk/nicemedia/live/11000/30365/30365.pdf) on 9 August 2011.
16. Desilets AR, Dhakal-Karki S, Dunican KC. Role of metformin for weight management in patients without type 2 diabetes. *Ann Pharmacother*. 2008;42:817-26. [PMID: 18477733]
17. McTigue KM, Harris R, Hemphill MB, Bunton AJ, Lux LJ, Sutton S, et al. Screening and interventions for overweight and obesity in adults. Contract no. 290-97-0011, task no. 3. Systematic evidence review no. 21. Rockville, MD: Agency for Healthcare Research and Quality; 2003.
18. Harris RP, Helfand M, Woolf SH, Lohr KN, Mulrow CD, Teutsch SM, et al; **Methods Work Group, Third US Preventive Services Task Force**. Current methods of the US Preventive Services Task Force: a review of the process. *Am J Prev Med*. 2001;20:21-35. [PMID: 11306229]
19. **American Diabetes Association**. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2004;27 Suppl 1:S5-S10. [PMID: 14693921]
20. **National Cholesterol Education Program, National Heart, Lung and Blood Institute, and National Institutes of Health**. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). NIH publication no. 02-5215. 2002. Accessed at [www.nhlbi.nih.gov/guidelines/cholesterol/atp3full.pdf](http://www.nhlbi.nih.gov/guidelines/cholesterol/atp3full.pdf) on 31 May 2011.
21. **U.S. Department of Health and Human Services, National Institutes of Health, and National Heart, Lung and Blood Institute**. Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. NIH publication no. 04-5230. 2004. Accessed at [www.nhlbi.nih.gov/guidelines/hypertension/jnc7full.pdf](http://www.nhlbi.nih.gov/guidelines/hypertension/jnc7full.pdf) on 31 May 2011.

22. Fontbonne A, Charles MA, Juhan-Vague I, Bard JM, André P, Isnard F, et al. The effect of metformin on the metabolic abnormalities associated with upper-body fat distribution. BIGPRO Study Group. *Diabetes Care*. 1996;19:920-6. [PMID: 8875083]
23. Mitsui T, Shimaoka K, Tsuzuku S, Kajioka T, Sakakibara H. Gentle exercise of 40 minutes with dietary counseling is effective in treating metabolic syndrome. *Tohoku J Exp Med*. 2008;215:355-61. [PMID: 18679010]
24. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21:1539-58. [PMID: 12111919]
25. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315:629-34. [PMID: 9310563]
26. Terrin N, Schmid CH, Lau J. In an empirical evaluation of the funnel plot, researchers could not visually identify publication bias. *J Clin Epidemiol*. 2005;58:894-901. [PMID: 16085192]
27. **The Diabetes Prevention Program**. Design and methods for a clinical trial in the prevention of type 2 diabetes. *Diabetes Care*. 1999;22:623-34. [PMID: 10189543]
28. The Hypertension Prevention Trial: three-year effects of dietary changes on blood pressure. Hypertension Prevention Trial Research Group. *Arch Intern Med*. 1990;150:153-62. [PMID: 2404477]
29. Andersen S, Holme I, Urdal P, Hjermann I. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press*. 1995;4:343-9. [PMID: 8746601]
30. Burke V, Beilin LJ, Cutt HE, Mansour J, Wilson A, Mori TA. Effects of a lifestyle programme on ambulatory blood pressure and drug dosage in treated hypertensive patients: a randomized controlled trial. *J Hypertens*. 2005;23:1241-9. [PMID: 15894901]
31. Christian JG, Bessesen DH, Byers TE, Christian KK, Goldstein MG, Bock BC. Clinic-based support to help overweight patients with type 2 diabetes increase physical activity and lose weight. *Arch Intern Med*. 2008;168:141-6. [PMID: 18227359]
32. Cohen MD, D'Amico FJ, Merenstein JH. Weight reduction in obese hypertensive patients. *Fam Med*. 1991;23:25-8. [PMID: 2001777]
33. Cussler EC, Teixeira PJ, Goings SB, Houtkooper LB, Metcalfe LL, Blew RM, et al. Maintenance of weight loss in overweight middle-aged women through the Internet. *Obesity (Silver Spring)*. 2008;16:1052-60. [PMID: 18309301]
34. Davis BR, Oberman A, Blaufox MD, Wassertheil-Smoller S, Hawkins CM, Cutler JA, et al. Effect of antihypertensive therapy on weight loss. The Trial of Antihypertensive Interventions and Management Research Group. *Hypertension*. 1992;19:393-9. [PMID: 1555871]
35. Fitzgibbon ML, Stolley MR, Schiffer L, Sharp LK, Singh V, Dyer A. Obesity Reduction Black Intervention Trial (ORBIT): 18-month results. *Obesity (Silver Spring)*. 2010;18:2317-25. [PMID: 20300081]
36. Frey-Hewitt B, Vranizan KM, Dreon DM, Wood PD. The effect of weight loss by dieting or exercise on resting metabolic rate in overweight men. *Int J Obes*. 1990;14:327-34. [PMID: 2361810]
37. Haapala I, Barengo NC, Biggs S, Surakka L, Manninen P. Weight loss by mobile phone: a 1-year effectiveness study. *Public Health Nutr*. 2009;12:2382-91. [PMID: 19323865]
38. Irwin ML, Yasui Y, Ulrich CM, Bowen D, Rudolph RE, Schwartz RS, et al. Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial. *JAMA*. 2003;289:323-30. [PMID: 12525233]
39. Jeffery RW, Wing RR, Thorson C, Burton LR, Raether C, Harvey J, et al. Strengthening behavioral interventions for weight loss: a randomized trial of food provision and monetary incentives. *J Consult Clin Psychol*. 1993;61:1038-45. [PMID: 8113481]
40. Jones DW, Miller ME, Wofford MR, Anderson DC Jr, Cameron ME, Willoughby DL, et al. The effect of weight loss intervention on antihypertensive medication requirements in the Hypertension Optimal Treatment (HOT) study. *Am J Hypertens*. 1999;12:1175-80. [PMID: 10619579]
41. Kastarinen MJ, Puska PM, Korhonen MH, Mustonen JN, Salomaa VV, Sundvall JE, et al; LIHEF Study Group. Non-pharmacological treatment of hypertension in primary health care: a 2-year open randomized controlled trial of lifestyle intervention against hypertension in eastern Finland. *J Hypertens*. 2002;20:2505-12. [PMID: 12473876]
42. Kulzer B, Hermanns N, Gorges D, Schwarz P, Haak T. Prevention of diabetes self-management program (PREDIAS): effects on weight, metabolic risk factors, and behavioral outcomes. *Diabetes Care*. 2009;32:1143-6. [PMID: 19509014]
43. Langford HG, Blaufox MD, Oberman A, Hawkins CM, Curb JD, Cutter GR, et al. Dietary therapy slows the return of hypertension after stopping prolonged medication. *JAMA*. 1985;253:657-64. [PMID: 3881608]
44. Martin PD, Dutton GR, Rhode PC, Horswell RL, Ryan DH, Brantley PJ. Weight loss maintenance following a primary care intervention for low-income minority women. *Obesity (Silver Spring)*. 2008;16:2462-7. [PMID: 18787526]
45. Mayer-Davis EJ, D'Antonio AM, Smith SM, Kirkner G, Levin Martin S, Parra-Medina D, et al. Pounds off with empowerment (POWER): a clinical trial of weight management strategies for black and white adults with diabetes who live in medically underserved rural communities. *Am J Public Health*. 2004;94:1736-42. [PMID: 15451743]
46. Mensink M, Blaak EE, Corpeleijn E, Saris WH, de Bruin TW, Feskens EJ. Lifestyle intervention according to general recommendations improves glucose tolerance. *Obes Res*. 2003;11:1588-96. [PMID: 14694225]
47. Moore H, Summerbell CD, Greenwood DC, Tovey P, Griffiths J, Henderson M, et al. Improving management of obesity in primary care: cluster randomised trial. *BMJ*. 2003;327:1085. [PMID: 14604931]
48. Narayan KM, Hoskin M, Kozak D, Kriska AM, Hanson RL, Pettitt DJ, et al. Randomized clinical trial of lifestyle interventions in Pima Indians: a pilot study. *Diabet Med*. 1998;15:66-72. [PMID: 9472866]
49. Parikh P, Simon EP, Fei K, Looker H, Goytia C, Horowitz CR. Results of a pilot diabetes prevention intervention in East Harlem, New York City: Project HEED. *Am J Public Health*. 2010;100 Suppl 1:S232-9. [PMID: 20147680]
50. Perri MG, McAllister DA, Gange JJ, Jordan RC, McAdoo G, Nezu AM. Effects of four maintenance programs on the long-term management of obesity. *J Consult Clin Psychol*. 1988;56:529-34. [PMID: 2848874]
51. Pritchard DA, Hyndman J, Taba F. Nutritional counselling in general practice: a cost effective analysis. *J Epidemiol Community Health*. 1999;53:311-6. [PMID: 10396539]
52. Silva MN, Vieira PN, Coutinho SR, Minderico CS, Matos MG, Sardinha LB, et al. Using self-determination theory to promote physical activity and weight control: a randomized controlled trial in women. *J Behav Med*. 2010;33:110-22. [PMID: 20012179]
53. Simkin-Silverman LR, Wing RR, Boraz MA, Kuller LH. Lifestyle intervention can prevent weight gain during menopause: results from a 5-year randomized clinical trial. *Ann Behav Med*. 2003;26:212-20. [PMID: 14644697]
54. Stevens VJ, Corrigan SA, Obarzanek E, Bernauer E, Cook NR, Hebert P, et al. Weight loss intervention in phase 1 of the Trials of Hypertension Prevention. The TOHP Collaborative Research Group. *Arch Intern Med*. 1993;153:849-58. [PMID: 8466377]
55. Stevens VJ, Obarzanek E, Cook NR, Lee IM, Appel LJ, Smith West D, et al; Trials for the Hypertension Prevention Research Group. Long-term weight loss and changes in blood pressure: results of the Trials of Hypertension Prevention, phase II. *Ann Intern Med*. 2001;134:1-11. [PMID: 11187414]
56. Svetkey LP, Stevens VJ, Brantley PJ, Appel LJ, Hollis JF, Loria CM, et al; Weight Loss Maintenance Collaborative Research Group. Comparison of strategies for sustaining weight loss: the weight loss maintenance randomized controlled trial. *JAMA*. 2008;299:1139-48. [PMID: 18334689]
57. ter Bogt NC, Bemelmans WJ, Beltman FW, Broer J, Smit AJ, van der Meer K. Preventing weight gain: one-year results of a randomized lifestyle intervention. *Am J Prev Med*. 2009;37:270-7. [PMID: 19765497]
58. Tuomilehto J, Lindström J, Eriksson JG, Valle TT, Hämäläinen H, Ilanne-Parikka P, et al; Finnish Diabetes Prevention Study Group. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med*. 2001;344:1343-50. [PMID: 11333990]
59. Villareal DT, Shah K, Banks MR, Sinacore DR, Klein S. Effect of weight loss and exercise therapy on bone metabolism and mass in obese older adults: a one-year randomized controlled trial. *J Clin Endocrinol Metab*. 2008;93:2181-7. [PMID: 18364384]
60. Werkman A, Hulshof PJ, Stafleu A, Kremers SP, Kok FJ, Schouten EG, et al. Effect of an individually tailored one-year energy balance programme on body weight, body composition and lifestyle in recent retirees: a cluster randomised controlled trial. *BMC Public Health*. 2010;10:110. [PMID: 20205704]
61. Whelton PK, Appel LJ, Espeland MA, Applegate WB, Ettinger WH Jr, Kostis JB, et al. Sodium reduction and weight loss in the treatment of hypertension in older persons: a randomized controlled trial of nonpharmacologic interventions in the elderly (TONE). TONE Collaborative Research Group. *JAMA*. 1998;279:839-46. [PMID: 9515998]
62. Wood PD, Stefanick ML, Dreon DM, Frey-Hewitt B, Garay SC, Williams


- PT, et al. Changes in plasma lipids and lipoproteins in overweight men during weight loss through dieting as compared with exercise. *N Engl J Med.* 1988;319:1173-9. [PMID: 3173455]
63. Wood PD, Stefanick ML, Williams PT, Haskell WL. The effects on plasma lipoproteins of a prudent weight-reducing diet, with or without exercise, in overweight men and women. *N Engl J Med.* 1991;325:461-6. [PMID: 1852180]
64. Woollard J, Burke V, Beilin LJ, Verheijden M, Bulsara MK. Effects of a general practice-based intervention on diet, body mass index and blood lipids in patients at cardiovascular risk. *J Cardiovasc Risk.* 2003;10:31-40. [PMID: 12569235]
65. Berne C, Orlistat Swedish Type 2 diabetes Study Group. A randomized study of orlistat in combination with a weight management programme in obese patients with type 2 diabetes treated with metformin. *Diabet Med.* 2005;22:612-8. [PMID: 15842517]
66. Broom I, Wilding J, Stott P, Myers N; UK Multimorbidity Study Group. Randomised trial of the effect of orlistat on body weight and cardiovascular disease risk profile in obese patients: UK Multimorbidity Study. *Int J Clin Pract.* 2002;56:494-9. [PMID: 12296610]
67. Davidson MH, Hauptman J, DiGirolamo M, Foreyt JP, Halsted CH, Heber D, et al. Weight control and risk factor reduction in obese subjects treated for 2 years with orlistat: a randomized controlled trial. *JAMA.* 1999;281:235-42. [PMID: 9918478]
68. Derosa G, Mugellini A, Ciccarelli L, Fogari R. Randomized, double-blind, placebo-controlled comparison of the action of orlistat, fluvastatin, or both an anthropometric measurements, blood pressure, and lipid profile in obese patients with hypercholesterolemia prescribed a standardized diet. *Clin Ther.* 2003;25:1107-22. [PMID: 12809960]
69. Derosa G, Maffioli P, Salvadeo SA, Ferrari I, Gravina A, Mereu R, et al. Comparison of orlistat treatment and placebo in obese type 2 diabetic patients. *Expert Opin Pharmacother.* 2010;11:1971-82. [PMID: 20569086]
70. Finer N, James WP, Kopelman PG, Lean ME, Williams G. One-year treatment of obesity: a randomized, double-blind, placebo-controlled, multicentre study of orlistat, a gastrointestinal lipase inhibitor. *Int J Obes Relat Metab Disord.* 2000;24:306-13. [PMID: 10757623]
71. Hanefeld M, Sachse G. The effects of orlistat on body weight and glycaemic control in overweight patients with type 2 diabetes: a randomized, placebo-controlled trial. *Diabetes Obes Metab.* 2002;4:415-23. [PMID: 12406041]
72. Hauptman J, Lucas C, Boldrin MN, Collins H, Segal KR. Orlistat in the long-term treatment of obesity in primary care settings. *Arch Fam Med.* 2000;9:160-7. [PMID: 10693734]
73. Hill JO, Hauptman J, Anderson JW, Fujioka K, O'Neil PM, Smith DK, et al. Orlistat, a lipase inhibitor, for weight maintenance after conventional dieting: a 1-y study. *Am J Clin Nutr.* 1999;69:1108-16. [PMID: 10357727]
74. Hollander PA, Elbein SC, Hirsch IB, Kelley D, McGill J, Taylor T, et al. Role of orlistat in the treatment of obese patients with type 2 diabetes. A 1-year randomized double-blind study. *Diabetes Care.* 1998;21:1288-94. [PMID: 9702435]
75. Krempf M, Louvet JP, Allanic H, Miloradovich T, Joubert JM, Attali JR. Weight reduction and long-term maintenance after 18 months treatment with orlistat for obesity. *Int J Obes Relat Metab Disord.* 2003;27:591-7. [PMID: 12704403]
76. Lindgärde F. The effect of orlistat on body weight and coronary heart disease risk profile in obese patients: the Swedish Multimorbidity Study. *J Intern Med.* 2000;248:245-54. [PMID: 10971792]
77. Miles JM, Leiter L, Hollander P, Wadden T, Anderson JW, Doyle M, et al. Effect of orlistat in overweight and obese patients with type 2 diabetes treated with metformin. *Diabetes Care.* 2002;25:1123-8. [PMID: 12087008]
78. Richelsen B, Tonstad S, Rössner S, Toubro S, Niskanen L, Madsbad S, et al. Effect of orlistat on weight regain and cardiovascular risk factors following a very-low-energy diet in abdominally obese patients: a 3-year randomized, placebo-controlled study. *Diabetes Care.* 2007;30:27-32. [PMID: 17192328]
79. Rössner S, Sjöström L, Noack R, Meinders AE, Noseda G. Weight loss, weight maintenance, and improved cardiovascular risk factors after 2 years treatment with orlistat for obesity. European Orlistat Obesity Study Group. *Obes Res.* 2000;8:49-61. [PMID: 10678259]
80. Sjöström L, Rissanen A, Andersen T, Boldrin M, Golay A, Koppeschaar HP, et al. Randomised placebo-controlled trial of orlistat for weight loss and prevention of weight regain in obese patients. European Multicentre Orlistat Study Group. *Lancet.* 1998;352:167-72. [PMID: 9683204]
81. Swinburn BA, Carey D, Hills AP, Hooper M, Marks S, Proietto J, et al. Effect of orlistat on cardiovascular disease risk in obese adults. *Diabetes Obes Metab.* 2005;7:254-62. [PMID: 15811142]
82. Torgerson JS, Hauptman J, Boldrin MN, Sjöström L. XENical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care.* 2004;27:155-61. [PMID: 14693982]
83. Gambineri A, Patton L, Vaccina A, Cacciari M, Morselli-Labate AM, Cavazza C, et al. Treatment with flutamide, metformin, and their combination added to a hypocaloric diet in overweight-obese women with polycystic ovary syndrome: a randomized, 12-month, placebo-controlled study. *J Clin Endocrinol Metab.* 2006;91:3970-80. [PMID: 16868063]
84. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med.* 2002;346:393-403. [PMID: 11832527]
85. Haffner S, Temprosa M, Crandall J, Fowler S, Goldberg R, Horton E, et al; Diabetes Prevention Program Research Group. Intensive lifestyle intervention or metformin on inflammation and coagulation in participants with impaired glucose tolerance. *Diabetes.* 2005;54:1566-72. [PMID: 15855347]
86. Crandall J, Schade D, Ma Y, Fujimoto WY, Barrett CE, et al; Diabetes Prevention Program Research Group. The influence of age on the effects of lifestyle modification and metformin in prevention of diabetes. *J Gerontol A Biol Sci Med Sci.* 2006;61:1075-81. [PMID: 17077202]
87. West DS, Elaine Prewitt T, Bursac Z, Felix HC. Weight loss of black, white, and Hispanic men and women in the Diabetes Prevention Program. *Obesity (Silver Spring).* 2008;16:1413-20. [PMID: 18421273]
88. Stevens J, Juhaeri, Cai J, Jones DW. The effect of decision rules on the choice of a body mass index cutoff for obesity: examples from African American and white women. *Am J Clin Nutr.* 2002;75:986-92. [PMID: 12036803]
89. Rubin RR, Knowler WC, Ma Y, Marrero DG, Edelstein SL, Walker EA, et al; Diabetes Prevention Program Research Group. Depression symptoms and antidepressant medicine use in Diabetes Prevention Program participants. *Diabetes Care.* 2005;28:830-7. [PMID: 15793181]
90. Ratner R, Goldberg R, Haffner S, Marcovina S, Orchard T, Fowler S, et al; Diabetes Prevention Program Research Group. Impact of intensive lifestyle and metformin therapy on cardiovascular disease risk factors in the diabetes prevention program. *Diabetes Care.* 2005;28:888-94. [PMID: 15793191]
91. Lindegarde F. The effect of orlistat on body weight and coronary heart disease risk profile in obese patients: the Swedish Multimorbidity Study. *J Intern Med.* 2000;248:245-54. [PMID: 10971792]
92. Kirk EP, Jacobsen DJ, Gibson C, Hill JO, Donnelly JE. Time course for changes in aerobic capacity and body composition in overweight men and women in response to long-term exercise: the Midwest Exercise Trial (MET). *Int J Obes Relat Metab Disord.* 2003;27:912-9. [PMID: 12861232]
93. Warren M, Schmitz KH. Safety of strength training in premenopausal women: musculoskeletal injuries from a two-year randomized trial. *Am J Health Promot.* 2009;23:309-14. [PMID: 19445432]
94. Williamson DA, Martin CK, Anton SD, York-Crowe E, Han H, Redman L, et al; Pennington CALERIE Team. Is caloric restriction associated with development of eating-disorder symptoms? Results from the CALERIE trial. *Health Psychol.* 2008;27:S32-42. [PMID: 18248104]
95. Bakris G, Calhoun D, Egan B, Hellmann C, Dolker M, Kingma I, et al; orlistat and resistant hypertension investigators. Orlistat improves blood pressure control in obese subjects with treated but inadequately controlled hypertension. *J Hypertens.* 2002;20:2257-67. [PMID: 12409965]
96. Broom I, Hughes E, Dodson P, Reckless J. The role of orlistat in the treatment of obese patients with mild to moderate hypercholesterolaemia: consequences for coronary risk. *Br J Cardiol.* 2002;9:460-68.
97. Kelley DE, Bray GA, Pi-Sunyer FX, Klein S, Hill J, Miles J, et al. Clinical efficacy of orlistat therapy in overweight and obese patients with insulin-treated type 2 diabetes: A 1-year randomized controlled trial. *Diabetes Care.* 2002;25:1033-41. [PMID: 12032111]
98. Muls E, Kolanowski J, Scheen A, Van Gaal L; ObelHyx Study Group. The effects of orlistat on weight and on serum lipids in obese patients with hypercholesterolemia: a randomized, double-blind, placebo-controlled, multicentre study. *Int J Obes Relat Metab Disord.* 2001;25:1713-21. [PMID: 11753595]
99. Van Gaal LF, Broom JI, Enzi G, Toplak H. Efficacy and tolerability of orlistat in the treatment of obesity: a 6-month dose-ranging study. Orlistat Dose-Ranging Study Group. *Eur J Clin Pharmacol.* 1998;54:125-32. [PMID: 9626916]

100. Acharya NV, Wilton LV, Shakir SA. Safety profile of orlistat: results of a prescription-event monitoring study. *Int J Obes (Lond)*. 2006;30:1645-52. [PMID: 16552401]
101. Gokcel A, Gumurdulu Y, Karakose H, Melek Ertorer E, Tanaci N, BascilTutuncu N, et al. Evaluation of the safety and efficacy of sibutramine, orlistat and metformin in the treatment of obesity. *Diabetes Obes Metab*. 2002; 4:49-55. [PMID: 11874442]
102. Trolle B, Flyvbjerg A, Kesmodel U, Lauszus FF. Efficacy of metformin in obese and non-obese women with polycystic ovary syndrome: a randomized, double-blinded, placebo-controlled cross-over trial. *Hum Reprod*. 2007;22:2967-73. [PMID: 17766923]
103. U.S. Food and Drug Administration. **Guidance for industry**. Developing products for weight management. Accessed at [www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm071612.pdf](http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm071612.pdf) on 31 May 2011.
104. Gregg EW, Gerzoff RB, Thompson TJ, Williamson DF. Trying to lose weight, losing weight, and 9-year mortality in overweight U.S. adults with diabetes. *Diabetes Care*. 2004;27:657-62. [PMID: 14988281]
105. Knudtson MD, Klein BE, Klein R, Shankar A. Associations with weight loss and subsequent mortality risk. *Ann Epidemiol*. 2005;15:483-91. [PMID: 16029840]
106. Williamson DF, Pamuk E, Thun M, Flanders D, Byers T, Heath C. Prospective study of intentional weight loss and mortality in never-smoking overweight US white women aged 40-64 years. *Am J Epidemiol*. 1995;141:1128-41. [PMID: 7771451]
107. Yaari S, Goldbourt U. Voluntary and involuntary weight loss: associations with long term mortality in 9,228 middle-aged and elderly men. *Am J Epidemiol*. 1998;148:546-55. [PMID: 9753009]
108. Williamson DF, Thompson TJ, Thun M, Flanders D, Pamuk E, Byers T. Intentional weight loss and mortality among overweight individuals with diabetes. *Diabetes Care*. 2000;23:1499-504. [PMID: 11023143]
109. Christou NV, Sampalis JS, Liberman M, Look D, Auger S, McLean AP, et al. Surgery decreases long-term mortality, morbidity, and health care use in morbidly obese patients. *Ann Surg*. 2004;240:416-23. [PMID: 15319713]
110. Peeters A, O'Brien PE, Laurie C, Anderson M, Wolfe R, Flum D, et al. Substantial intentional weight loss and mortality in the severely obese. *Ann Surg*. 2007;246:1028-33. [PMID: 18043106]
111. Sjöström CD, Lissner L, Sjöström L. Relationships between changes in body composition and changes in cardiovascular risk factors: the SOS Intervention Study. *Swedish Obese Subjects*. *Obes Res*. 1997;5:519-30. [PMID: 9449135]
112. Tonstad S, Pometta D, Erkelens DW, Ose L, Moccetti T, Schouten JA, et al. The effect of the gastrointestinal lipase inhibitor, orlistat, on serum lipids and lipoproteins in patients with primary hyperlipidaemia. *Eur J Clin Pharmacol*. 1994;46:405-10. [PMID: 7957533]
113. Neter JE, Stam BE, Kok FJ, Grobbee DE, Geleijnse JM. Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension*. 2003;42:878-84. [PMID: 12975389]
114. Collins R, MacMahon S. Blood pressure, antihypertensive drug treatment and the risks of stroke and of coronary heart disease. *Br Med Bull*. 1994;50:272-98. [PMID: 8205459]
115. Razak F, Anand SS, Shannon H, Vuksan V, Davis B, Jacobs R, et al. Defining obesity cut points in a multiethnic population. *Circulation*. 2007;115: 2111-8. [PMID: 17420343]
116. Sánchez-Castillo CP, Velázquez-Monroy O, Berber A, Lara-Esqueda A, Tapia-Conyer R, James WP; Encuesta Nacional de Salud (ENSA) 2000 Working Group. Anthropometric cutoff points for predicting chronic diseases in the Mexican National Health Survey 2000. *Obes Res*. 2003;11:442-51. [PMID: 12634443]
117. Tothill P. Dual-energy x-ray absorptiometry measurements of total-body bone mineral during weight change. *J Clin Densitom*. 2005;8:31-8. [PMID: 15722585]
118. Tothill P, Laskey MA, Orphanidou CI, van Wijk M. Anomalies in dual energy X-ray absorptiometry measurements of total-body bone mineral during weight change using Lunar, Hologic and Norland instruments. *Br J Radiol*. 1999;72:661-9. [PMID: 10624323]
119. Tothill P, Hannan WJ, Cowen S, Freeman CP. Anomalies in the measurement of changes in total-body bone mineral by dual-energy X-ray absorptiometry during weight change. *J Bone Miner Res*. 1997;12:1908-21. [PMID: 9383696]
120. U.S. Food and Drug Administration. Questions and answers: orlistat and severe liver injury. *FDA Drug Bulletin*. May 26, 2010. Accessed at [www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm213040.htm](http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm213040.htm) on 31 May 2011.
121. Weir MA, Beyea MM, Gomes T, Juurlink DN, Mamdani M, Blake PG, et al. Orlistat and acute kidney injury: an analysis of 953 patients [Letter]. *Arch Intern Med*. 2011;171:703-4. [PMID: 21482850]
122. Finer N; Executive Steering Committee of the Sibutramine Cardiovascular Outcome Trial. Withdrawal of sibutramine. Editorial in advance of the facts [Letter]. *BMJ*. 2010;340:c1346. [PMID: 20219799]
123. Healy M. Weight-loss drug Qnexa gets thumbs-down from FDA panel. *Los Angeles Times*. July 15, 2010. Accessed at <http://articles.latimes.com/2010/jul/15/news/la-heb-qnexa-20100715> on 31 May 2011.
124. U.S. Food and Drug Administration. Meridia (sibutramine): market withdrawal due to risk for serious cardiovascular events. October 8, 2010. Accessed at [www.fda.gov/safety/medwatch/safetyinformation/safetyalertsforhumanmedicalproducts/ucm228830.htm](http://www.fda.gov/safety/medwatch/safetyinformation/safetyalertsforhumanmedicalproducts/ucm228830.htm) on 31 May 2011.
125. Wing RR; Look AHEAD Research Group. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med*. 2010;170:1566-75. [PMID: 20876408]
126. Evidence-based Practice Center Systematic Review Protocol. Project title: Comparative effectiveness of bariatric surgery and non-surgical therapy in adults with metabolic conditions and body mass index of 30 to 34.9. Accessed at [www.effectivehealthcare.ahrq.gov/ehc/products/227/595/Bariatric%20protocol%20111010.pdf](http://www.effectivehealthcare.ahrq.gov/ehc/products/227/595/Bariatric%20protocol%20111010.pdf) on 31 May 2011.
127. Arderm CI, Janssen I, Ross R, Katzmarzyk PT. Development of health-related waist circumference thresholds within BMI categories. *Obes Res*. 2004; 12:1094-103. [PMID: 15292473]
128. Freemantle N, Holmes J, Hockey A, Kumar S. How strong is the association between abdominal obesity and the incidence of type 2 diabetes? *Int J Clin Pract*. 2008;62:1391-6. [PMID: 18557792]
129. Koster A, Leitzmann MF, Schatzkin A, Mouw T, Adams KF, van Eijk JT, et al. Waist circumference and mortality. *Am J Epidemiol*. 2008;167:1465-75. [PMID: 18417494]
130. Lear SA, Kohli S, Bondy GP, Tchernof A, Sniderman AD. Ethnic variation in fat and lean body mass and the association with insulin resistance. *J Clin Endocrinol Metab*. 2009;94:4696-702. [PMID: 19820012]
131. Lee CM, Huxley RR, Wildman RP, Woodward M. Indices of abdominal obesity are better discriminators of cardiovascular risk factors than BMI: a meta-analysis. *J Clin Epidemiol*. 2008;61:646-53. [PMID: 18359190]
132. Patel MR, Donahue M, Wilson PW, Califf RM. Clinical trial issues in weight-loss therapy. *Am Heart J*. 2006;151:633-42. [PMID: 16504625]
133. Qiao Q, Nyamadorj R. The optimal cutoff values and their performance of waist circumference and waist-to-hip ratio for diagnosing type II diabetes. *Eur J Clin Nutr*. 2010;64:23-9. [PMID: 19690576]
134. Vazquez G, Duval S, Jacobs DR Jr, Silventoinen K. Comparison of body mass index, waist circumference, and waist/hip ratio in predicting incident diabetes: a meta-analysis. *Epidemiol Rev*. 2007;29:115-28. [PMID: 17494056]
135. Janssen I, Katzmarzyk PT, Ross R. Waist circumference and not body mass index explains obesity-related health risk. *Am J Clin Nutr*. 2004;79:379-84. [PMID: 14985210]
136. Lear SA, James PT, Ko GT, Kumanyika S. Appropriateness of waist circumference and waist-to-hip ratio cutoffs for different ethnic groups. *Eur J Clin Nutr*. 2010;64:42-61. [PMID: 19672278]
137. Vetter ML, Faulconbridge LF, Webb VL, Wadden TA. Behavioral and pharmacologic therapies for obesity. *Nat Rev Endocrinol*. 2010;6:578-88. [PMID: 20680034]

**Current Author Addresses:** Drs. LeBlanc, O'Connor, Whitlock, Patnode, and Kapka: Center for Health Research, Kaiser Permanente Northwest, 3800 North Interstate Avenue, Portland, OR 97227.


**Author Contributions:** Conception and design: E.S. LeBlanc, E. O'Connor, E. Whitlock, C.D. Patnode.  
 Analysis and interpretation of the data: E.S. LeBlanc, E. O'Connor, E. Whitlock, T. Kapka.  
 Drafting of the article: E.S. LeBlanc, E. Whitlock, C.D. Patnode, T. Kapka.  
 Critical revision of the article for important intellectual content: E.S. LeBlanc, E. O'Connor, E. Whitlock, C.D. Patnode, T. Kapka.  
 Final approval of the article: E.S. LeBlanc, E. O'Connor, E. Whitlock, C.D. Patnode, T. Kapka.  
 Statistical expertise: E. O'Connor.  
 Obtaining of funding: E. Whitlock.  
 Collection and assembly of data: E.S. LeBlanc, E. O'Connor, T. Kapka.

**Appendix Figure 1. Analytic framework: primary care screening and interventions for obesity and overweight.**



KQ 1: Is there direct evidence that primary care screening programs for adult obesity or overweight improve health outcomes or result in short-term (12 to 18 mo) or sustained (>18 mo) weight loss or improved physiologic measures (i.e., glucose tolerance, blood pressure, and dyslipidemia)? a) How well is weight loss maintained after an intervention is completed? KQ 2: Do primary care-relevant interventions (behaviorally based interventions and/or pharmacotherapy) in obese or overweight adults lead to improved health outcomes (morbidity from diabetes mellitus, cardiovascular disease, cancer, arthritis, asthma, sleep apnea, depression, emotional functioning, physical fitness capacity or performance, physical functioning, disability, mortality)? a) What are common elements of efficacious interventions? b) Are there differences in efficacy between patient subgroups (i.e., age 65 y or older, sex, race/ethnicity, degree of obesity, baseline cardiovascular risk)? KQ 3: Do primary care-relevant interventions in obese or overweight adults lead to short-term or sustained weight loss, with or without improved physiologic measures? a) How well is weight loss maintained after an intervention is completed? b) What are common elements of efficacious interventions? c) Are there differences in efficacy between patient subgroups (i.e., age 65 y or older, sex, race/ethnicity, degree of obesity, baseline cardiovascular risk)? KQ 4: What are the adverse effects of primary care-relevant interventions in obese or overweight adults (e.g., nutritional deficits, cardiovascular disease, bone mass loss, injuries, death)? a) Are there differences in adverse effects between patient subgroups (i.e., age 65 y or older, sex, race/ethnicity, degree of obesity, baseline cardiovascular risk status)? KQ = key question.

Appendix Figure 2. Literature search and selection.



KQ = key question.

**Appendix Table. Outcomes Reported and Quality Issues for Included Trials**

Study, Year (Reference)	Outcomes								Allocation Concealment	Quality Issues			
	Weight Loss	Distal Health Outcome	DM	Glucose Tolerance	Lipids	Blood Pressure	Waist Circumference	Adverse Events		BOA	Retention*	ITT Analysis	Overall Quality and Other Quality Concerns or Clarifications
<b>Behavioral trials</b>													
With CV risk factor													
Diabetes													
Christian et al, 2008 (31)	XX		X	XX	XX	XX			Yes	NR	88%	Unclear	Fair Use of the term "ITT" not clear
Mayer-Davis et al, 2004 (45)	X		X	X	X				NR	NR	81%	RER	Fair
Hypertension													
Burke et al, 2005 (ADAPT) (30)	XX					XX	XX		Yes	NR	80%	No	Fair
Cohen et al, 1991 (32)	XX					X			NR	NR	100%	NA	Fair Small number of participants
Davis et al, 1992 (TAIM) (34)	X	X				X			Yes	Yes (blood pressure) NR (weight loss)	Unclear	No	Fair Retention NR at 12 mo, 89% at 6 mo, 59% at 24 mo
Jones et al, 1999 (HOT) (40)	X					X			Unclear	NR	91%	No	Fair Unclear information on outcome blinding (described as "single blind"), assessment and statistical methods not well described
Kastarinen et al, 2002 (LIHEF) (41)	X		X	X	X	X	X		Yes	NR	83%	LOCF	Fair Retention lower in control than intervention group at 12 mo (77% vs. 88%)
Langford et al, 1985 (DISH) (43)	XX								NR	NR	82%	No	Fair Attrition lower in control than intervention group (77% vs. 87%) Minimal description of control group contact
Whelton et al, 1998 (TONE) (61)	X	X				X		X	NR	Yes	98%	NR	Good
Multiple risk factors													
Anderssen et al, 1995 (ODES) (29)	X	X			X	XX			Yes	NR	95%	No	Fair Lack of information about assessment blinding, but well-defined procedures probably minimized bias
Svetkey et al, 2008 (WLM) (56)	X	X							Yes	Yes	95%	MImp	Good
ter Bogt et al, 2009 (57)	XX			XX	XX	XX	XX		NR	NR	91%	No	Fair Lack of information about assessment blinding, but well-defined procedures probably minimized bias
Woollard et al, 2003 (64)	XX				X				NR	No (weight loss), yes (lipid)	71%	RER	Fair Outcome measurement by nurse giving intervention Details of outcome measures NR
Subclinical													
Parikh et al, 2010 (Project HEED) (49)	XX		X	XX	XX	XX	XX		NR	NR	73%	No	Fair Small number of participants
HPT, 1990 (28)	X					X			Yes	Likely	91%	No	Good

Continued on following page

## Appendix Table—Continued

Study, Year (Reference)	Outcomes								Allocation Concealment	Quality Issues			
	Weight Loss	Distal Health Outcome	DM	Glucose Tolerance	Lipids	Blood Pressure	Waist Circumference	Adverse Events		BOA	Retention*	ITT Analysis	Overall Quality and Other Quality Concerns or Clarifications
DPP, 2005 (85)	XX	X	X	XX		XX	XX	X	Yes	Partial	95%	No	Good Outcomes assessment seems unblinded but required initial certification and annual recertification of assessment staff, which should minimize if not eliminate bias introduced by lack of blinding
Stevens et al, 1993 (TOHP I) (54)	XX					XX			Yes	Yes	94%	No	Good
Stevens et al, 2001 (TOHP II) (55)	XX					XX			Yes	Yes	92%	No	Good
Villareal et al, 2008 (59)	X	X		X	X	X		X	Yes	Yes	89%	LOCF	Fair
Mensink et al, 2003 (46)	XX			XX	XX		XX	X	Unclear	NR	81%	No	Fair
Tuomilehto et al, 2001 (58)	XX	X	X	XX	XX	XX	XX		Unclear	NR	97%	No	Good Uncertain if allocation concealed; person scheduling baseline appointments was blinded to randomization list, but there is uncertainty whether this was the person who conducted the assessment
Kulzer et al, 2009 (42)	XX	X		XX	XX	XX	XX		NR	NR	91%	BOCF	Fair Group-specific follow-up NR
Mitsui et al, 2008 (23)	X			XX	XX	XX	XX		NR	NR	94%	No	Fair Small number of participants higher proportion of control group had the metabolic syndrome than intervention group (32% vs. 21%)
Low risk or unselected Cussler et al, 2008 (33)	X								NR	NR	82%	BOCF	Fair Did not adjust analysis for cluster randomization
Fitzgibbon et al, 2010 (ORBIT) (35)	XX								Yes	No	89%	MImp	Fair Assessment not blinded, but well-defined procedures probably minimized bias
Haapala et al, 2009 (37)	XX							XX	Yes	NR	68%	BOCF or LOCF (value that was highest)	Fair
Irwin et al, 2003 (PATH) (38)	XX			X	XX		XX	X	Yes	Yes	98%	BOCF	Good
Jeffery et al, 1993 (39)	X								NR	NR	87%	No	Fair Measurement procedures not well described, attrition by group NR
Martin et al, 2008 (44)	XX								NR	NR	65%	LOCF	Fair Group-specific attrition not provided at 12 mo, retention was lower in intervention group than control group at 6-mo (71% vs. 88%) and 18-mo (54% vs. 77%) follow-up
Moore et al, 2003 (47)	X								Yes	Yes	67%	No	Fair
Narayan et al, 1998 (48)	X		X	X	X	X	X		NR	NR	93%	NR	Fair BMI, weight, waist circumference, and fasting glucose levels higher in intervention group at baseline; measurement procedures not clearly described.

Continued on following page

Appendix Table—Continued

Study, Year (Reference)	Outcomes								Allocation Concealment	Quality Issues			Overall Quality and Other Quality Concerns or Clarifications
	Weight Loss	Distal Health Outcome	DM	Glucose Tolerance	Lipids	Blood Pressure	Waist Circumference	Adverse Events		BOA	Retention*	ITT Analysis	
Perri et al, 1988 (50)	X								NR	NR	74%	No	Fair Little information about baseline comparability
Pritchard et al, 1999 (51)	X							Likely	NR	66%	No	Fair	Lower retention in dietitian group than other 2 groups (54.5% vs. 71% and 71%); outcomes reported only for overweight/obesity, hypertension, and DM subgroups, although randomization was not stratified by disease status; statistical procedures not well-described
Silva et al, 2010 (52)	X								NR	NR	81%	BOCF	Fair Lower retention in control group than intervention group (80% vs. 93%); lack of information about blinding of assessment and blinding, but well-defined procedures probably minimized bias
Simkin-Silverman et al, 2003 (WHLP) (53)	XX			XX	XX	XX			Yes	Yes	94%	No	Good
Werkman et al, 2010 (60)	XX						XX	XX	No	NR	95%	No	Fair Did not have blinded outcomes assessment, but well-defined procedures probably minimized bias; analyzed only male participants (so not truly randomized comparison)
Wood et al, 1991 (63)	XX	X			XX	XX			NR	NR	88%	No	Fair
Wood et al, 1988 (62)	XX				XX				Yes	NR	85%	No	Fair Retention higher in exercise than other groups (90% vs. 81% and 82%)
<b>Medication trials</b>													
<b>Orlistat</b>													
Berne et al, 2005 (65)	X			X	XX	X	XX	XX	NR	Yes	86%	LOCF	Fair Possible selective reporting of weight outcomes
Broom et al, 2002 (66)	XX			X	X	X	X	XX	NR	Yes	65%	LOCF	Fair Statistical procedures not well described
Davidson et al, 1999 (67)	XX			X	X	XX		XX	NR	Yes	66%	LOCF	Fair High dropout during run-in
Derosa et al, 2003 (68)	XX				XX	XX	XX	XX	Yes	Yes	96%	No	Fair Small sample measurement methods not clearly described

Continued on following page

## Appendix Table—Continued

Study, Year (Reference)	Outcomes								Allocation Concealment	Quality Issues			
	Weight Loss	Distal Health Outcome	DM	Glucose Tolerance	Lipids	Blood Pressure	Waist Circumference	Adverse Events		BOA	Retention*	ITT Analysis	Overall Quality and Other Quality Concerns or Clarifications
Derosa et al, 2010 (69)	XX			XX	XX		XX	XX	Yes	Yes	92%	NR	Good
Finer et al, 2000 (70)	X				XX			XX	Yes	Yes	61%	LOCF	Fair
Hanefeld and Sachse, 2002 (71)	XX			XX	XX	X	XX	XX	NR	Yes	69%	LOCF	Fair Measurement procedures not well described
Hauptman et al, 2000 (72)	XX	X		XX	XX	XX		XX	NR	Yes	67%	LOCF	Fair
Hill et al, 1999 (73)	X			X	X	X		XX	NR	Yes	74%	LOCF	Fair Measurement procedures not well described Did not report whether placebo matched active medicine
Hollander et al, 1998 (74)	XX			XX	XX		XX	XX	NR	Yes	79%	Unclear	Fair
Krempf et al, 2003 (75)	XX				X		XX	XX	NR	Yes	61%	No	Fair
Lindgärde, 2000 (76)	XX			XX	XX	XX	X	XX	NR	Yes	86%	NR	Fair
Miles et al, 2002 (77)	XX			XX	XX	XX		XX	NR	Yes	60%	LOCF	Fair Did not report whether placebo matched active medicine
Richelsen et al, 2007 (78)	X		X	X	X	X	X	XX	Yes	Yes	65%	LOCF	Fair
Rössner et al, 2000 (79)	XX	X		XX	XX	XX	X	XX	NR	Yes	72%	LOCF	Fair
Sjöström et al, 1998 (80)	X			XX	XX	XX		XX	Likely	Yes	79%	LOCF	Fair
Swinburn et al, 2005 (81)	XX	X		XX	XX	XX	XX	XX	NR	Yes	79%	LOCF	Fair Did not report whether placebo matched active medicine
Torgerson et al, 2004 (XENDOS) (82)	X		X	X	X	X	X	XX	Yes	Yes	83%	LOCF	Fair Higher retention in control than intervention group (90 vs. 78)
Metformin													
Fontbonne et al, 1996 (BIGPRO) (22)	XX	X	X	XX	XX	XX		XX	NR	Yes	71%	LOCF	Fair
Gambineri et al, 2006 (83)	XX			XX	XX		XX	XX	NR	NR	98%	No	Fair Not double blind—only participants blinded
DPP, 2005 (85)	XX	X	X	XX	X	XX	XX	X	Yes	Partial	95%	No	Good Nonlaboratory outcomes assessment seems unblinded but required initial certification and annual recertification of assessment staff, which should minimize if not eliminate bias introduced by lack of blinding

ADAPT = Activity, Diet and Blood Pressure Trial; BIGPRO = BIGuanides and Prevention of the Risks in Obesity; BMI = body mass index; BOA = blinding of outcomes assessment; BOCF = baseline-observation-carried-forward method; CV = cardiovascular; DM = diabetes mellitus; DISH = Dietary Intervention to Study Hypertension; DPP = Diabetes Prevention Program; HEED = Help Educate to Eliminate Diabetes; HOT = Hypertension Optimal Treatment; HPT = Hypertension Prevention Trial; ITT = intention-to-treat; LIHEF = Lifestyle Intervention against Hypertension in Eastern Finland; LOCF = last-observation-carried-forward method; MImp = multiple imputation method; NA = not applicable; NR = not reported; ODES = Oslo Diet and Exercise Study; ORBIT = Obesity Reduction Black Intervention Trial; PATH = Physical Activity for Total Health; RER = imputation of missing data through use of random-effects regression; TAIM = Trial of Antihypertensive Interventions and Management; TOHP = Trials of Hypertension Prevention; TONE = Trial of Nonpharmacologic Interventions in the Elderly; WHLP = Women's Healthy Lifestyle Project; WLM = Weight Loss Maintenance; X = outcome was reported; XENDOS = XENical in the Prevention of Diabetes in Obese Subjects; XX = outcome was included in the meta-analysis.

\* Proportion with follow-up data.