

Screening for Syphilis Infection in Pregnant Women

US Preventive Services Task Force Reaffirmation Recommendation Statement

US Preventive Services Task Force

IMPORTANCE Untreated syphilis infection in pregnant women can be transmitted to the fetus (congenital syphilis) at any time during pregnancy or at birth. Congenital syphilis is associated with stillbirth, neonatal death, and significant morbidity in infants (eg, bone deformities and neurologic impairment). After a steady decline from 2008 to 2012, cases of congenital syphilis markedly increased from 2012 to 2106, from 8.4 to 15.7 cases per 100 000 live births (an increase of 87%). At the same time, national rates of syphilis increased among women of reproductive age.

OBJECTIVE To update the US Preventive Services Task Force (USPSTF) 2009 recommendation on screening for syphilis infection in pregnant women.

EVIDENCE REVIEW The USPSTF commissioned a reaffirmation evidence update to identify new and substantial evidence sufficient enough to change its prior recommendation. Given the established benefits and practice of screening for syphilis in pregnant women, the USPSTF targeted its evidence review on the direct benefits of screening on the prevention of congenital syphilis morbidity and mortality and the harms of screening for and treatment of syphilis infection in pregnant women.

FINDINGS Using a reaffirmation process, the USPSTF found that accurate screening algorithms are available to identify syphilis infection. Effective treatment with antibiotics can prevent congenital syphilis and significantly decrease adverse pregnancy outcomes, with small associated harms, providing an overall substantial health benefit. Therefore, the USPSTF reaffirms its previous conclusion that there is convincing evidence that screening for syphilis infection in pregnant women provides substantial benefit.

CONCLUSIONS AND RECOMMENDATION The USPSTF recommends early screening for syphilis infection in all pregnant women. (A recommendation)

JAMA. 2018;320(9):911-917. doi:10.1001/jama.2018.11785

← Related article page 918 and JAMA Patient Page page 948

+ CME Quiz at jamanetwork.com/learning

+ Related article at jamadermatology.com

Group Information: The US Preventive Services Task Force members are listed at the end of this article.

Corresponding Author: Susan J. Curry, PhD, The University of Iowa, 111 Jessup Hall, Iowa City, IA 52242 (chair@uspstf.net).

The US Preventive Services Task Force (USPSTF) makes recommendations about the effectiveness of specific preventive care services for patients without obvious related signs or symptoms.

It bases its recommendations on the evidence of both the benefits and harms of the service and an assessment of the balance. The USPSTF does not consider the costs of providing a service in this assessment.

The USPSTF recognizes that clinical decisions involve more considerations than evidence alone. Clinicians should understand the evidence but individualize decision making to the specific patient or situation. Similarly, the USPSTF notes that policy and coverage decisions involve considerations in addition to the evidence of clinical benefits and harms.

Summary of Recommendation and Evidence

The USPSTF recommends early screening for syphilis infection in all pregnant women (A recommendation) (Figure 1).

Rationale

Importance

Syphilis is an infection that is primarily sexually transmitted. Untreated syphilis infection in pregnant women can also be transmitted to the fetus (congenital syphilis) at any time during pregnancy or at birth. Congenital syphilis is associated with stillbirth,

Figure 1. USPSTF Grades and Levels of Evidence

What the USPSTF Grades Mean and Suggestions for Practice		
Grade	Definition	Suggestions for Practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
B	The USPSTF recommends the service. There is high certainty that the net benefit is moderate, or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
C	The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.	Offer or provide this service for selected patients depending on individual circumstances.
D	The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.
I statement	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	Read the Clinical Considerations section of the USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

USPSTF Levels of Certainty Regarding Net Benefit	
Level of Certainty	Description
High	The available evidence usually includes consistent results from well-designed, well-conducted studies in representative primary care populations. These studies assess the effects of the preventive service on health outcomes. This conclusion is therefore unlikely to be strongly affected by the results of future studies.
Moderate	The available evidence is sufficient to determine the effects of the preventive service on health outcomes, but confidence in the estimate is constrained by such factors as the number, size, or quality of individual studies. inconsistency of findings across individual studies. limited generalizability of findings to routine primary care practice. lack of coherence in the chain of evidence. As more information becomes available, the magnitude or direction of the observed effect could change, and this change may be large enough to alter the conclusion.
Low	The available evidence is insufficient to assess effects on health outcomes. Evidence is insufficient because of the limited number or size of studies. important flaws in study design or methods. inconsistency of findings across individual studies. gaps in the chain of evidence. findings not generalizable to routine primary care practice. lack of information on important health outcomes. More information may allow estimation of effects on health outcomes.
The USPSTF defines certainty as “likelihood that the USPSTF assessment of the net benefit of a preventive service is correct.” The net benefit is defined as benefit minus harm of the preventive service as implemented in a general, primary care population. The USPSTF assigns a certainty level based on the nature of the overall evidence available to assess the net benefit of a preventive service.	

USPSTF indicates US Preventive Services Task Force.

neonatal death, and significant morbidity in infants (eg, bone deformities and neurologic impairment).¹ After a steady decline from 2008 to 2012, cases of congenital syphilis markedly increased from 2012 to 2016, from 8.4 to 15.7 cases per 100 000 live births (an increase of 87%).² At the same time, national rates of syphilis increased among women of reproductive age.

Reaffirmation

In 2009, the USPSTF reviewed the evidence on screening for syphilis infection in pregnant women and issued an A recommendation.³ The USPSTF has decided to use a reaffirmation deliberation process

to update this recommendation. The USPSTF uses the reaffirmation process for well-established, evidence-based standards of practice in current primary care practice for which only a very high level of evidence would justify a change in the grade of the recommendation.⁴ In its deliberation of the evidence, the USPSTF considers whether the new evidence is of sufficient strength and quality to change its previous conclusions about the evidence.

Detection

The USPSTF found adequate evidence that screening tests can accurately detect syphilis infection in pregnant women.

Figure 2. Clinical Summary: Screening for Syphilis Infection in Pregnant Women

Population	Pregnant women
Recommendation	Screen early for syphilis infection in all pregnant women. Grade: A
Risk Assessment	All pregnant women are at risk. All pregnant women should be tested for syphilis as early as possible when they first present to care. If a woman has not received prenatal care before delivery, she should be tested at the time she presents for delivery.
Screening Tests	Screening for syphilis infection is a 2-step process. The traditional approach is to perform an initial “nontreponemal” antibody test (ie, Venereal Disease Research Laboratory [VDRL] or rapid plasma reagin [RPR] test), followed by a confirmatory “treponemal” antibody detection test (ie, fluorescent treponemal antibody absorption or <i>Treponema pallidum</i> particle agglutination test). A newer alternative is the reverse sequence screening algorithm: an automated treponemal antibody test (eg, enzyme-linked, chemiluminescence, or multiplex flow immunoassay) is performed first, followed by a nontreponemal VDRL or RPR test. If the test results are discordant, a second treponemal test is performed.
Treatments and Interventions	The Centers for Disease Control and Prevention recommend parenteral benzathine penicillin G for the treatment of syphilis in pregnant women.
Relevant USPSTF Recommendations	The USPSTF has made recommendations on screening for other sexually transmitted infections, including chlamydia and gonorrhea, hepatitis B virus, genital herpes, and HIV.

For a summary of the evidence systematically reviewed in making this recommendation, the full recommendation statement, and supporting documents, please go to <https://www.uspreventiveservicestaskforce.org>.



USPSTF indicates US Preventive Services Task Force.

Benefits of Detection and Early Treatment

The USPSTF found convincing evidence that early universal screening for syphilis infection in pregnant women reduces the incidence of congenital syphilis and the adverse outcomes of pregnancy associated with maternal infection.

Harms of Detection and Early Treatment

Screening for syphilis infection in pregnant women may result in potential harms, including false-positive results that require clinical evaluation, anxiety, and harms of treatment with antibiotic medications. However, the USPSTF concluded that these harms of screening are no greater than small.

USPSTF Assessment

Using a reaffirmation process,⁴ the USPSTF concludes with high certainty that the net benefit of screening for syphilis infection in pregnant women is substantial.

Clinical Considerations

Patient Population Under Consideration

This recommendation applies to all pregnant women (Figure 2).

Screening Intervals

All pregnant women should be tested for syphilis as early as possible when they first present to care. If a woman has not received

prenatal care prior to delivery, she should be tested at the time she presents for delivery. In most cases of congenital syphilis, pregnant women received prenatal care but were not screened and treated for syphilis early enough during the pregnancy to prevent transmission to the fetus.

The USPSTF found no new studies that examined the effectiveness of repeated testing for syphilis during pregnancy. The Centers for Disease Control and Prevention (CDC)⁵ and joint guidelines from the American Academy of Pediatrics (AAP) and the American College of Obstetricians and Gynecologists (ACOG)⁶ endorse repeat screening. Specifically, these groups recommend that women at high risk for syphilis be rescreened early in the third trimester (at approximately 28 weeks of gestation) and again at delivery. Women at high risk for syphilis infection include those living in communities or geographic areas with higher prevalence of syphilis, those living with HIV, and those with a history of incarceration or commercial sex work.⁷ AAP and ACOG also recommend repeat screening after exposure to an infected partner.⁶ Clinicians should be aware of the prevalence of syphilis infection in the communities they serve.⁷ Most states mandate screening for syphilis in all pregnant women at the first prenatal visit, and some mandate screening at the time of delivery.⁸

Screening Tests

Syphilis infection is caused by the *Treponema pallidum* bacteria. Current screening tests for syphilis rely on detection of antibodies to the infection rather than direct detection of the bacteria. Screening for syphilis infection is a 2-step process. Traditionally,

screening involved an initial “nontreponemal” antibody test (ie, Venereal Disease Research Laboratory test or rapid plasma reagin [RPR] test) to detect biomarkers released from damage caused by syphilis infection, followed by a confirmatory “treponemal” antibody detection test (ie, fluorescent treponemal antibody absorption or *T pallidum* particle agglutination test). Because nontreponemal tests are complex, a reverse sequence screening algorithm has been developed in which an automated treponemal test (such as an enzyme-linked, chemiluminescence, or multiplex flow immunoassay) is performed first, followed by a nontreponemal test. If the test results of the reverse sequence algorithm are discordant, a second treponemal test (preferably using a different treponemal antibody) is performed. The USPSTF found no studies comparing the false-positive rate of the traditional screening algorithm with that of the reverse sequence screening algorithm among pregnant women. The CDC has provided more detailed guidance on testing for and treatment of sexually transmitted diseases, including syphilis.⁹

Treatment

In 2015, the CDC recommended parenteral benzathine penicillin G for the treatment of syphilis in pregnant women.⁵ Evidence on the efficacy or safety of alternative antibiotic medications for pregnant women and the fetus is very limited; therefore, women who report a penicillin allergy should be evaluated and, if found allergic, desensitized and treated with penicillin. Because the CDC updates its recommendations regularly, clinicians are encouraged to consult the CDC website for the most up-to-date information.⁹

Additional Approaches to Prevention

Trends in congenital syphilis incidence rates are closely related to trends in primary and secondary syphilis infection rates among all women. Screening for syphilis in nonpregnant populations is an important public health approach to preventing the sexual transmission of syphilis and subsequent vertical transmission of congenital syphilis. The USPSTF recommends screening for syphilis in nonpregnant adolescents and adults at increased risk for infection.¹⁰

Useful Resources

The USPSTF has made recommendations on screening for other sexually transmitted infections, including chlamydia and gonorrhea,¹¹ hepatitis B virus,¹² genital herpes,¹³ and HIV.¹⁴ National-, state-, and county-level data on syphilis infection rates are also available from the CDC.²

Other Considerations

Research Needs and Gaps

Although the benefits of screening for syphilis infection in pregnant women to prevent congenital syphilis are well established, additional studies on the use of different screening algorithms in pregnant women, as well as studies to help identify optimal rescreening intervals and populations to rescreen during pregnancy, could help inform implementation of screening

programs. Studies on treatment options besides penicillin could also be helpful.

Discussion

Burden of Disease

Although national rates of syphilis infection among pregnant women are not currently available, the incidence rates of primary and secondary syphilis infection among women and congenital syphilis among infants have been increasing, despite consistent recommendations and legal mandates to screen for syphilis in pregnant women.² In 2012, there were 0.9 cases of primary and secondary syphilis infection per 100 000 women and 8.4 cases of congenital syphilis per 100 000 live births. In 2016, the case rate had increased to 1.9 cases of primary and secondary syphilis infection per 100 000 women and 15.7 cases of congenital syphilis per 100 000 live births.¹⁵

Late or limited prenatal care has been associated with congenital syphilis.² Although nearly 70% of infants with congenital syphilis are born to mothers who received prenatal care, detection and treatment of maternal syphilis often occurs too late to treat the fetus and prevent congenital syphilis.¹⁵ Recent data suggest that while screening rates for syphilis infection are generally high, the proportion of women screened earlier in pregnancy remains low (eg, 20% of women are screened only at the time of delivery).

Primary, secondary, and congenital syphilis rates differ by race/ethnicity. Case rates of primary, secondary, and congenital syphilis are higher in black, American Indian/Alaska Native, and Hispanic populations than in white populations.² Syphilis rates also differ by geography, with generally higher rates of primary, secondary, and congenital syphilis in the Western and Southern states and lower rates in the Northeastern and Midwest states. However, clinicians should be aware of the prevalence of syphilis infection in their community, as rates can vary.²

Syphilis can be transmitted to the fetus during all stages of maternal infection, although the risk is highest with primary and secondary maternal syphilis infection, which is why detection early in pregnancy is important.^{5,16,17} Untreated syphilis infection during pregnancy greatly increases the risk of adverse pregnancy outcomes. A 2013 systematic review of 6 case-control studies found that compared with pregnancies that did not have maternal syphilis infection, untreated maternal syphilis infection during pregnancy was associated with an absolute difference of 21% for stillbirth or fetal loss, 9% for neonatal death, and 5% for prematurity or low birth weight.¹⁸ Although infants born with congenital syphilis are often asymptomatic at birth, some may develop signs within the first several weeks of life, including rash, hemorrhagic rhinitis, lymphadenopathy, hepatosplenomegaly, and skeletal abnormalities.¹⁹ Additional sequelae include anemia, neurologic impairment such as blindness or deafness, and meningitis.

Scope of Review

To reaffirm its 2009 recommendation on screening for syphilis in pregnant women, the USPSTF commissioned a reaffirmation evidence update. The aim of this update is to identify substantial new evidence that is sufficient enough to change the prior

recommendation. Given the established benefits and practice of screening for syphilis in pregnant women, the USPSTF targeted its evidence review to the direct benefits of screening on the prevention of congenital syphilis morbidity and mortality and the harms of screening for and treatment of syphilis infection in pregnant women.

Benefits of Early Detection and Treatment

The USPSTF found no new evidence inconsistent with the benefits of screening for syphilis infection in pregnant women. Evidence from observational studies demonstrates fewer adverse pregnancy outcomes among pregnant women screened and treated for syphilis infection compared with pregnant women not treated. In particular, treatment appears to be more beneficial when provided earlier rather than later in pregnancy.¹⁶ A 2014 systematic review of 54 observational studies found that incidence of congenital syphilis, preterm birth, low birth weight, stillbirth, and neonatal death was dramatically reduced in women treated for syphilis during pregnancy compared with women who had untreated syphilis.²⁰ However, only a slight reduction in stillbirth or fetal loss was observed in women who were only treated in the third trimester, compared with women who had untreated syphilis. Extended follow-up of a study previously considered by the USPSTF for its 2009 recommendation³ reported on the effects of implementing a free syphilis screening and treatment program for all pregnant women living in Shenzhen, China, from 2002 to 2012 (n = 2 441 237).²¹ During follow-up, screening uptake increased from 89.8% to 97.2%, and the congenital syphilis case rate decreased from 109.3 to 9.4 cases per 100 000 live births. During the same time, the incidence of adverse pregnancy outcomes decreased from 42.7% to 19.2%, and the incidence of stillbirth or fetal loss decreased from 19.0% to 3.3%. The USPSTF found this evidence to be consistent with findings from its previous evidence review.²²

Potential Harms of Screening and Treatment

Potential harms of screening for and treatment of syphilis infection include false-positive results that require clinical evaluation, unnecessary anxiety to the patient, and harms of antibiotic medication use.¹⁶ Consistent with previous reviews, the current, targeted review identified 5 studies that reported on false-positive rates of various screening tests²³⁻²⁷ and 1 study that reported on false-negative RPR test results.²⁸ Overall, these studies demonstrate that false-positive results with chemiluminescence or enzyme-linked immunoassay in pregnant women are common (false-positive rates ranged from 0% to 88.2%)²³⁻²⁷ and that false-negative RPR test results can result from undiluted serum with high titers (known as the prozone effect).²⁸ Because of the high potential for false-positive results with individual tests, a 2-step screening algorithm is recommended. The USPSTF found no studies that reported on the false-positive rate of the traditional or reverse sequence screening algorithm in pregnant women. Harms of treatment include rare adverse drug-related effects, such as anaphylaxis attributable to penicillin allergy and the Jarisch-Herxheimer reaction (febrile reaction with headache, myalgia, and other symptoms), which may occur within the first 24 hours after any type of syphilis therapy. The USPSTF found no studies that reported on harms to the fetus from treatment of syphilis infection.

Estimate of Magnitude of Net Benefit

The USPSTF considered the evidence using a reaffirmation process and found that accurate screening algorithms are available to identify syphilis infection. Effective treatment with antibiotics can prevent congenital syphilis and significantly decrease adverse pregnancy outcomes, with small associated harms, providing an overall substantial health benefit. Therefore, the USPSTF reaffirms its previous conclusion that there is convincing evidence that screening for syphilis infection in pregnant women provides substantial net benefit.

Response to Public Comment

A draft version of this recommendation statement was posted for public comment on the USPSTF website from February 6, 2018, to March 5, 2018. Most comments supported the recommendation statement. Several comments requested clearer guidance about the timing of initial syphilis screening (as early as possible after the first positive pregnancy test result). Other comments requested clarification on risk factors that would warrant repeat testing. The USPSTF provided more information in the recommendation statement to address these concerns. Several comments requested changes that are outside the scope of the USPSTF or not consistent with its methods and processes, and so no changes were made in response. These suggested changes included endorsement of guidelines from other organizations and discussion of evidence not considered in the USPSTF's deliberation on this recommendation. Some comments suggested additional citations; these were reviewed by the USPSTF but none met inclusion criteria.

Reaffirmation of Previous USPSTF Recommendation

This recommendation is a reaffirmation of the USPSTF 2009 recommendation statement.³ In 2009, the USPSTF reviewed the evidence on screening for syphilis infection in pregnant women and found that the benefits of screening substantially outweighed the harms.²² For the current recommendation, the USPSTF commissioned a targeted review¹⁶ to look for substantial new evidence on the benefits and harms of screening and determined that the net benefit of screening for syphilis infection in pregnant women continues to be well established. The USPSTF found no new substantial evidence that could change its recommendation and, therefore, reaffirms its recommendation to screen for syphilis infection in all pregnant women.

Recommendations of Others

This recommendation statement is consistent with those of other professional and public health organizations. The CDC recommends screening for syphilis infection in all pregnant women at their first prenatal visit.⁵ Joint guidelines from AAP and ACOG recommend screening for syphilis infection in pregnant women as early as possible in pregnancy.⁶ The CDC, AAP, and ACOG also recommend repeat screening at 28 weeks of gestation and again at delivery in high-risk women. Women at high risk for syphilis infection

include those living in high-prevalence communities, those living with HIV, and those with a history of incarceration or commercial sex work.¹⁰ AAP and ACOG also recommend repeat screening after ex-

posure to an infected partner.⁶ The American Academy of Family Physicians recommends screening for syphilis infection in all pregnant women.²⁹

ARTICLE INFORMATION

Accepted for Publication: July 25, 2018.

The US Preventive Services Task Force (USPSTF) members: Susan J. Curry, PhD; Alex H. Krist, MD, MPH; Douglas K. Owens, MD, MS; Michael J. Barry, MD; Aaron B. Caughey, MD, PhD; Karina W. Davidson, PhD, MASC; Chyke A. Doubeni, MD, MPH; John W. Epling Jr, MD, MSED; Alex R. Kemper, MD, MPH, MS; Martha Kubik, PhD, RN; Ann E. Kurth, PhD, RN, MSN, MPH; C. Seth Landefeld, MD; Carol M. Mangione, MD, MSPH; Maureen G. Phipps, MD, MPH; Michael Silverstein, MD, MPH; Melissa A. Simon, MD, MPH; Chien-Wen Tseng, MD, MPH, MSEE; John B. Wong, MD.

Affiliations of The US Preventive Services Task Force (USPSTF) members: University of Iowa, Iowa City (Curry); Fairfax Family Practice Residency, Fairfax, Virginia (Krist); Virginia Commonwealth University, Richmond (Krist); Veterans Affairs Palo Alto Health Care System, Palo Alto, California (Owens); Stanford University, Stanford, California (Owens); Harvard Medical School, Boston, Massachusetts (Barry); Oregon Health & Science University, Portland (Caughey); Columbia University, New York, New York (Davidson); University of Pennsylvania, Philadelphia (Doubeni); Virginia Tech Carilion School of Medicine, Roanoke (Epling); Nationwide Children's Hospital, Columbus, Ohio (Kemper); Temple University, Philadelphia, Pennsylvania (Kubik); Yale University, New Haven, Connecticut (Kurth); University of Alabama at Birmingham (Landefeld); University of California, Los Angeles (Mangione); Brown University, Providence, Rhode Island (Phipps); Boston University, Boston, Massachusetts (Silverstein); Northwestern University, Evanston, Illinois (Simon); University of Hawaii, Honolulu (Tseng); Pacific Health Research and Education Institute, Honolulu, Hawaii (Tseng); Tufts University, Medford, Massachusetts (Wong).

Author Contributions: Dr Curry had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. The USPSTF members contributed equally to the recommendation statement.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Authors followed the policy regarding conflicts of interest described at <https://www.uspreventiveservicestaskforce.org/Page/Name/conflict-of-interest-disclosures>. All members of the USPSTF receive travel reimbursement and an honorarium for participating in USPSTF meetings. No other disclosures were reported.

Funding/Support: The USPSTF is an independent, voluntary body. The US Congress mandates that the Agency for Healthcare Research and Quality (AHRQ) support the operations of the USPSTF.

Role of the Funder/Sponsor: AHRQ staff assisted in the following: development and review of the research plan, commission of the systematic evidence review from an Evidence-based Practice Center, coordination of expert review and public comment of the draft evidence report and draft

recommendation statement, and the writing and preparation of the final recommendation statement and its submission for publication. AHRQ staff had no role in the approval of the final recommendation statement or the decision to submit for publication.

Disclaimer: Recommendations made by the USPSTF are independent of the US government. They should not be construed as an official position of AHRQ or the US Department of Health and Human Services.

Additional Contributions: We thank Jill Huppert, MD, MPH (AHRQ), and Tina Fan, MD, MPH (AHRQ), who contributed to the writing of the manuscript, and Lisa Nicoletta, MA (AHRQ), who assisted with coordination and editing.

REFERENCES

- Cooper JM, Sánchez PJ. Congenital syphilis. *Semin Perinatol*. 2018;42(3):176-184. doi:10.1053/j.semperi.2018.02.005
- Centers for Disease Control and Prevention. *Sexually Transmitted Disease Surveillance, 2016*. Atlanta, GA: Centers for Disease Control and Prevention; 2017.
- U.S. Preventive Services Task Force. Screening for syphilis infection in pregnancy: U.S. Preventive Services Task Force reaffirmation recommendation statement. *Ann Intern Med*. 2009;150(10):705-709. doi:10.7326/0003-4819-150-10-200905190-00008
- US Preventive Services Task Force (USPSTF). Procedure Manual. USPSTF website. <https://www.uspreventiveservicestaskforce.org/Page/Name/procedure-manual>. 2018. Accessed July 17, 2018.
- Centers for Disease Control and Prevention. *Sexually Transmitted Diseases Treatment Guidelines, 2015*. Atlanta, GA: Centers for Disease Control and Prevention; 2015.
- American Academy of Pediatrics; American College of Obstetricians and Gynecologists. *Guidelines for Perinatal Care*. 8th ed. Elk Grove Village, IL: American Academy of Pediatrics; American College of Obstetricians and Gynecologists; 2017.
- Cantor AG, Pappas M, Daeges M, Nelson HD. Screening for syphilis: updated evidence report and systematic review for the US Preventive Services Task Force. *JAMA*. 2016;315(21):2328-2337. doi:10.1001/jama.2016.4114
- Warren HP, Cramer R, Kidd S, Leichter JS. State requirements for prenatal syphilis screening in the United States, 2016 [published online July 17, 2018]. *Matern Child Health J*. 2018. doi:10.1007/s10995-018-2592-0
- Centers for Disease Control and Prevention (CDC). Sexually transmitted diseases (STDs). CDC website. <https://www.cdc.gov/std/treatment/>. Updated November 30, 2017. Accessed July 17, 2018.
- Bibbins-Domingo K, Grossman DC, Curry SJ, et al; US Preventive Services Task Force (USPSTF). Screening for syphilis infection in nonpregnant adults and adolescents: US Preventive Services Task Force recommendation statement. *JAMA*. 2016;315(21):2321-2327. doi:10.1001/jama.2016.5824
- LeFevre ML; U.S. Preventive Services Task Force. Screening for chlamydia and gonorrhea: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2014;161(12):902-910. doi:10.7326/M14-1981
- LeFevre ML; U.S. Preventive Services Task Force. Screening for hepatitis B virus infection in nonpregnant adolescents and adults: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2014;161(1):58-66. doi:10.7326/M14-1018
- Bibbins-Domingo K, Grossman DC, Curry SJ, et al; US Preventive Services Task Force. Serologic screening for genital herpes infection: US Preventive Services Task Force recommendation statement. *JAMA*. 2016;316(23):2525-2530. doi:10.1001/jama.2016.16776
- Moyer VA; U.S. Preventive Services Task Force. Screening for HIV: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;159(1):51-60. doi:10.7326/0003-4819-159-1-201307020-00645
- Bowen V, Su J, Torrone E, Kidd S, Weinstock H. Increase in incidence of congenital syphilis—United States, 2012-2014. *MMWR Morb Mortal Wkly Rep*. 2015;64(44):1241-1245. doi:10.15585/mmwr.mm6444a3
- Lin JS, Eder M, Bean S. *Screening for Syphilis Infection in Pregnant Women: A Reaffirmation Evidence Update for the U.S. Preventive Services Task Force: Evidence Synthesis No. 167*. Rockville, MD: Agency for Healthcare Research and Quality; 2018. AHRQ publication 18-05238-EF-1.
- Lin JS, Eder M, Bean SI. Screening for syphilis infection in pregnant women: updated evidence report and systematic review for the US Preventive Services Task Force [published September 4, 2018]. *JAMA*. doi:10.1001/jama.2018.7769
- Gomez GB, Kamb ML, Newman LM, Mark J, Broutet N, Hawkes SJ. Untreated maternal syphilis and adverse outcomes of pregnancy: a systematic review and meta-analysis. *Bull World Health Organ*. 2013;91(3):217-226. doi:10.2471/BLT.12.107623
- Kingston M, French P, Higgins S, et al; Members of the Syphilis Guidelines Revision Group 2015. UK national guidelines on the management of syphilis 2015. *Int J STD AIDS*. 2016;27(6):421-446. doi:10.1177/0956462415624059
- Qin J, Yang T, Xiao S, Tan H, Feng T, Fu H. Reported estimates of adverse pregnancy outcomes among women with and without syphilis: a systematic review and meta-analysis. *PLoS One*. 2014;9(7):e102203. doi:10.1371/journal.pone.0102203
- Qin JB, Feng TJ, Yang TB, et al. Synthesized prevention and control of one decade for mother-to-child transmission of syphilis and determinants associated with congenital syphilis and adverse pregnancy outcomes in Shenzhen, South China. *Eur J Clin Microbiol Infect Dis*. 2014;33(12):2183-2198. doi:10.1007/s10096-014-2186-8

22. Wolff T, Shelton E, Sessions C, Miller T. Screening for syphilis infection in pregnant women: evidence for the U.S. Preventive Services Task Force reaffirmation recommendation statement. *Ann Intern Med*. 2009;150(10):710-716. doi:10.7326/0003-4819-150-10-200905190-00009
23. Boonchaoy A, Wongchampa P, Hirankarn N, Chaithongwongwatthana S. Performance of chemiluminescent microparticle immunoassay in screening for syphilis in pregnant women from low-prevalence, resource-limited setting. *J Med Assoc Thai*. 2016;99(2):119-124.
24. Henrich TJ, Yawetz S. Impact of age, gender, and pregnancy on syphilis screening using the Captia Syphilis-G assay. *Sex Transm Dis*. 2011;38(12):1126-1130. doi:10.1097/OLQ.0b013e31822e60e1
25. Mmeje O, Chow JM, Davidson L, Shieh J, Schapiro JM, Park IU. Discordant syphilis immunoassays in pregnancy: perinatal outcomes and implications for clinical management. *Clin Infect Dis*. 2015;61(7):1049-1053. doi:10.1093/cid/civ445
26. Wang KD, Xu DJ, Su JR. Preferable procedure for the screening of syphilis in clinical laboratories in China. *Infect Dis (Lond)*. 2016;48(1):26-31. doi:10.3109/23744235.2015.1044465
27. Wellinghausen N, Dietenberger H. Evaluation of two automated chemiluminescence immunoassays, the LIAISON Treponema Screen and the ARCHITECT Syphilis TP, and the *Treponema pallidum* particle agglutination test for laboratory diagnosis of syphilis. *Clin Chem Lab Med*. 2011;49(8):1375-1377. doi:10.1515/CCLM.2011.643
28. Liu LL, Lin LR, Tong ML, et al. Incidence and risk factors for the prozone phenomenon in serologic testing for syphilis in a large cohort. *Clin Infect Dis*. 2014;59(3):384-389. doi:10.1093/cid/ciu325
29. American Academy of Family Physicians (AAFP). Clinical Preventive Service Recommendation: syphilis. AAFP website. <https://www.aafp.org/patient-care/clinical-recommendations/all/syphilis.html>. 2018. Accessed July 17, 2018.